

113-2 General Physics II

李傳睿

Chuan-Jui Li



國立政治大學應用物理研究所

Graduate Institute of Applied Physics, NCCU

CJLI

About me



李傳睿 Chuan-Jui Li (CJLI)

✉ cjli@nccu.edu.tw

💻 <https://sites.google.com/view/cjli>

Research Area: Observational Astrophysics and Data analysis
(supernovae and their interaction with the interstellar medium)

2004 – 2008 B.S. in Transportation Technology & Management,
National Chiao Tung University

2008 – 2010 M.S. in Physics
National Chiao Tung University

2010 – 2018 Ph.D. in Astrophysics
National Taiwan University

2018 – 2021 ASIAA Postdoctoral Fellow (RDSS)

2022 – 2024 AS Postdoctoral Fellow


2024 NCCU Assistant Professor

This Lecture

- About Me
 - My Research
- About “General Physics II”
 - Fascinating Electricity & Magnetism


Electric fields

- Electric charges
- Conductors and Insulators
- Coulomb's Law
- Electric fields



References

- **R. A. Serway and J. W. Jewett Jr.,
Physics for Scientists and Engineers**
 - Part 4 Electricity and Magnetism
 - Part 5 Light and Optics
- R. Feynman, R. Leighton, and M. Sands,
The Feynman Lectures on Physics
<https://www.feynmanlectures.caltech.edu/>



Syllabus

Week 01	(02/17)	Electric fields	Part 4 Electricity and Magnetism
Week 02	(02/24)	Gauss's law	
Week 03	(03/03)	Electric potential	
Week 04	(03/10)	Capacitance and dielectrics	
Week 05	(03/17)	Current and resistance	
Week 06	(03/24)	Direct current circuits	
Week 07	(03/31)	Magnetic fields	
Week 08	(04/07)	Faraday's law	
Week 09	(04/14)	Midterm	
Week 10	(04/21)	Inductance	
Week 11	(04/28)	Alternating current circuits	
Week 12	(05/05)	Electromagnetic waves	
Week 13	(05/12)	The nature of light and the laws of geometric optics	
Week 14	(05/19)	Image formation	Part 5 Light and Optics
Week 15	(05/26)	Interference of light waves	
Week 16	(06/02)	Final Exam	
Week 17–18	(06/09, 06/16)	Capstone Self-Learning	

Gradings and Scores

- Homework (30%)
 - Assigned every 1 or 2 weeks (on Moodle)
 - Due 1 week (before the next class, printed or online submission)
 - One-week submission: 100% credit.
 - Afterwards: 70% credit.
- Midterm (30%)
- Final Exam (30%)
- Class participation (10%)

Office Hours

- 教師: 李傳睿 Chuan-Jui Li (CJLI)

✉ cjli@nccu.edu.tw

💻 <https://sites.google.com/view/cjli>



- 助教: 楊斯皓

113-2 哲學系大四 (雙主修電物)

114-1 應物所

✉ 110104035@nccu.edu.tw



- 請寄信: 1. 查閱郵件禮儀 2. 註明系級姓名
 3. 簡述問題 4. 預約時間
-

- 一起維持鼓勵學習的環境。


如果同學比較內向，歡迎寄信表達，合理都會聆聽。

- 信用比分數珍貴，請說真話。


Fascinating Electricity & Magnetism



Hydrogen bonding in water



Action potential (nerve impulse)



Fascinating Electricity & Magnetism

Any sufficiently advanced technology is indistinguishable from magic – C. Clarke

Smartphone

<https://en.wikipedia.org/wiki/Smartphone>

Computer

<https://en.wikipedia.org/wiki/Computer>

Electric motors

https://en.wikipedia.org/wiki/Electric_motor

Modern all-electric cars

Tesla Model 3

Nissan Leaf

Hyundai Ioniq 5

BMW i3

Television

<https://en.wikipedia.org/wiki/Smartphone>

Electronics






<https://en.wikipedia.org/wiki/Electronics>

High-energy accelerator


https://en.wikipedia.org/wiki/Large_Hadron_Collider

Fascinating Electricity & Magnetism

Any sufficiently advanced technology is indistinguishable from magic – C. Clarke



Fascinating Electricity & Magnetism




X-Men Magneto 萬磁王

https://en.wikipedia.org/wiki/X-Men:_Days_of_Future_Past
https://www.youtube.com/watch?v=pAApe_5Rewg




Fascinating Electricity & Magnetism

Gravity



Electric forces



https://en.wikipedia.org/wiki/Intermolecular_force <https://en.wikipedia.org/wiki/Biology> <https://en.wikipedia.org/wiki/Chemistry>

Biology



Chemistry




This Lecture

- About Me
 - My Research
- About “General Physics II”
 - Fascinating Electricity & Magnetism

Electric fields

- Electric charges
- Conductors and Insulators
- Coulomb's Law
- Electric fields



Lecture 01 – chapter 23

Electric Fields

蔡尚岳
政治大學應用物理所

Lecture 01 – chapter 23

Electric Fields

蔡尚岳

政治大學應用物理所



為了保留蔡老師的原創，以及不影響同學閱讀，
若左下角標註 CJLI，代表內容已作更動或新增。

Some History

Not until the early 19th century did scientists establish
electricity and magnetism as related phenomena.

- 1785
 - Charles Coulomb confirmed inverse square law form for electric forces


In the upcoming slides




Some History

Not until the early 19th century did scientists establish
electricity and magnetism as related phenomena.

- 1785
 - Charles Coulomb confirmed inverse square law form for electric forces
- 1819
 - Hans Oersted found a **compass needle** deflected when near a wire carrying an electric current



Hans Christian Ørsted




https://en.wikipedia.org/wiki/Hans_Christian_%C3%98rsted

Some History

Not until the early 19th century did scientists establish electricity and magnetism as related phenomena.

- 1785
 - Charles-Augustin de Coulomb found the inverse square law form for electric force
- 1819
 - Hans Christian Oersted showed that a magnetic needle deflected when near a wire carrying an electric current
- 1831
 - Michael Faraday and Joseph Henry showed that when a **wire is moved near a magnet**, an electric current is produced in the wire

Faraday's 1831 demonstration



A change in the magnetic flux of the left coil induces a current in the right coil.

Later this semester

https://en.wikipedia.org/wiki/Joseph_Henry

https://en.wikipedia.org/wiki/Michael_Faraday


https://en.wikipedia.org/wiki/Faraday%27s_law_of_induction

Some History

Not until the early 19th century did scientists establish electricity and magnetism as related phenomena.

- 1785
 - Charles-Augustin de Coulomb found the law of electrostatics
- 1819
 - Hans Christian Ørsted caused a compass needle to move when it was placed near a wire carrying electric current
- 1831
 - Michael Faraday and Joseph Henry **moved near a magnet**, an electric current was induced in the wire


Faraday's 1831 demonstration



https://en.wikipedia.org/wiki/Joseph_Henry

https://en.wikipedia.org/wiki/Michael_Faraday

https://en.wikipedia.org/wiki/Faraday%27s_law_of_induction





Later this semester

Some History

Not until the early 19th century did scientists establish
electricity and magnetism as related phenomena.

- 1785
 - Charles Coulomb confirmed the inverse square law for electric forces
- 1819
 - Hans Christian Oersted carried out experiments showing that a current in a wire creates a magnetic field
- 1831
 - Michael Faraday observed that a changing magnetic field induces an electromotive force in a wire
- 1873
 - James Clerk Maxwell used observations and other experimental facts as a basis for formulating the **laws of electromagnetism**
 - Unified electricity and magnetism



for electric forces
when near a wire
when a **wire is** moved in the wire

https://en.wikipedia.org/wiki/Electromagnetic_radiation
https://en.wikipedia.org/wiki/James_Clerk_Maxwell

Some History

Not until the early 19th century did scientists establish electricity and magnetism as related phenomena.

- 1785

- Char


愛因斯坦

1905 狹義相對論
1915 廣義相對論

114-1 Modern Physics

- 1831

- Michael Faraday



- 1873

- James Clerk Maxwell used observations and other facts as a basis for formulating the **laws of electromagnetism**
 - Unified electricity and magnetism

https://en.wikipedia.org/wiki/Electromagnetic_radiation
https://en.wikipedia.org/wiki/James_Clerk_Maxwell



Movie *Interstellar* 2014

Some History

- 1785
 - Charles Coulomb confirmed inverse square law form for electric forces
- 1819
 - Hans Oersted found a **compass needle** deflected when near a wire carrying an electric current
- 1831
 - Michael Faraday and Joseph Henry showed that when a **wire is moved near a magnet**, an electric current is produced in the wire
- 1873
 - James Clerk Maxwell used observations and other experimental facts as a basis for formulating the **laws of electromagnetism**
 - Unified electricity and magnetism

Electric Charges (電荷)

- Two kinds of electric charges
 - **Negative charges** : electrons
 - **Positive charges** : protons
 - Repel (same sign) or Attract (opposite signs)
↖ Charge conservation
- Electric charge is **conserved** in an isolated system
 - **Electrification** : transfer of charge from one object to another
 - Example : A glass rod is rubbed with silk
 - Electrons are transferred from the glass to the silk

Quantization of Electric Charges

- The electric charge, q , is said to be **quantized**

- $q = \pm Ne$ (Robert A. Millikan)

- N is an integer
- e is the fundamental **unit of charge**, $|e| = 1.6 \times 10^{-19}$ C
- The SI unit of charge is the **coulomb (C)**


Table 23.1 Charge and Mass of the Electron, Proton, and Neutron

Particle	Charge (C)	Mass (kg)
Electron (e)	$-1.602\ 176\ 5 \times 10^{-19}$ (-e)	$9.109\ 4 \times 10^{-31}$
Proton (p)	$+1.602\ 176\ 5 \times 10^{-19}$ (+e)	$1.672\ 62 \times 10^{-27}$
Neutron (n)	0	$1.674\ 93 \times 10^{-27}$

Quick Quiz 22.1


- Three objects are brought close to each other, two at a time. When objects A and B are brought together, they repel. When objects B and C are brought together, they also repel. Which of the following are true? (Multiple choices)
 - (a) Objects A and C possess charges of the same sign.
 - (b) Objects A and C possess charges of opposite sign.
 - (c) All three objects possess charges of the same sign.
 - (d) One object is neutral
 - (e) Additional experiments must be performed to determine the signs of the charges.

Answer: (a)(c)(e)




Conductors and Insulators

- Electrical conductors (導體)
 - Some of the electrons are **free electrons**
 - Move relatively **freely** through the material
- Electrical insulators (絕緣體)
 - All of the electrons are **bound to atoms**
- Semiconductors (半導體)



Section 23.2




9 Awesome Science Tricks Using Static Electricity!

Experiments demonstrate the existence of the electric force.




<https://www.youtube.com/watch?v=ViZNgU-Yt-Y>

Charles Coulomb




© INTERFOTO/Alamy

- 1736 – 1806
- French physicist
- Major contributions were in areas of electrostatics and magnetism
- Also investigated in areas of
 - Strengths of materials 、 Structural mechanics
- Twist angle → Torque → The force between particles
- Same apparatus used by Cavendish to measure the density of Earth



Charles Coulomb




© INTERFOTO/Alamy

- 1736
- French
- Major contributions in electricity
- Also interested in:
 - Static electricity
 - magnetism
- Twisted the torsion balance between charges
- Same principle used to measure the density of Earth

TutorVista.com

Coulomb's Torsion Balance



The diagram illustrates Coulomb's Torsion Balance. A hand holds a blue sphere at the top of a vertical glass tube. The tube is supported by a torsion head at the top and a lid at the bottom. The lid rests on a cylindrical glass case containing a scale and a small metal sphere. A torsion fibre runs vertically through the tube, connecting the top torsion head to the lid. Labels identify the components: Torsion head, Glass tube, Torsion fibre, Lid, Cylindrical glass case, Scale, and Coulomb's torsion balance. To the right, a side view shows the suspension head, fiber, and fiber holder. Control buttons at the bottom include PAUSE, REPLAY, BACK, and NEXT, along with a volume icon. The edurite logo is in the bottom left corner.

Suspension head

Fiber

A

PAUSE REPLAY BACK NEXT


edurite

Coulomb's Law (庫倫定律)

$$F_e = k_e \frac{|q_1| |q_2|}{r^2}$$

$$k = \frac{1}{4\pi\epsilon_0}$$

- Coulomb constant, $k_e = 8.9876 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2 = 1/(4\pi\epsilon_0)$
- Permittivity (介電常數) of free space, $\epsilon_0 = 8.8542 \times 10^{-12} \text{ C}^2 / \text{N}\cdot\text{m}^2$
- The force is a conservative force (保守力)



In physics, a **conservative force** is a force with the property that the total [work](#) done by the force in moving a particle between two points is independent of the path taken.^[1] Equivalently, if a particle travels in a closed loop, the total work done (the sum of the force acting along the path multiplied by the displacement) by a conservative force is zero.^[2]

A conservative force depends only on the position of the object. If a force is conservative, it is possible to assign a numerical value for the [potential](#) at any point and conversely, when an object moves from one location to another, the force changes the [potential energy](#) of the object by an amount that does not depend on the path taken, contributing to the [mechanical energy](#) and the overall [conservation of energy](#). If the force is not conservative, then defining a scalar potential is not possible, because taking different paths would lead to conflicting potential differences between the start and end points.

Gravitational force is an example of a conservative force, while frictional force is an example of a non-conservative force.

Other examples of conservative forces are: [force in elastic spring](#), [electrostatic force](#) between two electric charges, and [magnetic force](#) between two magnetic poles. The last two forces are called central forces as they act along the line joining the centres of two charged/magnetized bodies. A central force is conservative if and only if it is spherically symmetric.^[3]

For conservative forces,

$$\mathbf{F}_c = -\frac{dU}{ds}$$

where F_c is the conservative force, U is the potential energy, and s is the position.^[4]

Part of a series on
Classical mechanics

$$\mathbf{F} = \frac{d\mathbf{p}}{dt}$$

[Second law of motion](#)

[History](#) · [Timeline](#) · [Textbooks](#)

Branches [\[show\]](#)

Fundamentals [\[show\]](#)

Formulations [\[show\]](#)

Core topics [\[show\]](#)

Rotation [\[show\]](#)


Scientists [\[show\]](#)

[Physics portal](#) [Category](#)

V · T · E


Multiple Charges

- The vector sum of the forces $\vec{F}_1 = \vec{F}_{21} + \vec{F}_{31} + \vec{F}_{41}$
- Example 23.3 : $q_1 = 15.0 \mu C$, $q_2 = 6.00 \mu C$, and the net force acting on q_3 is zero. What is the x coordinate of q_3 ? 0.775



Electric Field (電場)

- Electric field exist in the region of space around a source charge
- The electric field vector, \vec{E} , is defined as the electric force on the test charge per unit charge (Newton/coulomb)




$$\vec{E} = \frac{\vec{F}_e}{q_0} = k_e \frac{q}{r^2} \hat{r}$$

\downarrow test charge

$$\vec{E} = k_e \sum_i \frac{q_i}{r_i^2} \hat{r}_i$$


More About Electric Field Direction



Quick Quiz 23.4


- A test charge of $+3 \mu\text{C}$ is at a point P where an external electric field is directed to the right and has a magnitude of $4 \times 10^6 \text{ N/C}$. If the test charge is replaced with another test charge of $-3 \mu\text{C}$, what happens to the external electric field at P ?
- (a) It is unaffected.
- (b) It reverses direction.
- (c) It changes in a way that cannot be determined.

- **Answer: (a)**



Electric Field Lines

- A pictorial representation of electric field
 - “+” → “-”
 - No cross
 - (Number of lines per unit area through a surface) $\propto E$




Dipole

Like charges

Section 23.6

Unequal charges

Lightning




https://en.wikipedia.org/wiki/Electrostatic_discharge
<https://en.wikipedia.org/wiki/Lightning>

Electron in a Uniform Field

$$\vec{F}_e = q\vec{E} = m\vec{a}$$
$$\vec{a} = \frac{q\vec{E}}{m}$$

The electron undergoes a downward acceleration (opposite \vec{E}), and its motion is parabolic while it is between the plates.

parabola: 抛物線



© Cengage Learning. All Rights Reserved.

Example 23.8: An Accelerated Electron


- An electron enters the region of a uniform electric field as shown in the figure, with $v_i = 3 \times 10^6 \text{ m/s}$ and $E = 200 \text{ N/C}$. The horizontal length of the plates is $\ell = 0.100 \text{ m}$.
- (A) Find the acceleration of the electron while it is in the electric field
- (B) Find the time at which the electron leaves the field.
- (C) What is the electron vertical position when it leaves the field?
(Assuming $y_i(0) = 0$)

$$a_y = -\frac{eE}{m_e}$$

(A)

$$a_y = -\frac{(1.60 \times 10^{-19} \text{ C})(200 \text{ N/C})}{9.11 \times 10^{-31} \text{ kg}}$$
$$= -3.51 \times 10^{13} \text{ m/s}^2$$


The electron undergoes a downward acceleration (opposite \vec{E}), and its motion is parabolic while it is between the plates.



Example 23.8: An Accelerated Electron

(B) $t = \frac{\ell - 0}{v_x} = \frac{0.100 \text{ m}}{3.00 \times 10^6 \text{ m/s}}$
 $= 3.33 \times 10^{-8} \text{ s}$

(c) $y_f = y_i + \left(v_{yi} t + \frac{1}{2} a_y t^2 \right)$




$$y_f = 0 + 0 + \frac{1}{2} \left(-3.51 \times 10^{13} \text{ m/s}^2 \right) \left(3.33 \times 10^{-8} \text{ s} \right)^2$$
$$= -0.0195 \text{ m} = -1.95 \text{ cm}$$

Formulas for Uniformly - Accelerated Motion

$$1. \quad V = U + at$$

$$2. \quad S = Ut + \frac{1}{2}at^2$$

$$3. \quad V^2 = U^2 + 2as$$



Quick quiz

- The diagrams below show two uniformly charged spheres. The charge on the right sphere is 3 times as large as the charge on the left sphere.
- (a) Which **force** diagram best represents the magnitudes and directions of the electric forces on the two spheres? **d**
- ~~(b)~~ Which choice best represents the magnitudes and directions of the **electric field vectors** created by one sphere at the other sphere? **e**



Answer (a): (d)

Answer (b): (e)

This Lecture

- About Me
 - My Research
- About “General Physics II”
 - Fascinating Electricity & Magnetism

Electric fields

- Electric charges
- Conductors and Insulators
- Coulomb's Law
- Electric fields

