

DIABETIC RETINOPATHY ASSESSMENT REPORT

Report Date: May 02, 2025

PATIENT INFORMATION

Image Filename: IDRiD_01.jpg

Patient ID: For clinical use, please enter patient information

ANALYSIS RESULTS

Grade 2: Moderate Non-Proliferative Diabetic Retinopathy


Clinical Description:

More microaneurysms are present along with other signs such as dot and blot hemorrhages, hard exudates (lipid deposits), and cotton wool spots (nerve fiber layer infarcts).

Medical Advice:

Progressive stage detected. Schedule a comprehensive eye examination within 6 months and maintain strict glycemic control.

RETINAL IMAGES

Original Image	AI-Processed Image with Lesions
	

ABOUT DIABETIC RETINOPATHY

What is Diabetic Retinopathy?

Diabetic retinopathy is a diabetes complication that affects the eyes. It's caused by damage to the blood vessels in the retina (the light-sensitive tissue at the back of the eye). Over time, too much sugar in the blood can lead to blockage of the tiny blood vessels that nourish the retina, cutting off its blood supply. As a result, the eye attempts to grow new blood vessels, but these don't develop properly.

Stages of Diabetic Retinopathy:

- 1. No Diabetic Retinopathy (Grade 0):** No abnormalities detected.
- 2. Mild NPDR (Grade 1):** Microaneurysms (small swellings in blood vessels) appear.
- 3. Moderate NPDR (Grade 2):** Some blood vessels that nourish the retina become blocked.
- 4. Severe NPDR (Grade 3):** Many blood vessels are blocked, depriving several areas of the retina of blood supply.
- 5. Proliferative DR (Grade 4):** New, abnormal blood vessels grow (proliferate) in the retina, which can leak and cause severe vision problems.

DISCLAIMER

This report is generated by an automated Diabetic Retinopathy Detection System using artificial intelligence algorithms to analyze retinal images. The results provided are meant to assist healthcare professionals and should NOT be used as the sole basis for medical decisions. This report MUST be reviewed and verified by a qualified ophthalmologist or healthcare professional before any treatment decisions are made. Early detection and treatment of diabetic retinopathy can significantly reduce the risk of vision loss.