

Automatic Differentiation Variational Inference

Philip Schulz and Wilker Aziz

[https:
//github.com/philschulz/VITutorial](https://github.com/philschulz/VITutorial)

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks
- ▶ Objective: a lowerbound on likelihood

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks
- ▶ Objective: a lowerbound on likelihood
- ▶ ELBO cannot be computed exactly, but we can obtain gradient estimates

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks
- ▶ Objective: a lowerbound on likelihood
- ▶ ELBO cannot be computed exactly, but we can obtain gradient estimates
- ▶ Main problem: the MC estimator is not differentiable

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks
- ▶ Objective: a lowerbound on likelihood
- ▶ ELBO cannot be computed exactly, but we can obtain gradient estimates
- ▶ Main problem: the MC estimator is not differentiable
 - ▶ Score function estimator: applicable to any model

What we know so far

- ▶ DGMs: probabilistic models parameterised by neural networks
- ▶ Objective: a lowerbound on likelihood
- ▶ ELBO cannot be computed exactly, but we can obtain gradient estimates
- ▶ Main problem: the MC estimator is not differentiable
 - ▶ Score function estimator: applicable to any model
 - ▶ Reparameterised gradients: so far seems applicable only to Gaussian variables

Reparameterised gradients: Gaussian

We have seen one case, namely,
if $\epsilon \sim \mathcal{N}(0, I)$ and $Z \sim \mathcal{N}(\mu, \sigma^2)$

Reparameterised gradients: Gaussian

We have seen one case, namely,
if $\epsilon \sim \mathcal{N}(0, I)$ and $Z \sim \mathcal{N}(\mu, \sigma^2)$

Then

$$Z \sim \mu + \sigma\epsilon$$

and

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)]$$

Reparameterised gradients: Gaussian

We have seen one case, namely,
if $\epsilon \sim \mathcal{N}(0, I)$ and $Z \sim \mathcal{N}(\mu, \sigma^2)$

Then

$$Z \sim \mu + \sigma\epsilon$$

and

$$\begin{aligned} & \frac{\partial}{\partial \lambda} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \lambda} g(z = \mu + \sigma\epsilon) \right] \end{aligned}$$

Reparameterised gradients: Gaussian

We have seen one case, namely,
if $\epsilon \sim \mathcal{N}(0, I)$ and $Z \sim \mathcal{N}(\mu, \sigma^2)$

Then

$$Z \sim \mu + \sigma\epsilon$$

and

$$\begin{aligned} & \frac{\partial}{\partial \lambda} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \lambda} g(z = \mu + \sigma\epsilon) \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma\epsilon) \frac{\partial z}{\partial \lambda} \right] \end{aligned}$$

Reparameterised gradients: Gaussian

Location

$$\frac{\partial}{\partial \mu} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] = \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \frac{\partial z}{\partial \mu} \right]$$

Reparameterised gradients: Gaussian

Location

$$\begin{aligned}\frac{\partial}{\partial \mu} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \mathbf{z}} g(\mathbf{z} = \mu + \sigma \epsilon) \frac{\partial \mathbf{z}}{\partial \mu} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \mathbf{z}} g(\mathbf{z} = \mu + \sigma \epsilon) \right]\end{aligned}$$

Reparameterised gradients: Gaussian

Location

$$\begin{aligned}\frac{\partial}{\partial \mu} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \mathbf{z}} g(\mathbf{z} = \mu + \sigma \epsilon) \frac{\partial \mathbf{z}}{\partial \mu} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \mathbf{z}} g(\mathbf{z} = \mu + \sigma \epsilon) \right]\end{aligned}$$

Scale

$$\frac{\partial}{\partial \sigma} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] = \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial \mathbf{z}} g(\mathbf{z} = \mu + \sigma \epsilon) \frac{\partial \mathbf{z}}{\partial \sigma} \right]$$

Reparameterised gradients: Gaussian

Location

$$\begin{aligned}\frac{\partial}{\partial \mu} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \frac{\partial z}{\partial \mu} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \right]\end{aligned}$$

Scale

$$\begin{aligned}\frac{\partial}{\partial \sigma} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \frac{\partial z}{\partial \sigma} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \epsilon \right]\end{aligned}$$

Reparameterised gradients: Gaussian

Location

$$\begin{aligned}\frac{\partial}{\partial \mu} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \frac{\partial z}{\partial \mu} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \right]\end{aligned}$$

Scale

$$\begin{aligned}\frac{\partial}{\partial \sigma} \mathbb{E}_{\mathcal{N}(z|\mu, \sigma^2)} [g(z)] &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \frac{\partial z}{\partial \sigma} \right] \\ &= \mathbb{E}_{\mathcal{N}(0, I)} \left[\frac{\partial}{\partial z} g(z = \mu + \sigma \epsilon) \epsilon \right]\end{aligned}$$

Multivariate calculus recap

Let $x \in \mathbb{R}^D$ and let $\mathcal{T} : \mathbb{R}^D \rightarrow \mathbb{R}^D$ be differentiable and invertible

► $y = \mathcal{T}(x)$

► $x = \mathcal{T}^{-1}(y)$

Jacobian

The Jacobian matrix $\mathbf{J} = J_{\mathcal{T}}(\mathbf{x})$ of \mathcal{T} assessed at \mathbf{x} is the matrix of partial derivatives

$$J_{ij} = \frac{\partial y_i}{\partial x_j}$$

Jacobian

The Jacobian matrix $\mathbf{J} = J_{\mathcal{T}}(x)$ of \mathcal{T} assessed at x is the matrix of partial derivatives

$$J_{ij} = \frac{\partial y_i}{\partial x_j}$$

Inverse function theorem

$$J_{\mathcal{T}^{-1}}(y) = (J_{\mathcal{T}}(x))^{-1}$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx = \int g(\mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)| dy$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx = \int g(\mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)| dy$$

and similarly for a function $h(y)$

$$\int h(y) dy$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx = \int g(\mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)| dy$$

and similarly for a function $h(y)$

$$\int h(y) dy = \int h(\mathcal{T}(x)) |\det J_{\mathcal{T}}(x)| dx$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx = \int g(\mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)| dy$$

and similarly for a function $h(y)$

$$\int h(y) dy = \int h(\mathcal{T}(x)) |\det J_{\mathcal{T}}(x)| dx$$

Rule of thumb

$$dx = |\det J_{\mathcal{T}^{-1}}(y)| dy$$

Integration by substitution

We can integrate a function $g(x)$
by substituting $x = \mathcal{T}^{-1}(y)$

$$\int g(x) dx = \int g(\mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)| dy$$

and similarly for a function $h(y)$

$$\int h(y) dy = \int h(\mathcal{T}(x)) |\det J_{\mathcal{T}}(x)| dx$$

Rule of thumb

$$dx = |\det J_{\mathcal{T}^{-1}}(y)| dy \quad dy = |\det J_{\mathcal{T}}(x)| dx$$

Change of density

Let X take on values in \mathbb{R}^D with density $f_X(x)$

Change of density

Let X take on values in \mathbb{R}^D with density $f_X(x)$
and recall that $y = \mathcal{T}(x)$ and $x = \mathcal{T}^{-1}(y)$

Change of density

Let X take on values in \mathbb{R}^D with density $f_X(x)$
and recall that $y = \mathcal{T}(x)$ and $x = \mathcal{T}^{-1}(y)$

Then \mathcal{T} induces a density $f_Y(y)$ expressed as

$$f_Y(y) = f_X(x = \mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)|$$

Change of density

Let X take on values in \mathbb{R}^D with density $f_X(x)$
and recall that $y = \mathcal{T}(x)$ and $x = \mathcal{T}^{-1}(y)$

Then \mathcal{T} induces a density $f_Y(y)$ expressed as

$$f_Y(y) = f_X(x = \mathcal{T}^{-1}(y)) |\det J_{\mathcal{T}^{-1}}(y)|$$

and then it follows that

$$f_X(x) = f_Y(y = \mathcal{T}(x)) |\det J_{\mathcal{T}}(x)|$$

Revisiting reparameterised gradients

Let Z take on values in \mathbb{R}^D with pdf $q(z|\lambda)$

Revisiting reparameterised gradients

Let Z take on values in \mathbb{R}^D with pdf $q(z|\lambda)$

The idea is to count on a *standardisation* procedure

Revisiting reparameterised gradients

Let Z take on values in \mathbb{R}^D with pdf $q(z|\lambda)$

The idea is to count on a *standardisation* procedure
a transformation $\mathcal{S}_\lambda : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that

Revisiting reparameterised gradients

Let Z take on values in \mathbb{R}^D with pdf $q(z|\lambda)$

The idea is to count on a *standardisation* procedure
a transformation $\mathcal{S}_\lambda : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that

$$\begin{aligned}\mathcal{S}_\lambda(z) &\sim \pi(\epsilon) \\ \mathcal{S}_\lambda^{-1}(\epsilon) &\sim f(z|\lambda)\end{aligned}$$

- $\pi(\epsilon)$ does not depend on parameters λ
we call it a *standard* density

Revisiting reparameterised gradients

Let Z take on values in \mathbb{R}^D with pdf $q(z|\lambda)$

The idea is to count on a *standardisation* procedure
a transformation $\mathcal{S}_\lambda : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that

$$\begin{aligned}\mathcal{S}_\lambda(z) &\sim \pi(\epsilon) \\ \mathcal{S}_\lambda^{-1}(\epsilon) &\sim f(z|\lambda)\end{aligned}$$

- ▶ $\pi(\epsilon)$ does not depend on parameters λ
we call it a *standard* density
- ▶ $\mathcal{S}_\lambda(z)$ absorbs dependency on λ

Reparameterised expectations

If we are interested in

$$\mathbb{E}_{q(z|\lambda)} [g(z)]$$

Reparameterised expectations

If we are interested in

$$\mathbb{E}_{q(z|\lambda)} [g(z)] = \int q(z|\lambda) g(z) dz$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}\mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) dz \\ &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) dz\end{aligned}$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}\mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) \mathrm{d}z \\ &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) \mathrm{d}z \\ &= \int \pi(\epsilon) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right|^{-1}\end{aligned}$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}\mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) \mathrm{d}z \\ &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) \mathrm{d}z \\ &= \int \pi(\epsilon) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right|^{-1} g(\mathcal{S}_\lambda^{-1}(\epsilon)) \mathrm{d}\epsilon\end{aligned}$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}
 \mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) dz \\
 &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) dz \\
 &= \int \pi(\epsilon) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right|^{-1} g(\mathcal{S}_\lambda^{-1}(\epsilon)) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right| d\epsilon
 \end{aligned}$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}
 \mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) dz \\
 &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) dz \\
 &= \int \pi(\epsilon) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right|^{-1} g(\mathcal{S}_\lambda^{-1}(\epsilon)) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right| d\epsilon \\
 &= \int \pi(\epsilon) g(\mathcal{S}_\lambda^{-1}(\epsilon)) d\epsilon
 \end{aligned}$$

Reparameterised expectations

If we are interested in

$$\begin{aligned}
 \mathbb{E}_{q(z|\lambda)} [g(z)] &= \int q(z|\lambda) g(z) dz \\
 &= \int \underbrace{\pi(\mathcal{S}_\lambda(z)) |\det J_{\mathcal{S}_\lambda}(z)|}_{\text{change of density}} g(z) dz \\
 &= \int \pi(\epsilon) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right|^{-1} g(\mathcal{S}_\lambda^{-1}(\epsilon)) \left| \det J_{\mathcal{S}_\lambda^{-1}}(\epsilon) \right| d\epsilon \\
 &= \int \pi(\epsilon) g(\mathcal{S}_\lambda^{-1}(\epsilon)) d\epsilon = \mathbb{E}_{\pi(\epsilon)} [g(\mathcal{S}_\lambda^{-1}(\epsilon))]
 \end{aligned}$$

Reparameterised gradients

For optimisation, we need tractable gradients

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{q(z|\lambda)} [g(z)] = \frac{\partial}{\partial \lambda} \mathbb{E}_{\pi(\epsilon)} [g(\mathcal{S}_{\lambda}^{-1}(\epsilon))]$$

Reparameterised gradients

For optimisation, we need tractable gradients

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{q(z|\lambda)} [g(z)] = \frac{\partial}{\partial \lambda} \mathbb{E}_{\pi(\epsilon)} [g(\mathcal{S}_{\lambda}^{-1}(\epsilon))]$$

since now the measure of integration does not depend on λ , we can obtain a gradient estimate

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{q(z|\lambda)} [g(z)] = \mathbb{E}_{\pi(\epsilon)} \left[\frac{\partial}{\partial \lambda} g(\mathcal{S}_{\lambda}^{-1}(\epsilon)) \right]$$

Reparameterised gradients

For optimisation, we need tractable gradients

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{q(z|\lambda)} [g(z)] = \frac{\partial}{\partial \lambda} \mathbb{E}_{\pi(\epsilon)} [g(\mathcal{S}_{\lambda}^{-1}(\epsilon))]$$

since now the measure of integration does not depend on λ , we can obtain a gradient estimate

$$\frac{\partial}{\partial \lambda} \mathbb{E}_{q(z|\lambda)} [g(z)] = \mathbb{E}_{\pi(\epsilon)} \left[\frac{\partial}{\partial \lambda} g(\mathcal{S}_{\lambda}^{-1}(\epsilon)) \right]$$

$$\stackrel{\text{MC}}{\approx} \frac{1}{M} \sum_{\substack{i=1 \\ \epsilon_i \sim \pi(\epsilon)}}^M \frac{\partial}{\partial \lambda} g(\mathcal{S}_{\lambda}^{-1}(\epsilon_i))$$

Standardisation functions

Location-scale family

- ▶ a family of distributions where for

$$F_X(x) = \Pr\{X \leq x\}$$

$$\text{if } Y = a + bX, \text{ then } F_Y(y|a, b) = F_X\left(\frac{y-a}{b}\right)$$

Standardisation functions

Location-scale family

- ▶ a family of distributions where for
 $F_X(x) = \Pr\{X \leq x\}$
if $Y = a + bX$, then $F_Y(y|a, b) = F_X(\frac{y-a}{b})$
- ▶ if we can draw from $f_X(x)$, we can draw from $f_Y(y|a, b)$

Standardisation functions

Location-scale family

- ▶ a family of distributions where for $F_X(x) = \Pr\{X \leq x\}$
if $Y = a + bX$, then $F_Y(y|a, b) = F_X(\frac{y-a}{b})$
- ▶ if we can draw from $f_X(x)$, we can draw from $f_Y(y|a, b)$
- ▶ the transformation absorbs the parameters a, b

Standardisation functions

Location-scale family

- ▶ a family of distributions where for
 $F_X(x) = \Pr\{X \leq x\}$
if $Y = a + bX$, then $F_Y(y|a, b) = F_X(\frac{y-a}{b})$
- ▶ if we can draw from $f_X(x)$, we can draw from $f_Y(y|a, b)$
- ▶ the transformation absorbs the parameters a, b

Examples: Gaussian, Laplace, Cauchy, Uniform

Standardisation functions (cont.)

Inverse cdf

- ▶ for univariate Z with pdf $f_Z(z)$ and cdf $F_Z(z)$

$$P \sim \mathcal{U}(0, 1) \quad Z \sim F_Z^{-1}(P)$$

where $F_Z^{-1}(p)$ is the *quantile function*

Standardisation functions (cont.)

Inverse cdf

- ▶ for univariate Z with pdf $f_Z(z)$ and cdf $F_Z(z)$

$$P \sim \mathcal{U}(0, 1) \quad Z \sim F_Z^{-1}(P)$$

where $F_Z^{-1}(p)$ is the *quantile function*

Gumbel distribution

- ▶ $f_Z(z|\mu, \beta) = \beta^{-1} \exp(-z - \exp(-z))$
- ▶ $F_Z(z|\mu, \beta) = \exp\left(-\exp\left(-\frac{z-\mu}{\beta}\right)\right)$
- ▶ $F_Z^{-1}(p) = \mu - \beta \log(-\log p)$

Beyond

Many interesting densities are not location-scale families

- ▶ e.g. Beta, Gamma

The inverse cdf of a multivariate rv is seldom known in closed-form

- ▶ Dirichlet, von Mises-Fisher

Automatic Differentiation VI

Motivation

- ▶ many models have intractable posteriors
their normalising constants (evidence) lacks
analytic solutions

Automatic Differentiation VI

Motivation

- ▶ many models have intractable posteriors
their normalising constants (evidence) lacks
analytic solutions
- ▶ but many models are differentiable
that's the main constraint for using NNs

Automatic Differentiation VI

Motivation

- ▶ many models have intractable posteriors
their normalising constants (evidence) lacks
analytic solutions
- ▶ but many models are differentiable
that's the main constraint for using NNs

Reparameterised gradients are a step towards
automatising VI for differentiable models

Automatic Differentiation VI

Motivation

- ▶ many models have intractable posteriors
their normalising constants (evidence) lacks
analytic solutions
- ▶ but many models are differentiable
that's the main constraint for using NNs

Reparameterised gradients are a step towards
automatising VI for differentiable models

- ▶ but not every model of interest employs rvs for
which a standardisation function is known

Example

Suppose we have some ordinal data which we assume to be Poisson-distributed

$$X|\lambda \sim \text{Poisson}(\lambda)$$

and suppose we want to impose

Differentiable models

We focus on *differentiable probability models*

$$p(x, z) = p(x|z)p(z)$$

Differentiable models

We focus on *differentiable probability models*

$$p(x, z) = p(x|z)p(z)$$

- ▶ members of this class have continuous latent variables z

Differentiable models

We focus on *differentiable probability models*

$$p(x, z) = p(x|z)p(z)$$

- ▶ members of this class have continuous latent variables z
- ▶ and the gradient $\nabla_z \log p(x, z)$ is valid within the *support* of the prior
$$\text{supp}(p(z)) = \{z \in \mathbb{R}^K : p(z) > 0\} \subseteq \mathbb{R}^K$$

VI optimisation problem

Let's focus on the design and optimisation of the variational approximation

$$\arg \min_{q(z)} \text{KL} (q(z) \parallel p(z|x))$$

VI optimisation problem

Let's focus on the design and optimisation of the variational approximation

$$\arg \min_{q(z)} \text{KL} (q(z) \parallel p(z|x))$$

Support matching constraint

To automatise the search for a variational approximation $q(z)$ we must ensure that

$$\text{supp}(q(z)) \subseteq \text{supp}(p(z|x))$$

- ▶ otherwise KL is not defined

Support matching constraint

To automatise the search for a variational approximation $q(z)$ we must ensure that

$$\text{supp}(q(z)) \subseteq \text{supp}(p(z|x))$$

- ▶ otherwise KL is not defined
- ▶ but what is the support of $p(z|x)$?

Support matching constraint

To automatise the search for a variational approximation $q(z)$ we must ensure that

$$\text{supp}(q(z)) \subseteq \text{supp}(p(z|x))$$

- ▶ otherwise KL is not defined
- ▶ but what is the support of $p(z|x)$?
- ▶ typically the same as the support of $p(z)$ as long as $p(x, z) > 0$ if $p(z) > 0$

Parametric family

So let's constrain $q(z)$ to a family \mathcal{Q} whose support is included in the support of the prior

$$\mathcal{Q} = \{q(z; \phi) : \phi \in \Phi\}$$

- ▶ a parameter vector ϕ picks out a member $q(z; \phi)$ of the family

Our objective now is

$$\arg \min_{\phi \in \Phi} \text{KL} (q(z; \phi) \parallel p(z|x))$$

ADVI

Minimising the original VI problem is equivalent to maximising the ELBO

$$\mathcal{L}(\phi) = \mathbb{E}_{q(z;\phi)} [\log p(x, z)] + \mathbb{H}(q(z|\phi)) \quad (1)$$

From the point of view of a black-box procedure, this objective poses two problems

1. intractable expectations

ADVI

Minimising the original VI problem is equivalent to maximising the ELBO

$$\mathcal{L}(\phi) = \mathbb{E}_{q(z;\phi)} [\log p(x, z)] + \mathbb{H}(q(z|\phi)) \quad (1)$$

From the point of view of a black-box procedure, this objective poses two problems

1. intractable expectations Reparameterised Gradients!
2. custom $\text{supp}(q(z; \phi))$

Idea

1. let's find a way to transform $\text{supp}(p(z))$ to the complete real coordinate space

