

NCERT 11.14 Q-25

EE23BTECH11207 -KAILASH.C

I. QUESTION:

A mass attached to a spring is free to oscillate, with angular velocity ω , in a horizontal plane without friction or damping. It is pulled to a distance x_0 and pushed towards the centre with a velocity v_0 at time $t = 0$. Determine the amplitude of the resulting oscillations in terms of the parameters ω , x_0 , and v_0 . [Hint : Start with the equation $x = a \cos(\omega t + \theta)$ and note that the initial velocity is negative.]

II. SOLUTION:

Symbols	Definition
x	Displacement
t	Time
v	Velocity
ω	Angular velocity
Θ	Phase constant

TABLE 0

SYMBOLS AND THEIR DEFINITIONS

$$x = A \cos(\omega t + \theta) \quad (1)$$

By differentiating equation-(1) with respect to time t, we get:

$$v = \frac{dx}{dt} \quad (2)$$

$$v = -A\omega \sin(\omega t + \theta) \quad (3)$$

Let at time $t=0$ s, $x=x_0$

Substituting $t=0$ s in equation (1):

$$x = A \cos(\omega(0) + \theta) \quad (4)$$

$$x_0 = A \cos(\theta) \quad (5)$$

$$v_0 = \frac{dx}{dt} \quad (6)$$

$$v_0 = -A\omega \sin(\omega(0) + \theta) \quad (7)$$

$$v_0 = -A\omega \sin(\theta) \quad (8)$$

$$-A \sin(\theta) = \frac{v_0}{\omega} \quad (9)$$

Squaring and Adding eq-(5) and eq-(9):

$$\begin{aligned} (A \cos(\theta))^2 + (A \sin(\theta))^2 \\ = (x_0)^2 + \left(\frac{v_0}{\omega}\right)^2 \end{aligned} \quad (10)$$

$$\begin{aligned} A^2(\cos^2(\theta) + \sin^2(\theta)) \\ = (x_0)^2 + \left(\frac{v_0}{\omega}\right)^2 \end{aligned} \quad (11)$$

$$A^2 = (x_0)^2 + \left(\frac{v_0}{\omega}\right)^2 \quad (12)$$

Therefore by solving the above equation we get:

$$\text{Amplitude} = A = \sqrt{(x_0)^2 + \left(\frac{v_0}{\omega}\right)^2}$$