

of discriminant 5 to the canonical form $X^2 + XY - Y^2$, where s_5 is the odd root of $x^2 = 5 \bmod p$.

This reduction algorithm, outlined by Gauss and cleverly reported in Buell [3], was described in a masterly fashion in Mathews' book [13], and thus it will be referred to as Mathews' reduction. It can be applied to any quadratic form of a non-perfect square discriminant. The square root s_5 may be computed with deterministic polynomial complexity by distinguishing whether p is congruent 3 or 1 modulo 4. In the first case, s_5 is one of the two square roots $\pm 5^{\frac{p+1}{4}} \bmod p$, and the computation clearly has deterministic polynomial complexity. In the second case Schoof's algorithm [18, 21] is applied to compute both $\sqrt{-1}$ and $\sqrt{-5} \bmod p$, thus s_5 is one of the two values $\pm \sqrt{-1}\sqrt{-5} \bmod p$. Schoof's algorithm evaluates a square root modulo p in a number of arithmetical operations of order $O(\log^t p)$ with $t \leq 10$, that is with deterministic polynomial complexity [18]. A description of Schoof's algorithm can be found in the original paper [18] or in Washington's book [21], and is based on properties of elliptic curves that are briefly summarized.

.....

1. Representation of primes in $Z(\omega)$

.....

2. Computational remarks

The representations of primes $\pi \in Q(\omega)$ as the sums of two squares, established by Theorems 2 and 3, are computed with deterministic polynomial complexity using procedures that will be described separately for the different cases.

4.1 Primes $p \equiv 13, 17 \bmod 20$

The representation $p = a^2 + b^2$ does not need any further comment, since $p \equiv 1 \bmod 4$ is the sum of two integer squares by Fermat's theorem, and its computation was shown in Section 2.

4.2 Primes p

$$+ y_0 F_{2k+1})$$

.....

5. Conclusions

The representation of primes as sums of two squares in $Z(\omega)$ allows us to formulate a theorem concerning the representation of any integer in $Z(\omega)$:

Theorem 4. *In the field $Q(\omega)$, any totally positive integer*

.....

The computation of the two-square representation of any element in $Z(\omega)$ is equivalent to factoring, as in Z , therefore at the present time only the two-square representations of primes can be computed with deterministic polynomial complexity.

References

- [1] B. Berndt and R. Evans, Sums of Gauss, Jacobi and Jacobsthal, *Journal of Number Theory*, Vol. 11 (1979), pp. 349–398.
- [2] J. Brillhart, Note on representing a prime as a sum of two squares, *Mathematics of Computation*, Vol. 26 (1972), pp. 1011–1013.
- [3] D. A. Buell, *Binary Quadratic Forms*, Springer-Verlag, New York, 1989.
- [4] D. A. Cox, *Primes of the Form*, Wiley, New York, 1989.
- [5] F. W. Dodd, *Number Theory in the Quadratic Field with Golden Section Unit*, Polygonal Pub. House, Paissac, 1983.
- [6] C.F. Gauss, *Disquisitiones Arithmeticae*, Springer-Verlag, New York, 1986.
- [7] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, Oxford, 1971.
- [8] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, New York, 1998.
- [9] D. E. Knuth, *The Art of Computer Programming: Seminumerical Algorithms*, Addison-Wesley, Reading, 1970.
- [10] S. Lang, *Elliptic Functions*, Springer-Verlag, New York, 1987.
- [11] F. Lemmermeyer, *Reciprocity Laws*, Springer-Verlag, New York, 2000.
- [12] R. Lidl and H. Niederreiter, *Finite Fields*, Addison-Wesley, Reading