

SINGULAR VALUE DECOMPOSITION

MATH 578 Mini-Seminar Talk

Kai Yang

McGill University
Montréal, Quebec Canada H3A 1A2

McGill University

October 22, 2020

Introduction

- *Singular Value Decomposition (SVD)* is one of most important tool for mathematical computation
- Contents of this presentation are taken from *Numerical linear algebra* and *Matrix Computations*
- As we'll see later, SVD is connect to eigenvalues; but in applications, eigenvalues are more connected to the behaviour of iterated forms of A , e.g., A^k or e^{tA} ; SVD is more related to the behavior of A or A^{-1}
- We'll start with the geometric observation as a motivation for SVD

Geometric Observation




Figure: SVD of 2×2 matrix [1]

- We assume A is a real matrix here for the sake of geometric interpretation
- *The image of unit sphere under any $m \times n$ matrix is a hyperellipse, – a hyperellipse is a m –dimensional generalization of an ellipse.*

Geometric Interpretation

- Define n *singular values* of A : these are the lengths of n principal semiaxes of AS – it's conventional to assume that singular values are numbered in a descending order, $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n > 0$
- Define n *left singular vectors* of A : these are the *unit* vectors $\{u_1, u_2, \dots, u_n\}$ oriented in the directions of principal semiaxes of AS , numbered corresponding to the singular values
- Define n *right singular vectors* of A : these are the *unit* vectors $\{v_1, v_2, \dots, v_n\} \in S$ that are preimages of principal semiaxes of AS , number so that $Av_j = \sigma_j u_j$
- For the moment, let's assume A of full column rank n .

Reduced SVD

- Write the linear map as $Av_j = \sigma_j u_j$, $\forall 1 \leq j \leq n$; and express in matrix form:

$$AV = \hat{U}\hat{\Sigma}$$

- $\hat{\Sigma}$ is a $n \times n$ diagonal matrix with diagonal entries $\sigma_1, \sigma_2, \dots, \sigma_n > 0$
- $\hat{U} \in \mathbb{C}^{m \times n}$ has orthonormal columns $u_1, u_2, \dots, u_n \in \mathbb{C}^m$
- $V \in \mathbb{C}^{n \times n}$ is a unitary matrix with columns v_1, v_2, \dots, v_n
- Then we have $A = \hat{U}\hat{\Sigma}V^*$
- Note the column vectors of \hat{U} does not form a basis of \mathbb{C}^n – this is why it's called "reduced"

Full SVD

- By adjoining $m - n$ orthonormal columns \hat{U}_{m-n} , \hat{U} can be extended to a unitary matrix – let's call the result U
- $\hat{\Sigma}$ also needs to change to accommodate this – by adjoining $m - n$ rows of 0
- As a result, the full SVD is

$$A = \hat{U} \hat{\Sigma} V^* = \begin{bmatrix} \hat{U} & \hat{U}_{m-n} \end{bmatrix} \begin{bmatrix} \hat{\Sigma} \\ 0 \end{bmatrix} V^* =: U \Sigma V^*$$

- Now we can discard the initial assumption that A has full (column) rank – all it changes is now not n but only r of the left singular vectors of A are determined by the geometry of hyperellipse, then we'll have $\hat{U} \in \mathbb{C}^{m \times r}$, $\hat{\Sigma} \in \mathbb{C}^{r \times r}$ will be a diagonal matrix with positive diagonal entries $\sigma_1, \sigma_2, \dots, \sigma_r$, put at the upper-left corner of the otherwise-0 matrix $\Sigma \in \mathbb{C}^{m \times n}$

Formal Definition

- Given $A \in \mathbb{C}^{m \times n}$, a SVD of A is a factorization

$$A = U \Sigma V^*$$

- where $U \in \mathbb{C}^{m \times m}$, $V \in \mathbb{C}^{n \times n}$ are both unitary; $\Sigma \in \mathbb{R}^{m \times n}$ is diagonal, with diagonal entries $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$ and $\sigma_{r+1}, \dots, \sigma_p = 0$, where $p = \min\{m, n\}$
- The image of a unit sphere in \mathbb{R}^n under linear map $A = U \Sigma V^*$ will indeed be a hyperellipse in \mathbb{R}^m : the unitary map V^* preserves the sphere, then the diagonal matrix Σ stretches the sphere into a hyperellipse aligned with canonical basis, finally unitary map U rotates or reflects the hyperellipse without changing its shape.

Existence and Uniqueness

Theorem

Every matrix $A \in \mathbb{C}^{m \times n}$ has a SVD. Furthermore, the singular values are uniquely determined, and if A is square and σ_j are distinct, the left and right singular vectors $\{u_j\}$ and $\{v_j\}$ are uniquely determined up to the complex signs.

- The existence statement can be proved by induction on the dimension of A , whose induction step is established by: for submatrix B of A ,

$$B = U_2 \Sigma_2 V_2^* \Rightarrow A = U_1 \begin{bmatrix} 1 & 0 \\ 0 & U_2 \end{bmatrix} \begin{bmatrix} \sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & V_2 \end{bmatrix}^* V_1^*$$

is a SVD of A .

- The uniqueness statement is justified by the geometric interpretation: should the semiaxis lengths of a hyperellipse are distinct, then the semiaxes are determined by the geometry signs.

SVD v.s. Eigen-Decomposition

- SVD represents a change of bases, i.e., every matrix is diagonal if using proper bases for domain and range spaces – it uses two different bases, the sets of left (for range space) and right (for domain space) singular vectors. Eigen-Decomposition uses one bases – the set of eigenvectors
- SVD uses orthonormal bases; while eigen-decomposition in general does not
- Not all matrices (even square ones) have an eigenvalue decomposition, but all matrices have a SVD

Matrix Properties via SVD I

- $\text{rank}(A) = r$, the number of nonzero singular values. (*note that U, V are full rank*)
- $\text{range}(A) = \langle u_1, \dots, u_r \rangle$ and $\text{null}(A) = \langle v_{r+1}, \dots, v_n \rangle$. (*observing the range and null spaces of Σ*)
- $\|A\|_2 = \sigma_1$ and $\|A\|_F = \sqrt{\sum_{i=1}^r \sigma_i^2}$. (*U, V are unitary, together with the fact that ℓ_2 norm and Frobenius norm is invariant under unitary multiplication*)
- The nonzero singular values of A square roots of nonzero eigenvalues of A^*A or AA^* . (*substitution by SVD*)
- If $A = A^*$, the singular values of A are the absolute values of eigenvalues of A . (*using the fact that the eigenvalues for every Hermitian matrix are real, and the eigenvectors corresponding to distinct eigenvalues are orthogonal*)

Matrix Properties via SVD II

- $\forall A \in \mathbb{C}^{m \times m}$, $|\det(A)| = \prod_{i=1}^m \sigma_i$. (substitution by SVD)
- $\forall A \in \mathbb{C}^{m \times n}$, $E \in \mathbb{C}^{m \times n}$,

$$\sigma_{\max}(A + E) \leq \sigma_{\max}(A) + \|E\|_2$$

$$\sigma_{\min}(A + E) \geq \sigma_{\min}(A) - \|E\|_2$$

(recall $Av_j = \sigma_j u_j$, which implies $\sigma_{\min}(A) \leq \frac{\|Ax\|_2}{\|x\|_2} \leq \sigma_{\max}(A)$)

- $\forall A \in \mathbb{C}^{m \times n}$, $m > n$, and $\forall z \in \mathbb{C}^m$,

$$\sigma_{\max}\left(\begin{bmatrix} A & z \end{bmatrix}\right) \geq \sigma_{\max}(A)$$

$$\sigma_{\min}\left(\begin{bmatrix} A & z \end{bmatrix}\right) \leq \sigma_{\min}(A)$$

()

Low Rank Approximations

- $A = \sum_{i=1}^r \sigma_i u_i v_i^*$. (*trivial*)
- Define $A_k = \sum_{i=1}^k \sigma_i u_i v_i^*$ for some $0 \leq k \leq r$, and define $\sigma_{k+1} = 0$ if $k = \min\{m, n\}$. Then

$$\min_{B \in \mathbb{C}^{m \times n}, \text{rank}(B) \leq k} \|A - B\|_2 = \|A - A_k\|_2 = \sigma_{k+1}$$

(*second equality is trivial; prove first equality by assuming $\|A - B\|_2 < \|A - A_k\|_2 = \sigma_{k+1}$, then $\forall w \in \text{null}(B)$ ($\dim \geq n - k$),*

$$\|Aw\|_2 = \|(A - B)w\|_2 \leq \|A - B\|_2 \|w\|_2 < \sigma_{k+1} \|w\|_2$$

on the other hand, the first $k + 1$ right singular vectors of A span a $k + 1$ -dimensional space s.t. for all w in it, $\|Aw\|_2 \geq \sigma_{k+1} \|w\|_2$, which leads to contradiction)

- Similar to above statement (*and proof*), we also have

$$\min_{B \in \mathbb{C}^{m \times n}, \text{rank}(B) \leq k} \|A - B\|_F = \|A - A_k\|_F = \sqrt{\sigma_{k+1}^2 + \cdots + \sigma_r^2}$$

Bibliography

- [1] Lloyd Trefethen. *Numerical linear algebra*. Philadelphia, Pa: Society for Industrial and Applied Mathematics (SIAM, 3600 Market Street, Floor 6, Philadelphia, PA 19104, 1997. ISBN: 0898719577.
- [2] Gene H. Golub. *Matrix Computations*. J. Hopkins Uni. Press, Jan. 7, 2013. ISBN: 1421407949. URL: https://www.ebook.de/de/product/20241149/gene_h_golub_matrix_computations.html.