



XXTitle

Bergur Snorrason



Faculty of XX
University of Iceland
2020

XXTITLE

Bergur Snorrason

XX ECTS thesis submitted in partial fulfillment of a
Magister Scientiarum degree in XX

Advisor

XXNN1

XXNN2

Faculty Representative

XXNN3

M.Sc. committee

XXNN4

XXNN5

Faculty of XX

School of Engineering and Natural Sciences

University of Iceland

Reykjavik, XXmonth 2020

XXTitle

XXShort title (50 characters including spaces)

XX ECTS thesis submitted in partial fulfillment of a M.Sc. degree in XX

Copyright © 2020 Bergur Snorrason

All rights reserved

Faculty of XX

School of Engineering and Natural Sciences

University of Iceland

XXFaculty street address

XXFaculty postal code, Reykjavik, Reykjavik
Iceland

Telephone: 525 4000

Bibliographic information:

Bergur Snorrason, 2020, XXTitle, M.Sc. thesis, Faculty of XX, University of Iceland.

ISBN XX

Printing: Háskólaprent, Fálkagata 2, 107 Reykjavík
Reykjavik, Iceland, XXmonth 2020

Dedication

Abstract

Útdráttur á ensku sem er að hámarki 250 orð.

Útdráttur

Hér kemur útdráttur á íslensku sem er að hámarki 250 orð. Reynið að koma útdráttum á eina blaðsíðu en ef tvær blaðsíður eru nauðsynlegar á seinni blaðsíða útdráttar að hefjast á oddatölusíðu (hægri síðu).

Preface

Formála má sleppa og skal þá fjarlægja þessa blaðsíðu. Formáli skal hefjast á odd-atölu blaðsíðu og nota skal Section Break (Odd Page).

Ekki birtist blaðsíðutal á þessum fyrstu síðum ritgerðarinnar en blaðsíðurnar teljast með og hafa áhrif á blaðsíðutal sem birtist með rómverskum tölum fyrst á efnisyfirliti.

Contents

List of Figures	xi
List of Tables	xiii
Abbreviations	xv
Acknowledgments	1
1. Introduction	3
2. Preliminaries	5
2.1. Measure theory	5
2.2. Functional analysis	5
2.2.1. Hahn-Banach	5
2.2.2. Riesz representation theorem	5
3. Rudin-Carleson theorem	7
4. F. and M. Riesz theorem	9
5. A generalization of the Rudin-Carleson theorem	11
A. Annađ	15

List of Figures

List of Tables

Abbreviations

Í þessum kafla mega koma fram listar yfir skammstafanir og/eða breytuheiti. Gefið kaflanum nafn við hæfi, t.d. Skammstafanir eða Breytuheiti. Þessum kafla má sleppa ef hans er ekki þörf.

The section could be titled: Glossary, Variable Names, etc.

Acknowledgments

Í þessum kafla koma fram þakkir til þeirra sem hafa styrkt rannsóknina með fjárframlögum, aðstöðu eða vinnu. T.d. styrktarsjóðir, fyrirtæki, leiðbeinendur, og aðrir aðilar sem hafa á einhvern hátt aðstoðað við gerð verkefnisins, þ.m.t. vinir og fjölskylda ef við á. Þakkir byrja á oddatölusíðu (hægri síðu).

1. Introduction

2. Preliminaries

2.1. Measure theory

2.2. Functional analysis

2.2.1. Hahn-Banach

2.2.2. Riesz representation theorem

3. Rudin-Carleson theorem

Theorem 1. *Let E be a closed subset of \mathbb{T} of Lebesgue-measure 0, let f be a continuous function on E and let T be a simply connected subset of \mathbb{C} such that $f(\overline{\mathbb{D}}) \subset T$. Then there exists an $F \in \mathcal{A}$, such that $F = f$ on E and $F(\overline{\mathbb{D}}) \subset T$. (TODO skilgreina allt)*

We will break the proof into several (TODO how many?) lemmas.

Lemma 1. *Let E be a closed set of Lebesgue-measure 0. Then there exists an integrable function $\mu > 1$ such that μ is continuous on $\mathbb{T} \setminus E$, $\mu = +\infty$ on E , if $w \in E$ then $\mu(z) \xrightarrow{z \rightarrow w} +\infty$, and μ has a bounded derivative on any closed subarc of $\mathbb{T} \setminus E$.*

Proof. The function μ is found by solving the Dirichlet problem. (TODO Thomas Ransford p. 95 4.2.6) (TODO finish this) \square

Lemma 2. *If f is a simple continuous function on E such that $\operatorname{Re} f \geq 0$, then there exists an $F \in \mathcal{A}$ such that $F = f$ on E and $\operatorname{Re} F \geq 0$ on $\overline{\mathbb{D}}$.*

Proof. We will show that this holds in the case where $E = E_0 \cup E_1$, $f = 0$ on E_0 , $f = \alpha \neq 0$ on E_1 and $\operatorname{Re} \alpha \geq 0$. This suffices since simple functions are finite linear combinations of characteristic functions. \square

Proof. TODO \square

Corollary 1 (Fatou). *Let E be a closed subset of \mathbb{T} of Lebesgue-measure 0. There exists a function $f \in \mathcal{A}$ that vanishes on E and nowhere else.*

Proof. It's clear from the theorem (TODO add ref) that there exists a function $f \in \mathcal{A}$ that vanishes on E . TODO \square

4. F. and M. Riesz theorem

Theorem 2. *All measures in \mathcal{A}^\perp are absolutely continuous with regards to the Lebesgue-measure on \mathbb{T} . (TODO define all)*

Proof. TODO

□

5. A generalization of the Rudin-Carleson theorem

This borrows from Bishop (reference TODO).

Theorem 3 (General Rudin-Carleson theorem). *Let X be a compact Hausdorff space, $V = (C(X), \|\cdot\|_\infty)$, B be a closed subspace of $C(X)$, B^\perp be the annihilating measures for B , S be a closed subset of X , and f be a continuous function on S . If $\int_S f d\mu = 0$ holds for all $\mu \in B^\perp$ then there exists a function $F \in B$ such that $F = f$ on S .*

Proof. Since f is continuous and S is a closed subset of a compact set, and therefore also compact, f is bounded. So we can, with out loss of generality, assume that $|f| < r < 1$ on S . Let U_r be the subset of B defined by $U_r = \{g; \|g\| < r\}$ and ϕ be the mapping from B to $C(S)$ that sends a member of B to its restriction on S . It suffices to show that $f \in \phi(U_r)$. Let's first show that $f \in \overline{\phi(U_r)} =: V_r$, by assuming otherwise, and showing it leads to a contradiction.

We now assume $f \notin V_r$. By Hahn-Banach (TODO ref) we can define a bounded linear functional α , such that $\alpha(f) > 1$ and $|\alpha(h)| < 1$, for $h \in V_r$. We can then define a measure μ_1 by the Riesz-representation theorem (TODO ref) that fulfills

$$\alpha(g) = \int g d\mu_1$$

for all $g \in C(S)$. We will refer to the associated functional on B by $\beta(g) = \phi(\alpha(g))$. Since $\phi(g) \in V_r$ for all $g \in U_r$ we have that

$$\beta(g) = \alpha(\phi(g)) < 1,$$

for all $g \in U_r$, due to the construction of α . From this we get

$$\begin{aligned} \|\beta\| &= \sup\{|\beta(g)|; |g| < 1\} \\ &= \sup\{(1/r)|\beta(g)|; |g| < r\} \\ &\leq \sup\{(1/r); |g| < r\} \\ &= 1/r. \end{aligned}$$

5. A generalization of the Rudin-Carleson theorem

Let's denote the Riesz representation of β by μ_2 , set $\mu = \mu_1 - \mu_2$ and see that $\mu \in B^\perp$. But

$$0 = \left| \int_S f d\mu \right| \geq \int_S f d\mu_1 - r \|\mu_2\| \geq \int_S f d\mu_1 - r \frac{1}{r} > 1 - r \frac{1}{r} = 0,$$

where the first equality is the assumption in the theorem. This is the contradiction that gives that $f \in V_r$. We can now take a F_1 in U_r , and therefore also in B such that $|f - F_1| < \lambda/2$ on S , with $\lambda := 1 - r$. Remember that $F_1 \in U_r$ implies that $\|F_1\| < r$. Now let $f_1 = f - F_1$ and use the same method as above to obtain an F_2 such that $\|F_2\| < \lambda/2$ and $|f - F_2| < \lambda/4$ on S . Iterating this process yields a series $(F_n)_{n \in \mathbb{N}}$ from B that fulfill $\|F_n\| < 2^{1-n}\lambda$ for $n > 1$ and

$$\left| f - \sum_{k=1}^n F_k \right| < 2^{-n}\lambda$$

on S for $n > 1$. We finally let

$$F = \sum_{k=1}^{\infty} F_k.$$

Now $F \in B$,

$$\|F\| \leq \|F_1\| + \|F - F_1\| = r + \sum_{k=2}^{\infty} 2^{1-k}\lambda = r + \lambda = 1,$$

and $F = f$ on S . (TODO bæta við matinu, svo línan að ofan meiki sens) □

Corollary 2. *Let X be a compact Hausdorff space, $V = (C(X), \|\cdot\|_\infty)$, B be a closed subspace of $C(X)$, B^\perp be the annihilating measures for B , S be a closed subset of X , and f be a continuous function on S . If S is B^\perp -null (TODO define this) then there exists a function $F \in B$ such that $F = f$ on S .*

Proof. If S is B^\perp -null we have that $\int_S f d\mu = 0$ for all $\mu \in B^\perp$ (TODO add ref to above). □

Remark 1. *The corollary is the version of the theorem from Bishop (TODO add ref here). Note also that if we set $X = \mathbb{T}$ and $B = \mathcal{A}$ we can use F and M . Riesz (TODO add ref here) to prove the classical Rudin-Carleson theorem (TODO add ref here).*

It is of course worth noting an applications of (TODO REF to theorem) where the corollary fails.

Example 1. Let $X = \mathbb{T}$, $B = \mathcal{A}$ (TODO define this), E be a closed m -null (TODO define this, an maybe change to m_σ) subset of $\partial\mathbb{T}$ that is not dense in E , $F = \{e^{i\theta}; a \leq \theta \leq b\}$, and choose a and b such that E and F are disjoint and $a \neq b$. The last assumption restricts us to E that are not dense in the \mathbb{E} . Since $a \neq b$ we obtain that $S := E \cup F$ does not fulfill the requirements of the Rudin-Carelsen theorem (TODO add ref here) nor the above corollary (TODO add ref). Let's choose f such that $f = 0$ on F , and f is continuous on S . We now have for all $\mu \in \mathcal{A}^\perp$

$$\begin{aligned} \left| \int_S f d\mu \right| &= \left| \int_E f d\mu + \int_F f d\mu \right| \\ &\leq \left| \int_E f d\mu \right| + \left| \int_F f d\mu \right| \\ &= 0 + \left| \int_F f d\mu \right| \\ &= 0. \end{aligned}$$

The F. and M. Riesz theorem (TODO add ref here) tells us that since E is m -null it is also μ -null, which gives the third step. The final step stems from the fact that f vanishes on F . We now see that X , B , and f are all as in the theorem (TODO add ref) so there exists a $F \in B$, such that $F = f$ on S .

A. Annaď