

Smoothed complexity of local max-cut for special graphs

December 5, 2017

Abstract

The problem of finding a max-cut in graphs has many applications and motivation. As this problem to be NP-hard, research was made on simpler problems such as a local max-cut.

1 Introduction

Let $G = (V, E)$ be a graph with n vertices and m edges. Let $w : E \rightarrow [-1, 1]$ be an edge weight function. The local max-cut problem consists in finding a partition of the vertices σ such that the total cut weight, defined as :

$$h(\sigma) = \frac{1}{2} \sum_{uv \in E} (1 - \sigma(u)\sigma(v))w(uv)$$

is locally optimal. By locally optimal, we mean that there exists no vertex v such that, if we flip the vertex, i.e. change the sign of $\sigma(v)$, then $h(\sigma)$ increases.

A really naive algorithm called FLIP solves this problem. It finds a vertex which if flipped leads to an amelioration of the total weight cut, flips it and repeat until no such vertex exists. Some instances of graphs have been found to have an exponential number of steps before terminating. However, for most of the graphs, FLIP terminates in a reasonable time. Moreover, when adding a small amount of noise to the complicated graphs, FLIP's running time improved greatly. This lead to the study of the smoothed complexity

of FLIP, in which noise is added to the edge weights.

Etscheid and Röglin (2014) [?] proved that this complexity was at most quasi-polynomial in n for arbitrary graphs, with the insight that it may be polynomial. Angel et al. (2016) [?] proved that the complexity was polynomial for complete graphs.

We study here other special graphs, for which the complexity is polynomial with the hope that it would extend for general graphs.

2 Notation and preliminary lemmas

Let $X = (X_e)_{e \in E} \in [-1, 1]^E$ a random vector with independent entries, corresponding to the edge weights. We assume that X_e has density f_e with respect to the Lebesgue measure, and we denote $\phi = \max_{e \in E} \|f_e\|_\infty$.

Lemma 2.1 (*Lemma 2.1 [?]*)

Let $\alpha_1, \dots, \alpha_k$ be k linearly independent vectors in \mathbb{Z}^E . Then the joint density of $(\langle \alpha_i, X \rangle)_{i \leq k}$ is bounded by ϕ^k . In particular, if sets $J_i \in \mathbb{R}$ have measure at most ϵ each, then

$$\mathcal{P}(\forall i \in [k], \langle \alpha_i, X \rangle \in J_i) \leq (\phi\epsilon)^k$$

We define a move vector α_v as a vector indexed by E whose entries are :

$$\alpha_{uv} = \begin{cases} \sigma(u)\sigma(w) & \text{if } uw \in E \text{ and } ((u = v) \text{ or } (w = v)) \\ 0 & \text{otherwise} \end{cases}$$

For a sequence $L = (v_1, \dots, v_l)$ of l moves and initial state σ_0 , let $\alpha_i, i \in [l]$ be the corresponding move vectors. Let σ_t be the state just after flip of vertex v_t . We define matrix A_L as the concatenation of the move vectors as columns. We call a sequence ϵ -slowly improving if all moves yield an improvement of at most ϵ .

We introduce here the concept of critical block. A block B is defined as a substring of a sequence L .

Let $S(L)$ be the set containing the distinct vertices in L and $s(L)$ be its cardinality, $s_1(L)$ the number of distinct vertices in L that appear only once, $s_2(L)$ the number of distinct vertices in L that appear multiple times. Let $l(B)$ be the length of the block, i.e. the number of moves. A block B is

critical if $l(B) \geq (1 + \beta)s(B)$ and every block B' strictly contained in B has $l(B') < (1 + \beta)s(B')$.

Lemma 2.2 (Lemma 4.1 [?])

For complete graphs with n vertices: fix any positive integer $n \geq 2$ and a constant $\beta > 0$. Given a sequence consisting of $s(L) < n$ letters and with length $l(L) \geq (1 + \beta)s$, there exists a critical block B in L . Moreover, a critical block satisfies $l(B) = \lceil (1 + \beta)s(B) \rceil$. Moreover $\text{rank}(B) \geq \frac{1 + 4\beta}{1 + 3\beta}s(B)$

Lemma 2.3

If $s(L) < n$, and $\text{rank}(L) \geq (1 + \theta)s(L)$ then

$$P(L \text{ is } \epsilon\text{-slowly improving from some } \sigma_0) \leq (2n/\epsilon)^{s_0} (64\phi\epsilon)^{\text{rank}(A_L)}$$

Where s_0 is the number of vertices that have at least one neighbor which is not in L .

Proof. Let I be the set of the edges corresponding to independent rows in A_L . They do not depend on σ_0 since the starting configuration only multiplies each row by 1 or -1.

Define $T = \{v \in V : vw \in I \text{ or } v \in S(L)\}$. $|T| \leq 2\text{rank}(A_L) + s(L)$.

We split $h(\sigma)$ in three part $h_0(\sigma), h_1(\sigma), h_2(\sigma)$ Where h_i is the restriction of h to the edges that have i endpoints in T .

$h_0(\sigma_t) - h_0(\sigma_{t-1}) = 0$. Since the edges whose both endpoints are not flipped do not provoke a change in the total weight cut.

$$\begin{aligned} h_1(\sigma_t) - h_1(\sigma_{t-1}) &= -\sigma_t(v_t) \sum_{u \notin T, uv_t \in E} \sigma_0(u) X_{uv_t} \\ &= \sigma_t(v_t) Q(v_t). \end{aligned}$$

where $Q(v_t) = -\sum_{u \notin T, uv_t \in E} \sigma_0(u) X_{uv_t}$. Since $|X_e| \leq n$ and the maximum neighbours of a vertex is n , $Q(v_t) \in [-n, n]$. By defining $D = 2\epsilon\mathbb{Z} \cap [-n, n]$, there exists some $d(v_t) \in D$ such that $|Q(v_t) - d(v_t)| \leq \epsilon$.

$h_2(\sigma_t) - h_2(\sigma_{t-1}) = \langle \alpha', X \rangle$ where

$$\alpha'_t = \begin{cases} -\sigma_t(u)\sigma_t(w) & uw \in E, v_t \in \{u, w\}, \{u, w\} \subseteq T \\ 0 & \text{otherwise} \end{cases}$$

Since α'_t concerns only the rows linearly independent from A_L , $\text{rank}([\alpha'_t]_{t \leq l}) = \text{rank}(A_L)$.

$$h(\sigma_t) - h(\sigma_{t-1}) = \langle \alpha', X \rangle + \sigma_t(v_t)d_t + \delta_t \text{ where } |\delta_t| \leq \epsilon$$

Since $|\delta_t| \leq \epsilon$, L is ϵ -slowly improving implies that:

$$|\langle \alpha', X \rangle + \sigma_t(v_t)d_t| \leq 2\epsilon \quad \forall t \leq l$$

We need that $\langle \alpha', X \rangle \in [-d_t - 2\epsilon, -d_t + 2\epsilon] \cup [d_t - 2\epsilon, d_t + 2\epsilon]$, Using lemma 2.1, this is at most $8\phi\epsilon^{\text{rank}(A_L)}$. Using union bound over $\sigma_{t \in T}$ and d :

$$P(L \text{ is } \epsilon\text{-slowly improving from some } \sigma_0) \leq 2^{2\text{rank}(A_L)+s} \left(\frac{2n}{\epsilon}\right)^{s_0} (8\phi\epsilon)^{\text{rank}(A_L)}$$

Remark the s_0 instead of s in exponent, since vertices that have no-non flipped neighbors need not be taken in this bound.

By using $\text{rank}(L) \geq (1 + \theta)s(L)$, we get the desired bound.

3 Proof for graphs with one clique and low degrees vertices

Let $G = (V, E)$ be a graph with n vertices and m edges. Assume that this graph contains a clique H of r vertices and that the degree of vertices in the set $G \setminus H$ is at most $\log(n)$.

Proposition 3.1 *With high probability, there exists no ϵ -slowly improving sequence of length $2n$ from any starting configuration σ_0 , for ϵ is $O(1/\text{poly}(n))$.*

Proving this proposition implies that the smoothed complexity is $O(\text{poly}(n))$. If there exists no such sequence, then $2n^2/\epsilon$ sequence of $2n$ moves yield an improvement of at least $2n^2$ which is the maximum improvement possible since $h(\sigma) \in [-n^2, n^2]$.

Thus number of steps is $O(n^3/\epsilon)$ which is $O(\text{poly}(n))$.

We will prove the proposition by considering different sequences of size $2n$.

Let $p > 1$, $l(L) = 2n$:

E is the event corresponding to $\exists L, \sigma_0$, s.t. L is ϵ -slowly improving from σ_0

E_1 is the event corresponding to $\exists L, \sigma_0$, s.t. L is ϵ -slowly improving from σ_0 and $S(L) \not\subseteq H$.

E_2 is the event corresponding to $\exists L, \sigma_0$, s.t. L is ϵ -slowly improving from σ_0 and $S(L) \subseteq H$ and $s(L) < r$.

E_3 is the event corresponding to \exists critical block B, σ_0 , s.t. B is ϵ -slowly improving from σ_0 and $S(L) \subseteq H$, $s(B) = r$. By the lemma 2.2 on existence of critical blocks and the fact that if $s(L) = n$ implies that some vertex with degree at most $\log(n)$ is chosen we can have this bound:

$$P(E) \leq P(E_1) + P(E_2) + P(E_3)$$

Consider E_1 . Fix L and σ_0 then there must be a vertex $v \in S(L)$ whose degree is at most $\log(n)$. By lemma 2.1, the probability that $\alpha_v \in [0, \epsilon] = \phi\epsilon$. Using union bound on the number of vertices and the starting configuration of those vertices we have:

$$P(E_1) \leq 2^{\log(n)} n \phi \epsilon \quad (1)$$

By taking $\epsilon = n^{-2-\eta} \phi^{-1}$ with $\eta > 0$ there exists no such sequence with high probability.

Now consider E_2 . We consider the subgraph G' induced by the restriction to the clique H . We observe that $\text{rank}'_G(A_L) \leq \text{rank}_G(A_L)$. Since $G'(A_L)$ is a submatrix of $G(A_L)$.

By lemma 2.2, there exists a critical block B whose rank is at least $1.25 s(B)$. We now can use lemma ?? to have this bound:

$$P(B \text{ is } \epsilon\text{-slowly improving from some } \sigma) \leq 2 \left(\frac{4n}{\epsilon} \right)^{s(B)} (8\phi\epsilon)^{\frac{5s(B)}{4}}$$

Because the number of blocks using s letters is n^{2s} , we have:

$$P(E_2) \leq 2 \sum_{s < n} (64\phi^{5/4} n^3 \epsilon^{1/4})^s$$

By choosing $\epsilon = n^{-(12+\eta)} \phi^{-5}$ this sums goes to zero.

For E_3 , we use a trick to show that the $rank(A_L) \geq 1.25s(B) - (s(B) - s_0(B))$. We choose some $w \in V \setminus H$. For each vertex v which has a non-flipped neighbour, we delete that edge and add vw to the graph. This does not change $rank(A_L)$ since the row added is the same as the row deleted times 1 or -1. Now we add edges from w to the remaining vertices on the clique, increasing the rank by at most $s(B) - s_0(B)$. The subgraph G' determined by $H \cup \{w\}$ is thus complete and we can use lemma 2.2 to have $rank_{G'}(A_L) \geq 1.25s(B)$. Then $rank_G(A_L) \geq 1.25s(B) - (s(B) - s_0(B))$. By lemma 2.3 we have :

$$P(E_3) \leq \sum_{s < n} n^{2s} \left(\frac{2n}{\epsilon}\right)^{s_0} (64\phi\epsilon)^{s/4+s_0} \leq \sum_{s < n} (Cn^3\phi^{5/4}\epsilon^{1/4})^s$$

By choosing $\epsilon = n^{-(12+\eta)}\phi^{-5}$ this sum goes to zero, thus concluding the proof.

4 Proof for graphs with multiple edge-disjoint cliques

Let $G = (V, E)$ be a graph with n vertices and m edges. Assume that this graph contains cliques H_1, \dots, H_o of r_1, \dots, r_o vertices and that the degree of vertices in the set $G \setminus H$ is at most $\log(n)$. Furthermore, there exists no edge going from one clique to another.

The precedent proof can be easily extended to those graphs. We will prove a similar proposition, but on longer sequences.

Proposition 4.1

With high probability, there exists no ϵ -slowly improving sequence of length $2n^2$ from any starting configuration σ_0 , for ϵ is $O(1/\text{poly}(n))$.

Lemma 4.2 *Let L be a sequence of q moves such that $S(L) \subseteq \bigcup_{i \leq k} A_i \subseteq V$, where A_1, \dots, A_k are edge-disjoint sets. Then, there exists a sequence with the same vertices but a different ordering on the moves such that $\forall l < q, l < j \leq q$, if $v_l \in A_i$ and $v_j \in A_i$, then $v_d \in A_i \quad \forall l \leq d \leq j$*

Proof. The proof is very straightforward. Suppose we have v_t and v_{t+1} which are edge-disjoint, the amelioration brought by v_t is equal to :

$$-\sigma(v_t) \sum_{u \in V, uv_t \in E} w_{uv_t} \sigma(u_t) = -\sigma(v_{t+1}) \sum_{u \in V, uv_{t+1} \in E} w_{uv_{t+1}} \sigma(u_{t+1})$$

Since v_t and v_{t+1} are edge-disjoint. We can then swap them, and both are still improving moves with the same amelioration of total weight. By repeating the swaps, we reach a sequence where all moves of vertices $\in A_i \quad \forall i \leq k$ are consecutive.

If we consider now sequences of length $2n^2$. Either there is a vertex with logarithmic degree and the whole sequence is not ϵ -slowly improving either we can reorder them with the previous lemma, such that vertices belonging to the same clique are consecutive. Since the number of cliques is upperbounded by n , by pigeonhole argument we have at least one sequence of size at least $2n$, which contains vertices from only a clique. We showed that with high probability such a sequence is not ϵ -slowly improving. The whole sequence is then not ϵ -slowly improving, concluding the proof.

5 Proof for two cliques with a matching

Now that we know how to deal with cliques without connection between them, we want to study a simple case where cliques are connected.

Let $G = (V, E)$ be a graph with n vertices and m edges. Assume that this graph consist of two cliques H_1, H_2 of $n/2$ vertices. Furthermore, each vertex in H_1 has a unique neighbour in H_2 .

Here we would like to use critical blocks to find a good lower bound on the rank. However, the proof on the rank [?] highly depends on the following notions:

A Transition block T is a substring of the sequence such that every vertex in T appear more than once in the sequence.

A Singleton block S is a substring of the sequence such that every vertex in S appear only once in the sequence.

Without repeating their proof here, we give a general explanation of it. The sequence is separated into Transition blocks and Singleton blocks in alternance. We pick a vertex v that appear in T_i and in T_j for $j > i$ and a vertex w appearing in a Singleton block in between, the row of the edge vw is independent of many other rows. This helps finding a bound on the rank depending of s_1 which is hard for arbitrary graphs. Along with a bound on the rank depending on s_2 which exists for arbitrary graphs and the criticality of the block, they manage to find this lower bound on the rank.

However, in the case of two cliques with a matching, a critical block may

contains vertices from both cliques and there is no guarantee that a vertex present in two transition block will have an edge to a singleton vertex in between, (e.g $S_1(B) \subset H_1$ and $S_2(B) \subset H_2$).

Since there are edges between the cliques, we cannot use a reordering argument like in the proof in the previous section. Indeed, reordering the moves could lead to illegal moves where the improvement is negative. Proving then that a move is not in $[0, \epsilon]$ would not suffice to say that the whole sequence is not ϵ -slowly improving since it could be cancelled by another move.

Denote $\alpha(v_t)_{E_i}$ the improvement made by a move, considering only the edges in $E_i \subseteq E$, similarly denote X_{E_i} the restriction of X (the matrix of the weights indexed by the rows) to the rows corresponding only to E_i

Lemma 5.1 *Let $\alpha(v_t)$ be a move and let E_1, E_2 be a partition on the edges touching the vertex v .*

The probability that $\alpha(v_t)$ is in $[0, \epsilon]$ is less or equal than the probability that $\alpha(v_t)_{E_1} \in \bigcup_{u \in \{-1, 1\}^{E_2}} [\langle u, X_{E_2} \rangle, \langle u, X_{E_2} \rangle + \epsilon]$

The proof follows from the fact that $\alpha(v_t) = \alpha(v_t)_{E_1} + \alpha(v_t)_{E_2}$, and that $\alpha(v_t)_{E_2}$ cannot take values outside $\bigcup_{u \in \{-1, 1\}^{E_2}} \{\langle u, X_{E_2} \rangle\}$