ME488 Homework 1

Kaleb Rice

Chapter 2 Problems

2.1 Basic descriptive statistics / standard error / variance

2.7 Normality + one-sample inference

2.10 Interpreting hypothesis-test output

2.15 Two-sample t-test output and interpretation

2.19 One-sample t-test in an engineering context

2.20 Normality assumption and robustness

2.21 Full one-sample t-test + interpretation

2.23 Two-sample comparison with known, unequal σ's

2.25 Two-sample inference / practical interpretation

2.29 Two-sample engineering experiment

Problem 2.1

Problem Statement:

Computer output for a random sample of data is shown below. Some of the quantities are missing. Compute the values of the missing quantities.

Variable	N	Mean	SE Mean	Std. Dev.	Variance	Minimum	Maximum
Y	9	19.96	NA	3.12	NA	15.94	27.16

Solution:

```
# Create table of data from problem statement
variableName <- c("Y")
nn <- c(9)
meanResult <- c(19.96)
SEMeanResult <- c(NA) # Solve for this
stdDevResult <- c(3.12)
varianceResult <- c(NA) # Solve for this
minResult <- c(15.94)
```

```
maxResult <- c(27.16)

# Put variables into data frame
p1Data <- data.frame(aa = variableName,
                     bb = nn,
                     cc = meanResult,
                     dd = SEMeanResult,
                     ee = stdDevResult,
                     ff = varianceResult,
                     gg = minResult,
                     hh = maxResult)
colnames(p1Data)  <-  c("Variable",  "N",  "Mean",  "SE  Mean",  "Std.  Dev.",
"Variance", "Minimum", "Maximum")

# Calculate SE Mean and variance
SEM <- stdDevResult/sqrt(nn)
variance <- stdDevResult^2

# Fill in table values and output results
p1Data[4] <- SEM
p1Data[6] <- variance
knitr::kable(p1Data, format = "pipe", align = 'c')
```

Variable	N	Mean	SE Mean	Std. Dev.	Variance	Minimum	Maximum
Y	9	19.96	1.04	3.12	9.7344	15.94	27.16

Problem 2.7

Problem Statement:

Consider the following sample data:


```
9.37, 13.04, 11.69, 8.21, 11.18, 10.41, 13.15, 11.51, 13.21, 7.75
```

Is it reasonable to assume that this data is a sample from a normal distribution? Is there evidence to support a claim that the mean of the population is 10?

Solution:

To test if the data is normally distributed, I will make a normal QQ plot to see if the data points lie in normal quantiles.

```r
p7Data = c(9.37, 13.04, 11.69, 8.21, 11.18, 10.41, 13.15, 11.51, 13.21, 7.75)
qqnorm(p7Data)
qqline(p7Data)
```



Plotting the normal QQ plot suggests that the data set is normally distributed, although the 13.21 data point is a slight outlier. Continuing with the assumption that the data is normally distributed, I will run a one-sample two-sided t-test with a standard 95% confidence interval in order to test if the mean of 10 lies inside the normally distributed data.

```r
t.test(p7Data, mu = 10) # Defaults to two-sided 95% confidence test
```

```
    One Sample t-test

data:  p7Data
t = 1.5102, df = 9, p-value = 0.1653
alternative hypothesis: true mean is not equal to 10
95 percent confidence interval:
  9.525956 12.378044
sample estimates:
mean of x
   10.952
```

The t-test tells us that there is a 95% chance that the mean is between 9.52 and 12.37, which is statistically significant enough to reject the alternative hypothesis that the mean is not equal to 10.

Conclusion:

It is statistically likely that the data set is normally distributed and that the mean does not differ from 10.

Problem 2.10

Problem Statement:

A two-sample t-test has been conducted and the sample sizes are $n_1 = n_2 = 10$. The computed value of the test statistic is to $t_0 = 2.15$. If the null hypothesis is two-sided, an upper bound on the P-value is:

(a) 0.10 (b) 0.05 (c) 0.025 (d) 0.01 (e) None of the above

Solution:

First I want to find out what P-value is associated with t_0. Since each sample has $n - 1$ degrees of freedom, the combined degrees of freedom for the two sample t-test is 18. I can then use R's distribution function to determine the P-value of t_0 approached from the upper tail.

```
pt(2.15, df = 18, lower.tail = FALSE)
```

```
[1] 0.02269846
```

This function outputs the associated P-value as 0.0227. The nearest upper bound from the list of answers in the problem statement is 0.025. Testing this P-value with R's quantile function yields the following t-value:

```
qt(0.025, df = 18, lower.tail = FALSE)
```

```
[1] 2.100922
```

As expected, this t-value is lower than t_0 because $P = 0.025$ contains a slightly smaller portion of the distribution.

Conclusion:

`(c) 0.025` is the most reasonable upper bound for the P-value.

Problem 2.15

Problem Statement:

Two-Sample t-Test and Cl: Y1, Y2

```
Two-sample T for Y1 vs Y2
Mean Std. Dev. SE Mean
```

```
Y1 20 50.19 1.71 0.38
Y2 20 52.52 2.48 0.55
Difference = mu (X1) - mu (X2)
Estimate for difference: -2.33341
95% CI for difference: (-3.69547, -0.97135)
T-Test of difference = 0 (vs not =): T-Value = -3.47
P-Value = 0.001 DF = 38
Both use Pooled Std. Dev. = 2.1277
```

(a) Can the null hypothesis be rejected at the 0.05 level? Why?

(b) Is this a one-sided or a two-sided test?

(c) If the hypotheses had been $H_0 : \mu_1 - \mu_2 = 2$ versus $H_1 : \mu_1 - \mu_2 \neq 2$, would you reject the null hypothesis at the 0.05 level?

(d) If the hypotheses had been $H_0 : \mu_1 - \mu_2 = 2$ versus $H_1 : \mu_1 - \mu_2 < 2$, would you reject the null hypothesis at the 0.05 level? Can you answer this question without doing any additional calculations? Why?

(e) Use the output and the t table to find a 95 percent upper confidence bound on the difference in means.

(f) What is the P-value if the hypotheses are $H_0 : \mu_1 - \mu_2 = 2$ versus $H_1 : \mu_1 - \mu_2 \neq 2$?

Problem 2.19

Problem Statement:

The shelf life of a carbonated beverage is of interest. Ten bottles are randomly selected and tested, and the following results are obtained:

```
Shelf Life (days)
108, 124, 124, 106, 115, 138, 163, 159, 134, 139
```

(a) We would like to demonstrate that the mean shelf lifeexceeds 120 days. Set up appropriate hypotheses for investigating this claim.

(b) Test these hypotheses using a = 0.01. What are your conclusions?

(c) Find the P-value for the test in part (b).

(d) Construct a 99 percent confidence interval on the meanshelf life.

Problem 2.20

Problem Statement:

Consider the shelf life data in Problem 2.19. Can shelf life be a described or modeled adequately by a normal distribution? What effect would the violation of this assumption have on the test procedure you used in solving Problem 2.15?

Problem 2.21

Problem Statement:

The time to repair an electronic instrument is a normally distributed variable measured in hours. The repair times for 16 such instruments chosen at random are as follows:

(a) You wish to know if the mean repair time exceeds 225 hours. Set up appropriate hypotheses for investigating this issue.

(b) Test the hypotheses you formulated in part (a). What are your conclusions? Use $\alpha = 0.05$.

(c) Find the P-value for the test.

(d) Construct a 95% confidence interval on mean repair time.

```
Hours
159, 224, 222, 149, 280, 379, 362, 260, 101, 179, 168, 485, 212, 264, 250, 170
```

Problem 2.23

Problem Statement:

Two machines are used for filling plastic bottles with a net volume of 16.0 ounces. The filling processes can be assumed to be normal, with standard deviations of $\sigma_1 = 0.015$ and $\sigma_2 = 0.018$. The quality engineering department suspects that both machines fill to the same net volume, whether or not this volume is 16.0 ounces. An experiment is performed by taking a random sample form the output of each machine.

(a) State the hypotheses that should be tested in this experiment.

(b) Test these hypotheses using $\alpha = 0.05$. What are your conclusions?

(c) Find the P-Value for this test.

(d) Find a 95% confidence interval on the difference in mean fill volume for the two machines.

Machine 1	Machine 2
16.03	16.02
16.04	15.97
16.05	15.96
16.05	16.01
16.02	15.99

Machine 1	Machine 2
16.01	16.03
15.96	16.04
15.98	16.02
16.02	16.01
15.99	16.00

Problem 2.25

Problem Statement:

The following are the burning times (in minutes) of chemical flares of two different formulations. The design engineer are interested in both the mean and variance of the burning times.

(a) Test the hypothesis that the two variances are equal. Use $\alpha = 0.05$.

(b) Using the results of (a), test the hypothesis that the mean burning times are equal. Use $\alpha = 0.05$. What is the P-value for this test?

(c) Discuss the role of the normality assumption in this problem. Check the assumption of normality for both types of flares.

Type 1	Type 2
65	64
81	71
57	83
66	59
82	65
82	56
67	69
59	74
75	82
70	79

Problem 2.29

Problem Statement:

The diameter of a ball bearing was measured by 12 inspectors, each using two different kinds of calipers. The results are as follows:

(a) Is there a significant difference between the means of the population of measurements from which the two samples were selected? Use $\alpha = 0.05$.

(b) Find the P-value for the test in part (a).

(c) Construct a 95% confidence interval on the difference in mean diameter measurements for the two types of calipers.

Caliper 1	Caliper 2
0.265	0.264
0.265	0.265
0.266	0.264
0.267	0.266
0.267	0.267
0.265	0.268
0.267	0.264
0.267	0.265
0.265	0.265
0.268	0.267
0.268	0.268
0.265	0.269