



# Irradiated three-dimensional Luttinger semimetal: A factory for engineering Weyl semimetals

**Srimaya Sekhar Padhi, 2011166**  
**Kaling Vikram Singh, 2011074**

**4th Year Integrated MSc., Batch-2020**

**National Institute Of Science Education And Research(NISER), Bhubaneswar**



**Course Instructor - Dr.Kush Saha**

# Contents

1. Band Structure and its Importance
2. Topology and Connection to Band Structure
3. Topological Semimetals
4. Luttinger Hamiltonian and Semimetals
5. Floquet Theory
6. Formalism to Study Irradiation of Luttinger Semimetals
7. Effects of Circularly Polarized Light on Luttinger Semimetals
8. Conclusion

# Band Structure

- Mathematical representation of all occupied and unoccupied energy levels in the lattice of a crystal
- Indispensable apparatus in study of properties of a crystalline solid
- Two types of bands: Valence and Conduction band




1

<sup>1</sup>Solid State Physics - University of Cambridge Part II Mathematical Tripos  
by David Tong

# Classification of Solids













- Classification the basis of gap between valence and conduction band
- 4 types of solids: Metal, semimetal, semiconductors and insulators




- Properties of spaces that are invariant under continuous deformations
- Genus( $g$ ) - Simply, number of holes present on the surface
- **Gauss-Bonnet Theorem:**

$$\int_S \kappa dA = 4\pi(1 - g)$$


# Some Illustrations

Object		Topologically Equivalent	Topologically Non-equivalent
			
			
			

# Connection of Topology to Band Structure




## Types of Materials Based on Topology





# Topological Semimetals

- Topologically nontrivial gapless materials where the conduction and the valence bands cross in the Brillouin zone and the crossing cannot be removed by perturbations preserving concerned crystalline symmetries
- Protected band crossings are present near the Fermi energy and exhibit nonzero topological charges
- The topological charges are basically integer invariants calculated around lines or spheres surrounding the nodes




5

<sup>5</sup>Momentum and Real-Space Study of Topological Semimetals and Topological Defects by Haim Beidenkopf

# Weyl Semimetal

- Semimetals whose low-energy excitations are the Weyl fermions
- Weyl fermions are massless particles proposed by Hermann Weyl in 1929 but discovered very recently in condensed matter systems
- Band dispersion of Weyl semimetal is linear in 3D momentum space through a node known as Weyl point
- The Berry curvature becomes singular at Weyl points that act as monopoles in the momentum space with a fixed chirality, i.e., a source ("+" chirality) or a sink ("- " chirality)
- Weyl points always appear in pairs to avoid the divergence of Berry flux
- The total Berry phase in the 2D  $k$  plane between a pair of Weyl points gives nonzero Chern number while it is zero for other 2D planes



# Types of Weyl Semimetals






Figure: Type I(left) and Type II(right) Weyl Semimetal

# Types OF Weyl Nodes

(a) Type-I Weyl cone



(c) Type-II Weyl cone



8

<sup>8</sup>Quasiparticle interference on type-I and type-II Weyl semimetal surfaces: A review by Hao Zheng and M. Zahid Hasan

# Luttinger-Kohn Hamiltonian and K.P Perturbation Theory

Using Bloch wavefunction in the Schrodinger's equation we get:

$$\left[ \frac{\hbar^2 k^2}{2m} + \frac{\hbar}{m} \vec{k} \vec{p} + \frac{p^2}{2m} + V(r) \right] u_{n,k}(r) = E_n u_{n,k}(r)$$

The k.p term can be taken as the perturbation to the gross Hamiltonian to get the description of heavy and light holes, given by:

$$\frac{1}{m^*} = \frac{1}{m} + \frac{2}{m^2 k^2} \sum_{m \neq n} \frac{\langle u_{n,0} | K \cdot p | u_{m,0} \rangle}{E_{n0} - E_{m0}}$$


Luttinger-Kohn Hamiltonian describes a system with doubly degenerate 4-band structure. Spin orbit coupling breaks the sixfold degenerate  $P_{3/2}$  and  $P_{1/2}$  valence orbitals into 4 high energy and 2 low energy bands, and with further K.P perturbation, the 4 fold degeneracy is lifted to give heavy and light hole bands.

Using group theory and representation theory, one can derive the Luttinger-Kohn Hamiltonian given by:


$$H_{LK} = \frac{1}{2} \int_{\mathbf{k}} c^\dagger(\mathbf{k}) \left( \left( \lambda_1 + \frac{5}{2} \lambda_2 \right) k^2 - 2\lambda_2 (\mathbf{J} \cdot \mathbf{k})^2 - \mu \right) c(\mathbf{k})$$

where  $\alpha$  and  $\beta$  are phases and  $\mathbf{J}$  are spin 3/2 operators.

- The Luttinger-Kohn Hamiltonian incorporates a quadratic band touching, includes spin orbit coupling and relativistic effects which describes the semi metal physics. For our case, the spin orbit coupling and relativistic terms are ignored.
- $\lambda_1, \lambda_2$  are related to the mass of the light and heavy hole in the band structure.
- The bands are doubly degenerate and show bending depending on  $\lambda_i$  values.



$$\lambda_1 = 0.1, \lambda_2 = 0.5$$



$$\lambda_1 = 0.1, \lambda_2 = 0.5$$



# Floquet Theory

- **Floquet Theorem:** Solutions of a periodic linear differential equation:

$$\frac{\partial \vec{x}}{\partial t} = X(t) \vec{x}, X(t+T) = X(t)$$

given by:

$$\vec{x} = \exp(-i\alpha t) \vec{y}(t)$$

where  $\vec{y}(t+T) = \vec{y}(t)$

- For a time periodic Hamiltonian, i.e.,  $H(t) = H(t+T)$ , Floquet-Schrodinger equation is given by:

$$(H(x, t) - i\hbar \frac{\partial}{\partial t}) |\Psi(x, t)\rangle = 0$$


where  $H(x, t) = H_0(x) + V(x, t)$ ,  $V(x, t) = V(x, t+T)$ ,  $T$  is time period of Hamiltonian

- Unitary operator:  
 $U(T) = \mathcal{T} \exp[-i \int_0^T H(t) dt] = \exp[-i H_F T]$ , where  $\mathcal{T}$  is time ordering operator
- $\hat{U}(t_0 + T, t_0) |\Psi_\alpha(t_0)\rangle = \exp(-i \epsilon_n T / \hbar) |\Psi_\alpha(t_0)\rangle$
- $\exp(\frac{-i}{\hbar} T \hat{H}_{t_0}^F) = \hat{U}(t_0 + T, t_0), \hat{H}_{t_0}^F |u_n(t_0)\rangle = \epsilon_n |u_n(t_0)\rangle$
- $H_{\text{eff}} = H_F = \frac{i}{T} \text{Log}(U)$
- Eigenvalues of  $H_F$  are  $\epsilon_\alpha$ , diagonalizing  $\hat{U}(t_0 + T, t_0)$
- $\hat{H}_{t_0}^F$ , acts as  $\hat{H}_{\text{eff}}$ , if observation is stroboscopic, i.e.,  
 $t_0, t_0 + T, t_0 + 2T, \dots, t_0 + nT$
- $H(\mathbf{k}, t) = \sum_m H_m(\mathbf{k}) \exp(im\omega t)$
- $\hat{H}_{\pm m}(\mathbf{k}) = \frac{1}{T} \int_0^T dt \exp(\pm im\omega t) \hat{H}(\mathbf{k}, t), \hat{H}_m = \hat{H}_{-m}^\dagger$


# Floquet Perturbation Theory

- In real systems the Hamiltonian is not time periodic when unperturbed
- Add a time periodic perturbation/drive to the systems(say a sinusoidal vector potential).
- Using different approaches like Floquet-Magnus expansion, degenerate perturbation and other methods to calculate the required parameters
- Different results for different regimes of frequency like high frequency and low frequency regime
- In high frequency regime the effective Hamiltonian:  
$$H_{\text{eff}}(\mathbf{k}) = H_0 + \sum_{n \geq 1} \frac{[H_{+n}, H_{-n}]}{n\omega} + O\left(\frac{1}{\omega^2}\right)$$
 where  $H_0$  is unperturbed Hamiltonian

# Formalism to Study Irradiation of Luttinger Semimetals



- Periodic driving is induced by laser light with a vector potential  $A(t) = (A_x \cos(\omega t), A_y \eta \sin(\omega t), 0)$ , where  $\eta$  - handedness of polarization
- $H(\mathbf{k}, t) \rightarrow H(\mathbf{k} - e\mathbf{A}/\hbar, t)$
- $H_{eff}(k) = H_0(k) + \frac{[H_1, H_{-1}]}{\omega} + \frac{[H_2, H_{-2}]}{2\omega}$
- $H_1 = (\lambda_1 + \frac{5}{2}\lambda_2)\mathbf{k} \cdot \mathbf{A} - 2\lambda_2\{\mathbf{J} \cdot \mathbf{k}, \mathbf{J} \cdot \mathbf{A}\}$ ,  
 $H_2 = \frac{1}{4}[(\lambda_1 + \frac{5}{2}\lambda_2)\mathbf{A}^2 - 2\lambda_2(\mathbf{J} \cdot \mathbf{A})^2]$
- Floquet perturbation series controlled by  $\gamma = \lambda e^2 E^2 / \hbar \omega^3$ , where  $\lambda = \lambda_1 \text{ or } \lambda_2$ .






# Effect of Circularly Polarized Light

- Here, LSM is irradiated by off-resonant circularly polarized light where  $A_x = A_y = A$

- Rotational symmetry ensures appearance of Weyl points appear on only  $k_z$  axis. so taking  $k_x = k_y = 0$  we get:

$$H_{\text{eff}}(k_z) = H_0(k_z) + \frac{2i\eta A^2 \lambda_2^2}{\omega} (-k_z^2 [\{J_x, J_z\}, \{J_y, J_z\}] + \frac{A^2}{8} [J_y^2 - J_x^2, \{J_x, J_y\}])$$

- $E_{1,\pm} = (\lambda_1 + 2\lambda_2)k_z^2 \pm [3A^2\lambda_2^2\eta(A^2 - 8k_z^2)]/2\omega - \mu$   
 $E_{2,\pm} = (\lambda_1 - 2\lambda_2)k_z^2 \pm [3A^2\lambda_2^2\eta(A^2 + 8k_z^2)]/2\omega - \mu$
- In  $H_{\text{eff}}(k_z)$  time reversal symmetry is broken but inversion symmetry is present
- Double degeneracy is lifted and four non degenerate bands appear which intersect in pairs to give rise to Weyl nodes at  $\vec{K}_1 = (0, 0, \pm A/2\sqrt{2})$  and  $\vec{K}_2 = (0, 0, \mp A^2\sqrt{3\lambda_2/\omega})/2$



- $W_n = \frac{1}{8\pi} \int_S d^2k \epsilon^{ijk} \mathbf{n} \cdot (\partial_j \mathbf{n} \times \partial_k \mathbf{n})$  where  $\mathbf{n}$  is a unit vector
- $W_n(K_1) = \pm 1, W_n(K_2) = \pm 2 \implies$  single and double-Weyl nodes co-exist at different energies
- At point  $A_m = \pm \sqrt{\omega/6\lambda_2}$  two types of nodes merge to form triply degenerate point (TDP) .

# Evolution of Luttinger Semimetal to Weyl Semimetal(Bands Bending Opposite Way)




Figure:  $A=2, \lambda_1=0.1, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$






Figure:  $A=2.58, \lambda_1=0.1, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$




Figure:  $A=2.7, \lambda_1=0.1, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$




Figure:  $A=3, \lambda_1=0.1, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$

# Evolution of Luttinger Semimetal to Weyl Semimetal(Bands Bending Same Way)




Figure:  $A=2, \lambda_1=1.8, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$




Figure:  $A=2.58, \lambda_1=1.8, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$




Figure:  $A=2.7, \lambda_1=1.8, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$




Figure:  $A=5, \lambda_1=1.8, \lambda_2=0.5, \omega=20, \mu=0, \eta=1$

# Summary

- Successful conversion of trivial semimetal to a topologically non-trivial semimetal
- Simultaneous accessibility of both types of Weyl nodes through controlled doping and tuning of laser light
- Provides another way to achieve TRS breaking Weyl semimetal using a wide variety of materials
- Adds to the repertoire of types of ways to convert a trivial semimetal to non-trivial topological semimetal(strain-induced[9], magnetic field induced[9], photo-induced[9])
- High scope of experimental realization as Luttinger Hamiltonian describes a wide variety of materials including semiconductors, pyrochlore iridates and half Heuslers which have abundant experimental samples available

10

<sup>10</sup>Engineering topological phases in the Luttinger semimetal  $\alpha - \text{Sn}$  by Dongqin Zhang et al



# THANK YOU FOR YOUR ATTENTION