



Supplementary Material




Supplementary Figure 1. Bootstrap and sample means, including QIs (only for the times the parameter was not set to zero) around the bootstrap mean for edge weights of the primary GGM. The values in the boxes represent the probability of how often the parameter was estimated set to zero.



Supplementary Figure 2. Difference plot of edge weights of the primary GGM. Black squares depict significant different in edge weights ($p < .05$), whereas grey squares illustrate non-significant comparisons ($p > .05$). Comparisons were deemed significant if zero was not in the QIs that was constructed on the difference of the two edge weights. Green dots highlight the most relevant comparisons that are mentioned in the main text.



Supplementary Figure 3. Bootstrap and sample means, including quantile intervals (only for the times the parameter was not set to zero) around the bootstrap mean for edge weights of the primary directed relative importance networks.



Supplementary Figure 4. Difference plot of the directed edge weights of the primary relative importance network. Black squares depict significant difference in the edges ($p < .05$), whereas grey squares illustrate non-significant comparisons ($p > .05$). Dots (green = significant, red = insignificant) highlight the most relevant comparisons.

Supplementary Table 1. Edge weights (r) of the connections among resilience factors in the second GGM.


	BC	PAS	SMA	PHY
BC		0.12	0.28	-0.09
PAS			0.29	0
SMA				0.18

Supplementary Table 2. Edge weights (r) of the connections among QoL facets in the second GGM.


	SAB	AUT	PPF	SOP	DAD	INT
SAB		0	0.06	0	0.07	0
AUT			0.40	0	0	0
PPF				0.48	0.14	0.28
SOP					0	0
DAD						0

Supplementary Table 3. Edge weights (r) of the connections between QoL facets and the resilience factors in the second GGM.


	SAB	AUT	PPF	SOP	DAD	INT
BC	0	0	0	0	0	0.11
PAS	0	0	0.11	0	0.13	0
SMA	0.12	0.10	0	0.24	0	0.11
PHY	0	0	0	0	0	0



Supplementary Figure 5.. Bootstrap and sample means, including quantile intervals (only for the times the parameter was not set to zero) around the bootstrap mean for edge weights of the second GGM. The values in the boxes represent the probability of how often the parameter was estimated set to zero.




Supplementary Figure 6. Difference plot of edge weights of the second GGM. Black squares depict significant difference in edge weights ($p < .05$), whereas grey squares illustrate non-significant comparisons ($p > .05$). Dots (green = significant, red = insignificant) highlight the most relevant comparisons that are mentioned in the main text.



Supplementary Figure 7. Bootstrap and sample means, including quantile intervals (only for the times the parameter was not set to zero) around the bootstrap mean for edge weights of the second

directed relative importance networks. Note that this figure only includes the directed edges that are included in the network, to improve readability.



Supplementary Figure 8. Difference plot of the directed edge weights of the second relative importance network. Black squares depict significant difference in the edges ($p < .05$), whereas grey squares illustrate non-significant comparisons ($p > .05$). Dots (green = significant, red = insignificant) highlight the most relevant comparisons.

Supplementary Table 4. Edge weights (r) of the connections among QoL facets in the third, exploratory GGM.


	SAB	AUT	PPF	SOP	DAD	INT
SAB		0	0.05	0	0.08	0
AUT			0.41	0	0	0
PPF				0.45	0.12	0.25
SOP					0.08	0
DAD						0

Supplementary Table 5. Edge weights (r) of the connections among all resilience factors in the third, exploratory GGM.

	BC	PAS	INI	INV	VAR	MUL	SEF	PFM	PHY
BC		0.15	0.12	0	0.07	0	0.14	0	0
PAS			0	0	0	0.08	0	0.49	0
INI				0.47	0	0.17	0.17	0	0
INV					0.16	0.08	0.15	0.10	0
VAR						0.26	0.14	0	0.18
MUL							0.12	0	0
SEF								0.07	0
PFM									0


Supplementary Table 6. Edge weights (r) of the connections between the resilience factors and QoL facets in the third, exploratory GGM.

	SAB	AUT	PPF	SOP	DAD	INT
BC	0	0	0	0	0	0.09
PAS	0	0	0.1	0	0.12	0
INI	0	0	0	0	-0.06	0
INV	0	0	0	0.12	0	0
VAR	0	0	0	0.06	0	0
MUL	0	0	0	0	0	0.10
SEF	0.09	0	0.13	0.08	0	0.10
PFM	0.07	0.08	0	0	0	0
PHY	0	0	0	0	0	0



Supplementary Figure 9. Bootstrap and sample means, including quantile intervals (only for the times the parameter was not set to zero) around the bootstrap mean for edge weights of the third,

exploratory GGM. The values in the boxes represent the probability of how often the parameter was estimated set to zero. Note that this figure only includes the edges that are included in the network, to improve readability.



Supplementary Figure 10. Difference plot of edge weights of the third, exploratory GGM. Black squares depict significant differences in edge weights ($p < .05$), whereas grey squares illustrate non-significant comparisons ($p > .05$). Dots (green = significant, red = insignificant) highlight the most relevant comparisons that are mentioned in text.