

Supplemental Material

Supplementary Table 1

Demographics for Study Sample and Excluded Sample

Variable		Study Sample <i>M</i> ± <i>SD</i> / Frequency	Excluded sample <i>M</i> ± <i>SD</i> / Frequency	Test Statistic	<i>p</i>
<i>N</i>		707 (62.62%)	422 (37.37%)		
Sex	♀ ♂	476 (67.33%) 231 (32.67%)	272 (64.61%) 149 (35.39%)	$X^2(1) = 0.76^a$	.385
Age		29.20 (8.27)	27.49 (7.17)	$W = 132604^b$	.002
Marital Status	Non-Married Married Separated, Divorced or Widowed	546 (79.48%) 120 (17.47%) 21 (1.16%)	293 (81.62%) 59 (16.43%) 7 (1.94%)	$X^2(2) = 1.361^a$	.506
Number of Persons Living in the Same Household		2.32 (1.22)	2.36 (1.31)	$X^2(5) = 7.245^a$	.203
Employment Status	Full Time Part Time No Employment Student/ in Training	226 (32.85%) 93 (13.52%) 16 (2.32%) 353 (51.31%)	105 (29.17 %) 40 (11.11%) 10 (2.78%) 205 (56.94%)	$X^2(3) = 3.70^a$	.296
Highest Level of Education	Certificate of Secondary Education School-Leaving Examination Completed Vocational Training University Degree	15 (2.18%) 258 (37.50%) 94 (13.66 %) 321 (46.66 %)	14 (3.86%) 150 (41.32%) 50 (13.77 %) 149 (41.05%)	$X^2(3) = 4.9894^a$	.173
Nationality	German Other European Country Others (Asia, South America)	650 (92.36%) 31 (4.39%) 25 (3.54%)	387 (92.36%) 19 (4.53%) 13 (3.10%)	$X^2(2) = 0.164^a$	.921
Mental Health Problems	GHQ	16.67 (7.36)	16.45 (8.45)	$W = 140111^b$	.132
Sense of Coherence	SOC	149.94 (19.53)	150.85 (20.32)	$W = 152138^b$	.341
Life Events	LE	2.35 (1.89)	2.47 (1.76)	$W = 27697^b$	<.001
Daily Hassles	DH	62.45 (28.83)	64.00 (28.69)	$W = 61975^b$	.035

Note. ^aPearson's Chi-squared test, ^bMann-Whitney-U-Test. Regarding missing values, please note that the sum of answers within a variable may not always align with the sample size. Percentages are reported for every category within a subsample and without accounting for missing values. *M* and *SD* for GHQ and SOC were calculated based on the scores of the initial laboratory visit, while for LE and DH, calculations were performed over the observation time window of the trajectories (pre-anchor to post-anchor 6M). Significant differences were observed in both, life events (LE) and daily hassles (DH). This can be attributed to the study sample being composed based on life events and the known correlation between LEs and DHs ($r = .34$). Additionally, there was a significant difference in age, although the effect size was small ($d = 0.22$).

Supplementary Table 2
Demographics for 2-Cluster Separation

Variable		Cluster A <i>M</i> ± <i>SD</i> / Frequency	Cluster B <i>M</i> ± <i>SD</i> / Frequency	Test Statistic	<i>p</i>
<i>n</i>		457 (64.64%)	250 (35.36%)		
Sex	♀	284 (62.14%)	192 (76.80%)	$X^2(1) = 15.119^a$ $W = 59440^b$	<.001 .372
	♂	173 (37.86%)	58 (23.20%)		
Age		29.43 (8.35)	28.79 (8.11)		
Marital Status	Non-Married	349 (78.43%)	197 (81.40%)	$X^2(2) = 3.519$ $X^2(5) = 5.15$	.172 .398
	Married	85 (19.10%)	35 (14.46%)		
	Separated, Divorced or Widowed	11 (2.41%)	10 (4.13%)		
Number of Persons living in the Same Household		2.36 (1.19)	2.26 (1.26)	$X^2(3) = 1.071^a$	.784
	Full Time	152 (34.08%)	74 (30.57 %)		
	Part Time	61 (13.68%)	32 (13.22%)		
	No Employment	10 (2.24 %)	6 (2.48%)		
Employment Status	Student/ in Training	223 (50.00%)	130 (53.72%)	$X^2(3) = 1.175^a$	.759
	Certificate of Secondary Education	9 (2.02%)	6 (2.48%)		
	School-Leaving Examination	162 (36.32%)	96 (39.67%)		
	Completed Vocational Training	64 (14.35%)	30 (12.40%)		
Highest Level of Education	University Degree	211 (47.31%)	110 (45.45%)	$X^2(2) = 0.838^a$	.658
	German	422 (92.54%)	228 (91.20%)		
	Other European Country	20 (4.39%)	11 (4.40 %)		
	Others (Asia, South America)	14 (3.07%)	11 (4.40%)		
Nationality				$W = 37003^b$	<.001
Mental Health Problems	GHQ	15.00 (6.22)	19.72 (8.25)	$W = 77569^b$	<.001
Sense of Coherence	SOC	154.54 (17.40)	141.59 (20.45)	$W = 37085^b$	<.001
Life Events	LE	2.27 (1.85)	2.48 (1.96)	$W = 48636^b$	.262
Daily Hassles	DH	61.10 (28.55)	64.94 (29.19)		.036

Note. ^aPearson's Chi-squared test, ^bMann-Whitney-U test. In some categories, missing values can result in a discrepancy between the cluster's sample size and the total sum of responses within a category. Percentages are reported for every category within a cluster and without accounting for missing values. *M* and *SD* for GHQ and SOC were calculated based on the scores of the initial laboratory visit, while for LE and DH, calculations were performed over the observation time window of the trajectories (pre-anchor to post-anchor 6M).

Supplementary Table 3
Demographics for 5-Cluster Separation

Variable		Cluster A	Cluster B	Cluster C	Cluster D	Cluster E	Test statistic	<i>p</i>
		<i>M</i> ± <i>SD</i> / Frequency	<i>M</i> ± <i>SD</i> / Frequency	<i>M</i> ± <i>SD</i> / Frequency	<i>M</i> ± <i>SD</i> / Frequency	<i>M</i> ± <i>SD</i> / Frequency		
<i>n</i>		215 (30.41%)	169 (23.90%)	140 (19.80%)	92 (13.01%)	91 (12.87%)		
Sex	♀	122 (56.74%)	119 (70.41%)	93 (66.43%)	68 (73.11%)	74 (10.55%)	<i>H</i>(4) = 21.64^a	< .001
	♂	93 (43.29%)	50 (29.59%)	47 (33.57%)	24 (26.09%)	17 (18.68%)		
Age		30.04 (8.41)	28.96 (8.23)	28.96 (8.55)	28.77 (7.84)	28.49 (7.99)	<i>F</i> (1,705) = 2.59 ^b	.108
Marital Status	Non-married	163 (77.62%)	123 (81.99%)	102 (74.45%)	74 (82.22%)	75 (84.27%)	<i>H</i> (4) = 4.53 ^a	.339
	Married	41 (19.52%)	26 (16.15%)	30 (21.90%)	14 (15.56%)	9 (15.56%)		
	Separated, Divorced or Widowed	6 (2.86%)	3 (1.86%)	5 (3.56%)	2 (2.22%)	5 (5.62%)		
Number of Persons Living in the Same Household		2.28 (1.10)	2.42 (1.26)	2.38 (1.26)	2.36 (1.25)	2.12 (1.28)	<i>H</i> (4) = 5.97 ^a	.201
Employment Status	Full Time	81 (38.57%)	45 (27.78%)	43 (31.39%)	31 (34.44%)	26 (29.21%)	<i>H</i> (4) = 5.86 ^a	.210
	Part Time	31 (14.76 %)	23 (14.20%)	16 (11.68%)	12 (13.33%)	11 (12.36%)		
	No Employment	5 (2.38 %)	3 (1.85%)	3 (2.19%)	3 (3.33%)	2 (2.24%)		
	Student/ in Training	93 (44.29%)	91 (56.17%)	75 54.74%)	44 (48.89%)	50 (56.18%)		
Highest Level of Education	Certificate of Secondary Education	6 (2.85%)	2 (1.23%)	2 (1.46%)	3 (3.33%)	2 (2.25%)	<i>H</i> (4) = 2.12 ^a	.714
	School-Leaving Examination	72 (34.29%)	57 (35.19%)	57 (41.61%)	37 (41.11%)	35 (39.33%)		
	Completed Vocational Training	30 (14.29%)	24 (14.81%)	17 (12.41%)	9 (10.00%)	14 (15.73%)		
	University Degree	102 (48.57%)	79 (48.77%)	61 (44.53%)	41 (45.56%)	38 (42.70%)		
Nationality	German	200 (93.02%)	154 (91.12%)	129 (92.81%)	86 (93.48%)	81 (89.48%)	<i>H</i> (4) = 2.14 ^a	.710
	Other European	9 (4.19%)	8 (4.73%)	6 (4.32%)	4 (4.35%)	4 (4.40%)		
	Country							
	Others (Asia, South America)	6 (2.79%)	7 (4.14%)	4 (2.88%)	2 (2.17%)	6 (6.59%)		
Mental Health Problems	GHQ	13.73 (5.62)	16.42 (6.56)	16.62 (6.72)	18.41 (8.04)	22.41 (8.04)	<i>F</i>(1, 703) = 99.23^b	< .001
	SOC	156.38 (17.68)	151.41 (17.05)	150.67 (19.56)	145.62 (17.94)	135.34 (21.20)	<i>F</i>(1, 702) = 80.27^b	< .001
Life Events	LE	2.37 (1.89)	2.11 (1.75)	2.30 (1.87)	2.45 (1.91)	2.71 (2.12)	<i>F</i> (1, 589) = 1.72 ^b	.190
Daily Hassles	DH	62.73 (29.52)	57.43 (26.36)	63.49 (28.61)	64.82 (28.87)	67.16 (30.68)	<i>F</i> (1, 685) = 3.18 ^b	.075

Note. ^aKruskal-Wallis test, ^bAnalysis of variance. In some categories, missing values can result in a discrepancy between the cluster's sample size and the total sum of responses within a category. Percentages are reported for every category within a cluster and without accounting for missing values. *M* and *SD* for GHQ and SOC were calculated based on the scores of the initial laboratory visit, while for LE and DH, calculations were performed over the observation time window of the trajectories (pre-anchor to post-anchor 6M).

Supplementary Table 4*Fisher's Exact Test - Relationship Between Cluster Membership and Mental Dysfunction*

	Fisher's Exact Test	Pairwise Fisher's Test		
	<i>p</i>	Cluster	Cluster	<i>p</i> _{adj}
5-Cluster	<.001	A	B	1
		A	C	.995
		A	D	.005
		A	E	.021
		B	C	1
		B	D	.033
		B	E	.169
		C	D	.773
		C	E	1
		D	E	1

Note. *N* = 628. Fisher's exact test as post-hoc test to compare each cluster combination with p-values, corrected using the Bonferroni correction.

Supplementary Table 5*Relationship Between Clusters and Mental Dysfunction*

	Variable	<i>b</i>	<i>p</i>	<i>OR</i>	95% <i>CI</i>
2-Cluster Separation	Intercept	-2.90	<.001	0.06	0.03; 0.10
	Cluster (B)	0.68	.004	1.97	1.24; 3.12
	Number of MINI interviews	0.46	.003	1.59	1.17; 2.16
	Sex (female)	0.91	.003	2.48	1.38; 4.48
	Age	-0.03	.797	0.97	0.78; 1.21
5-Cluster Separation	Intercept	-3.10	<.001	0.04	0.02; 0.09
	Cluster (B)	0.15	.067	1.16	0.55; 2.43
	Cluster (C)	0.58	.116	1.78	0.87; 3.66
	Cluster (D)	1.18	.001	3.27	1.59; 6.73
	Cluster (E)	1.09	.005	2.98	1.40; 6.36
	Number of MINI interviews	0.47	.003	1.61	1.18; 2.19
	Sex (female)	0.86	.005	2.37	1.30; 4.29
	Age	-0.02	.866	0.98	0.78; 1.23

Note. *N* = 628. Dependent variable is mental dysfunction. Cluster A serves as reference cluster.

Supplementary Table 6*Clusters and Mental Dysfunction Controlling for Pre-Anchor SR*

	Variable	<i>b</i>	<i>p</i>	<i>OR</i>	95% <i>CI</i>
2-Cluster Separation	Intercept	-2.83	<.001	0.06	0.03; 0.10
	Cluster (B)	0.50	.061	1.65	0.98; 2.80
	Number of MINI interviews	0.47	.003	1.60	1.17; 2.17
	Sex (female)	0.90	.003	2.46	1.36; 4.44
	Age	-0.04	.759	0.97	0.77; 1.21
	Pre anchor SR	0.17	.182	1.19	0.92; 1.52
5-Cluster Separation	Intercept	-3.03	<.001	0.05	0.02; 0.10
	Cluster (B)	0.09	.806	1.10	0.52; 2.33
	Cluster (C)	0.46	.249	1.59	0.72; 3.48
	Cluster (D)	1.08	.006	2.94	1.35; 6.39
	Cluster (E)	0.91	.043	2.51	1.03; 6.09
	Number of MINI interviews	0.47	.002	1.61	1.18; 2.19
	Sex (female)	0.86	.004	2.37	1.30; 4.29
	Age	-0.02	.832	0.98	0.78; 1.22
	Pre-anchor SR	0.10	.463	1.11	0.85; 1.44

Note. *N* = 628. Dependent variable is mental dysfunction. Cluster A serves as reference cluster.

Growth Mixture Modeling (GMM)

In order to establish comparability with the commonly used method in resilience research, (Latent) Growth Mixture Modeling (GMM, Muthén & Muthén, 2000), and the method employed in this paper, k-means for longitudinal data (KML, Genolini et al., 2015; Genolini & Falissard, 2011) for identifying differential trajectories, the results of applying GMM to the dataset are presented here.

GMM Method: Growth Mixture Models calculate a mean growth curve for each class, with variances estimated around the growth parameters within each latent trajectory (Muthén & Muthén, 2000). Thus, GMM account for the heterogeneity in variation within each class. GMMs were computed using the R package *lcmm* (Proust-Lima et al., 2017). To model the response to the anchor life event, a quadratic trend was chosen. To avoid local maxima, 50 iterations were conducted and 250 random vectors of initial values were employed.

As recommended by van De Schoot et al. (2017), we gave the greatest emphasis to the Bayesian Information Criterion (BIC) as the primary fit index. Additionally, following the suggestions of Wickrama et al. (2016), only classes were considered as meaningful, that include more than 5% of the participants.

GMM Results: The resulting class separations are illustrated in Supplementary Figure 1. Considering the fit indices (as presented in Supplementary Table 7) and taking into account the size of the smallest classes, the 3-class separation was chosen.

Comparison between results of GMM and KML: The Adjusted Rand Index (ARI; Hubert & Arabie, 1985) was used to compare the agreement between the classifications of GMM with the classifications of KML, while 1 indicates similar clustering. While the two-class separation, in particular, exhibits the most significant overlap with the 2-cluster separation generated by KML (agreement = 94%, also indicated by ARI = .77). The other cluster/class separations differ markedly from the KML solutions (ARI = .41 for the 3-cluster/class separation, ARI = .19 for the 4-cluster/class separation, ARI = .19 for the 5-cluster separation).

Supplementary Table 7 provides an overview of the underlying data within the clusters. Compared to the preferred 5-cluster solution derived from the KML approach, as depicted in Figure 3 in the main text, it appears that the KML clusters provide a better description of the underlying data. It is important to note that this should not be interpreted as a general preference for KML. Rather, for this specific dataset and considering the exploratory nature of our analyses, we favor the use of KML for these analyses.

A comprehensive discussion of the advantages and disadvantages of GMM and KML can be found in the manuscript in section “Limitations of clustering approaches: KML vs. GMM”.

Supplementary Table 7*Fit Indices of Growth Mixture Models*

	Loglik	AIC ^a	BIC ^b	SABIC ^c	Entropy ^d	ICL ^e	% Class 1	% Class 2	% Class 3	% Class 4	% Class 5
2-Classes	-3784.696	7599.393	7667.808	7620.180	0.403	6512.528	65.77	34.23			
3-Classes	-3768.279	7576.558	7667.779	7604.274	0.503	6586.956	40.74	47.52	11.74		
4-Classes	-3763.622	7577.244	7691.270	7611.889	0.488	6708.107	14.57	51.06	30.27	4.10	
5-Classes	-3757.990	7575.979	7712.810	7617.554	0.605	6685.498	13.01	53.47	27.16	1.70	4.67

Note. Fit indices that should be maximized: Loglik = Maximum log-likelihood. Entropy (should be close to 1). Fit indices that should be minimized: AIC, BIC, SABIC, ICL. ^aAkaike (1974) ^bSchwarz (1978), ^cSolve (1987), ^dCeleux et al. (1996), ^eBiernacki et al. (2000).

Supplementary Table 8*Fisher's Exact Test - Relationship Between Class Membership (GMM) and Mental Dysfunction*

	Fisher's Exact Test	Pairwise Fisher's Test		
	<i>p</i>	Cluster	Cluster	<i>p</i> _{adj}
3-Classes	<.001	A	B	.144
		A	C	<.001
		B	C	<.001

Note. *N* = 628. Fisher's exact test as post-hoc test to compare each class combination with *p*-values, corrected using the Bonferroni correction.

Supplementary Table 9*Classes (GMM) and Mental Dysfunction*

	Variable	<i>b</i>	<i>p</i>	OR	95% CI
3-Classes Separation	Intercept	-3.10	<.001	0.05	0.02; 0.09
	Cluster (B)	0.48	.087	1.62	0.93; 2.82
	Cluster (C)	1.62	<.001	5.05	2.58; 9.87
	Number of MINI interviews	0.48	.002	1.62	1.19; 2.22
	Sex (female)	0.87	.004	2.38	1.31; 4.32
	Age	-0.04	.723	0.96	0.77; 1.20

Note. *N* = 628. Dependent variable is mental dysfunction. Class A serves as reference class.

Supplementary Table 10*Classes (GMM) and Mental Dysfunction Controlling for Pre-Anchor SR*

	Variable	<i>b</i>	<i>p</i>	OR	95% CI
3-Classes Separation	Intercept	-3.11	<.001	0.04	0.02; 0.09
	Cluster (B)	0.50	.108	1.64	0.90; 3.01
	Cluster (C)	1.65	<.001	5.20	2.26; 11.93
	Number of MINI interviews	0.48	.002	1.62	1.19; 2.22
	Sex (female)	0.87	.004	2.38	1.31; 4.32
	Age	-0.04	.729	0.96	0.77; 1.21
	Pre-anchor SR	-0.02	.907	0.98	0.75; 1.29


Note. *N* = 628. Dependent variable is mental dysfunction. Class A serves as reference class.

Supplementary Table 11
Construct Validity of Classes (GMM)


3-Classes Separation	Variable	<i>b</i>	<i>p</i>	<i>OR</i>	95% <i>CI</i>
				A vs. B	
	Intercept	0.17	.335	1.19	0.84; 1.68
	SOC	-0.24	.036	0.78	0.63; 0.98
	Sex (female)	0.69	.001	1.99	1.31; 3.04
	Age	-0.18	.110	0.84	0.67; 1.04
	Pre-Anchor SR	1.79	<.001	5.97	4.29; 8.31
	Life-Time LE Exposure	-0.06	.584	0.94	0.75; 1.17
	LEs Between the Initial Laboratory Visit and Pre-Anchor	0.08	.419	1.09	0.89; 1.34
				A vs. C	
	Intercept	-2.75	<.001	0.06	0.03; 0.13
	SOC	-0.94	<.001	0.39	0.28; 0.56
	Sex (female)	1.31	<.001	3.69	1.74; 7.83
	Age	0.12	.514	1.12	0.79; 1.60
	Pre Anchor SR	2.88	<.001	17.73	11.16; 28.18
	Life-Time LE Exposure	-0.30	.119	0.74	0.50; 1.08
	LEs Between the Initial Laboratory Visit and Pre-Anchor	0.19	.246	1.21	0.88; 1.66

Note. *N* = 675.

Supplementary Figure 1 *Resulting Classes Based on Growth Mixture Models*




Supplementary Figure 2
Class Separation Preferred by GMM



Note. Classes do not fit the underlying data as good as clusters identified by KML algorithm (see Figure 3).


Supplementary Figure 3
Resulting Clusters After Applying the KML Algorithm on Raw Scores of Mental Health Problems



Note. Mental health problems measured with GHQ-28. For a discussion of the differences, see main manuscript section “SR vs. Raw Scores of Mental Health Problems in Trajectory Modeling”.

Supplementary Figure 4

Life Event Categories for 2-Cluster and 5-cluster Separation



Note. Frequency refers to the proportion of life events that can be assigned to this category in each cluster. Statistical tests conducted separately for each of the 27 life events showed that life events do not significantly differ between clusters ($p > .056$), except for one life event within the 2-cluster solution (financial problems: $p = .049$).

R-Code to compute the Stressor Reactivity (SR) score

```
##### data preparing #####
# read sav-file
raw_data <- haven::read_sav("data.sav", encoding="latin1")

# scale variables
data_scaled <- scale(raw_data)

# Convert dataframe to long format
data_long <- data_scaled %>% tidyr::pivot_longer(
  names_to = c(".value", "time"),
  names_sep = "\\_")

##### calculate stressor exposure #####
# Data exclusion if life events and daily hassles are missing
data <- data_long %>%
  dplyr::filter(is.na(le) == FALSE | is.na(dh) == FALSE)

# Calculate combined stressor exposure
data$E <- rowMeans(cbind(data$dh, data$le), na.rm = TRUE)

# scale stressor exposure
data$E <- scale(data$E)

##### calculate stressor reactivity #####
# check model assumptions
cor.test(data$le, data$dh)
PlotQQ(data$E)
plot(ghq ~ E, data)

# multilevel model with random slope and random intercept
mlm <- lmer(ghq ~ E + (E | id), data)
summary(mlm)

# fixed effects
intercept <- summary(mlm)$coefficients[1]
slope <- summary(mlm)$coefficients[2]

# predicted values
data$pred <- (intercept + (data$E*slope))

# residuals = Stressor Reactivity Score
data$SR <- as.numeric(scale(data$ghq - pred))

# save sav-file
haven::write_sav(data, "SR.sav")
```

Supplementary References

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE transactions on automatic control*, 19(6), 716-723.
- Biernacki, C., Celeux, G., & Govaert, G. (2000). Assessing a mixture model for clustering with the integrated completed likelihood. *IEEE transactions on pattern analysis and machine intelligence*, 22(7), 719-725.
- Celeux, G., & Soromenho, G. (1996). An entropy criterion for assessing the number of clusters in a mixture model. *Journal of classification*, 13, 195-212.
- Genolini, C., Alacoque, X., Sentenac, M., & Arnaud, C. (2015). kml and kml3d: R Packages to Cluster Longitudinal Data. *Journal of Statistical Software*, 65(4), 1–34. <https://doi.org/10.18637/JSS.V065.I04>
- Genolini, C., & Falissard, B. (2011). KmL: a package to cluster longitudinal data. *Computer Methods and Programs in Biomedicine*, 104(3). <https://doi.org/10.1016/J.CMPB.2011.05.008>
- Hubert, L., & Arabie, P. (1985). Comparing partitions. *Journal of Classification*, 2(1), 193–218. <https://doi.org/10.1007/BF01908075>
- Muthén, B., & Muthén, L. K. (2000). Integrating Person-Centered and Variable-Centered Analyses: Growth Mixture Modeling With Latent Trajectory Classes. *Alcoholism: Clinical and Experimental Research*, 24(6), 882–891. <https://doi.org/10.1111/j.1530-0277.2000.tb02070.x>
- Proust-Lima, C., Philipps, V., & Lique, B. (2017). Estimation of Extended Mixed Models Using Latent Classes and Latent Processes: The R Package lcmm. *Journal of Statistical Software*, 78, 1–56. <https://doi.org/10.18637/jss.v078.i02>
- Schwarz, G. (1978). Estimating the dimension of a model. *The annals of statistics*, 461-464.
- Sclove, S. L. (1987). Application of model-selection criteria to some problems in multivariate analysis. *Psychometrika*, 52, 333-343.
- van De Schoot, R., Sijbrandij, M., Winter, S. D., Depaoli, S., & Vermunt, J. K. (2017). The GRoLTS-Checklist: Guidelines for Reporting on Latent Trajectory Studies. *Structural Equation Modeling: A Multidisciplinary Journal*, 24(3), 451–467. <https://doi.org/10.1080/10705511.2016.1247646>
- Wickrama, K., Lee, T. K., O’Neal, C. W., & Lorenz, F. (2016). *Higher-Order Growth Curves and Mixture Modeling with Mplus: A Practical Guide* (1st ed.). Routledge.