Crop Disease Detection Report

Generated on March 24, 2026 | 07:35 PM

Disease Detected	Tomato - Early Blight
Confidence Score	90.8%
Confidence Level	High
Analysis Status	Complete

Analyzed Leaf Image

Symptoms

Dark brown spots with concentric rings (target-like pattern) on older leaves, yellowing and dropping of lower leaves, lesions on stems and fruits, leaves curl and die

Causes

Caused by Alternaria solani fungus, favored by warm temperatures (75-85°F), high humidity, wet foliage, poor air circulation, stressed plants

Treatment Methods

Remove infected lower leaves, apply fungicides containing chlorothalonil or mancozeb, use copper-based fungicides, ensure proper plant nutrition

Preventive Measures

Mulch around plants to prevent soil splash, water at base of plants, stake plants for air circulation, practice 3-year crop rotation, use resistant varieties

Recommended Pesticides

Chlorothalonil (Daconil), Mancozeb (Dithane), Azoxystrobin (Quadris), Copper fungicides, Apply every 7-14 days preventively

Organic Solutions

Baking soda spray (1 tbsp per gallon), Compost tea foliar spray, Copper fungicide, Sulfur dust, Remove affected leaves promptly

This report was generated automatically by the Crop Disease Detection System. Please consult an agricultural expert for professional advice.