

Abstract nonsense?

Kalle Heinonen

Aalto university (sivari), Moodi ry

Integraatiofest 2025

Abstract nonsense?



No you can't use abstract nonsense in functional analysis



haha splitting lemma go
brr... $L = \text{im}T + \text{ker}T$

in any **abelian category**, the following statements are **equivalent**: for a **short exact sequence**



$$0 \longrightarrow A \xrightarrow{q} B \xrightarrow{r} C \longrightarrow 0.$$

Then the following statements are equivalent:

- There exists a morphism $u: C \rightarrow B$ such that ru is the identity on C , id_C .
- There is an isomorphism h from B to the **direct sum** of A and C , such that hq is the natural monomorphism of A in the direct sum, and rh^{-1} is the natural projection of the direct sum onto C .

Origins in topology

How do we know these spaces are not homeomorphic formally?



Definition

A **category \mathbf{C}** consists of

- **Objects** A, B, C, \dots

Definition

A **category \mathbf{C}** consists of

- **Objects** A, B, C, \dots
- **Morphisms** or **arrows** between objects (a morphism $f: A \rightarrow B$ with domain A , and codomain B).

Definition

A **category C** consists of

- **Objects** A, B, C, \dots
- **Morphisms** or **arrows** between objects (a morphism $f: A \rightarrow B$ with domain A , and codomain B).
- Morphisms are **composable** meaning if $g: B \rightarrow C$, then $g \circ f: A \rightarrow C$ is a morphism where \circ is associative.

Definition

A **category \mathbf{C}** consists of

- **Objects** A, B, C, \dots
- **Morphisms** or **arrows** between objects (a morphism $f: A \rightarrow B$ with domain A , and codomain B).
- Morphisms are **composable** meaning if $g: B \rightarrow C$, then $g \circ f: A \rightarrow C$ is a morphism where \circ is associative.
- Every object has an **identity** arrow $1_A: A \rightarrow A$. If a composition of morphisms equals the identity, it is called an **inverse** (left and right inverse respectfully). If it has both inverses, it is an **isomorphism**.

Definition

A **category \mathbf{C}** consists of

- **Objects** A, B, C, \dots
- **Morphisms** or **arrows** between objects (a morphism $f: A \rightarrow B$ with domain A , and codomain B).
- Morphisms are **composable** meaning if $g: B \rightarrow C$, then $g \circ f: A \rightarrow C$ is a morphism where \circ is associative.
- Every object has an **identity** arrow $1_A: A \rightarrow A$. If a composition of morphisms equals the identity, it is called an **inverse** (left and right inverse respectfully). If it has both inverses, it is an **isomorphism**.

In "regular" mathematics one basically always assumes that our category is **locally small** (as in this presentation) meaning, the morphisms between any two objects $\text{hom}(A, B)$ are of set size.

Definition

A **subcategory** \mathbf{D} of \mathbf{C} is category that includes some of the objects and morphism of \mathbf{C} . A subcategory is **full** if it contains the maximal amount of morphism's with respect ot the objects.


Definition

A **subcategory** \mathbf{D} of \mathbf{C} is category that includes some of the objects and morphism of \mathbf{C} . A subcategory is **full** if it contains the maximal amount of morphism's with respect ot the objects.

We can form a category from any directed multigraph and it's subgraphs will be subcategories

Diagrams of morphisms example

Given two commutative squares,



they make up a commutative rectangle.

Familiar examples of categories

- **Set** of sets, functions, function composition.

Familiar examples of categories

- **Set** of sets, functions, function composition.
- **Grp** of groups, homomorphisms, composition of homomorphisms.

Familiar examples of categories

- **Set** of sets, functions, function composition.
- **Grp** of groups, homomorphisms, composition of homomorphisms.
- **Top** of topological spaces, continuous maps and composition of continuous maps.

Familiar examples of categories

- **Set** of sets, functions, function composition.
- **Grp** of groups, homomorphisms, composition of homomorphisms.
- **Top** of topological spaces, continuous maps and composition of continuous maps.
- **Vect $_{\mathbb{R}}$** of real vector spaces, linear maps, and composition of linear maps.

Familiar examples of categories

- **Set** of sets, functions, function composition.
- **Grp** of groups, homomorphisms, composition of homomorphisms.
- **Top** of topological spaces, continuous maps and composition of continuous maps.
- **Vect $_{\mathbb{R}}$** of real vector spaces, linear maps, and composition of linear maps.

These are something we would call **concrete** categories (i.e. in "some sense" underlying structure is a set and morphism a set theoretical function). A non concrete category would be for example **hTop** which is **Top** but morphisms are homotopy classes of morphisms.

Familiar examples of categories

- **Set** of sets, functions, function composition.
- **Grp** of groups, homomorphisms, composition of homomorphisms.
- **Top** of topological spaces, continuous maps and composition of continuous maps.
- **Vect _{\mathbb{R}}** of real vector spaces, linear maps, and composition of linear maps.

These are something we would call **concrete** categories (i.e. in "some sense" underlying structure is a set and morphism a set theoretical function). A non concrete category would be for example **hTop** which is **Top** but morphisms are homotopy classes of morphisms. A metacategory on the otherhand doesn't care about any underlying set theory. Take the category of Haskell types and functions as an example.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.
- A **discrete** category has objects, but only morphisms are identities.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.
- A **discrete** category has objects, but only morphisms are identities.
- A group G becomes a category $\mathbf{B}G$ with one element, and morphism being the group multiplication.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.
- A **discrete** category has objects, but only morphisms are identities.
- A group G becomes a category $\mathbf{B}G$ with one element, and morphism being the group multiplication.
- The opposite category \mathbf{C}^{op} of \mathbf{C} has same objects, but order of every arrow is reversed ($\text{hom}^{op}(A, B) := \text{hom}(B, A)$).

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.
- A **discrete** category has objects, but only morphisms are identities.
- A group G becomes a category $\mathbf{B}G$ with one element, and morphism being the group multiplication.
- The opposite category \mathbf{C}^{op} of \mathbf{C} has same objects, but order of every arrow is reversed ($\text{hom}^{op}(A, B) := \text{hom}(B, A)$).
- A **poset** (P, \leq) is a category where $A \rightarrow B$ if $A \leq B$ when (P, \leq) is viewed as partially ordered set. For example $\text{Op}(X)$ the category of open sets of a space X , with inclusions as morphisms.

More "funky?" examples (that are actually useful)

- **1** one object, identity morphism.
- **2** two object, one non trivial morphism.
- A **discrete** category has objects, but only morphisms are identities.
- A group G becomes a category $\mathbf{B}G$ with one element, and morphism being the group multiplication.
- The opposite category \mathbf{C}^{op} of \mathbf{C} has same objects, but order of every arrow is reversed ($\text{hom}^{op}(A, B) := \text{hom}(B, A)$).
- A **poset** (P, \leq) is a category where $A \rightarrow B$ if $A \leq B$ when (P, \leq) is viewed as partially ordered set. For example $\text{Op}(X)$ the category of open sets of a space X , with inclusions as morphisms.
- **Cat** the category of small categories (category is **small** if it has set amount of objects). We will soon find out what the morphisms in this category are.

Example of duality

$$C \begin{array}{c} \xrightarrow{a} \\ \xrightarrow{b} \end{array} A \xrightarrow{f} B$$

f is a monomorphism if for all C, a, b

$$C \begin{array}{c} \xleftarrow{b} \\ \xleftarrow{b} \end{array} A \xleftarrow{f} B$$

f is an epimorphism if for all C, a, b

Now if f is a monomorphism in a category \mathbf{C} iff it's associated morphism is an epimorphism in the opposite category \mathbf{C}^{op} .

Diagram chasing example

Suppose the outer most square below commutes.

$$\begin{array}{ccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & F \\ \downarrow g & & \downarrow f & & \downarrow g' \\ B & \xrightarrow{a'} & D & \xrightarrow{b'} & E \end{array}$$

This data defines a commutative rectangle if either the RHS square commutes and b' is a monomorphism or LHS square commutes and a is an epimorphism.

Example of duality 2

Proving a theorem in category theory proves two theorems at once.
A proposition holds in a category \mathbf{C} iff it's dual proposition holds in \mathbf{C}^{op}

Example of duality 2

Proving a theorem in category theory proves two theorems at once. A proposition holds in a category \mathbf{C} iff it's dual proposition holds in \mathbf{C}^{op}

Definition

An object T in a category \mathbf{C} is **initial**, if there exists a unique morphism $T \rightarrow C$ for any other object C in \mathbf{C} . The dual object of terminal object is called **terminal**. An object satisfies an **universal property** if it is initial (or dually terminal) in some ambient category.

Example of duality 2

Proving a theorem in category theory proves two theorems at once. A proposition holds in a category \mathbf{C} iff it's dual proposition holds in \mathbf{C}^{op}

Definition

An object T in a category \mathbf{C} is **initial**, if there exists a unique morphism $T \rightarrow C$ for any other object C in \mathbf{C} . The dual object of terminal object is called **terminal**. An object satisfies an **universal property** if it is initial (or dually terminal) in some ambient category.

Theorem

\mathbf{C} has an initial object iff \mathbf{C}^{op} has a terminal object.


Theorem

Initial objects are unique up to a unique isomorphism.

Visual proof of the second theorem

$$\begin{array}{ccccccc} & & & 1_{T'} & & & \\ & & \nearrow & & \searrow & & \\ T' & \xrightarrow{\exists!} & T & \xrightarrow{\exists!} & T' & \xrightarrow{\exists!} & T \\ & & \searrow & & \nearrow & & \\ & & & 1_T & & & \end{array}$$

**Noooooooo! You can't just
draw a picture and claim
it's a mathematical proof. You
need to rigorously prove the claim
instead!**



Definition

A (covariant) **functor** $F: \mathbf{C} \rightarrow \mathbf{D}$ takes an object in \mathbf{C} and returns an object $F(C)$ in \mathbf{D} . For any morphism $f: C \rightarrow C'$ there is an associated morphism $F(f): F(C) \rightarrow F(C')$ where F preserves the composition of morphisms $F(f \circ g) = F(f) \circ F(g)$ and $F(1_A) = 1_{F(A)}$.

It is easy to see from the composition laws that functors preserve isomorphisms. Functors preserve commutative diagrams up to variance.

Examples of functors

- The dual space functor on vector spaces
 $(-)^*: \mathbf{Vect}_{\mathbb{K}}^{\text{op}} \rightarrow \mathbf{Vect}_{\mathbb{K}}$, where $V \mapsto V^*$ and
 $L: V \rightarrow W \mapsto L^*: W^* \rightarrow V^*$ where the associated morphism is
the precomposition by L . Not that $(gf)^* = f^*g^*$ because the
domain of the functor is an opposite category.

Examples of functors

- The dual space functor on vector spaces
 $(-)^*: \mathbf{Vect}_{\mathbb{K}}^{\text{op}} \rightarrow \mathbf{Vect}_{\mathbb{K}}$, where $V \mapsto V^*$ and
 $L: V \rightarrow W \mapsto L^*: W^* \rightarrow V^*$ where the associated morphism is
the precomposition by L . Not that $(gf)^* = f^*g^*$ because the
domain of the functor is an opposite category.
- A list in Haskell (or other functional programming languages)
behaves functorially by `fmap`

$$\text{fmap } (+1) [1, 2, 3] == [2, 3, 4] .$$

Examples of functors

- The dual space functor on vector spaces
 $(-)^*: \mathbf{Vect}_{\mathbb{K}}^{\text{op}} \rightarrow \mathbf{Vect}_{\mathbb{K}}$, where $V \mapsto V^*$ and
 $L: V \rightarrow W \mapsto L^*: W^* \rightarrow V^*$ where the associated morphism is
the precomposition by L . Not that $(gf)^* = f^*g^*$ because the
domain of the functor is an opposite category.
- A list in Haskell (or other functional programming languages)
behaves functorially by `fmap`
$$\text{fmap } (+1) [1, 2, 3] == [2, 3, 4] .$$
- A forgetful functor (literally forgets structure). Example
`Forget: Ring → Ab` forgets the multiplicative structure.

- The fundamental group $\pi_1(X): \mathbf{hTop}_* \rightarrow \mathbf{Grp}$ is a functor.

Examples

- The fundamental group $\pi_1(X): \mathbf{hTop}_* \rightarrow \mathbf{Grp}$ is a functor.
- The free group is a functor $\mathbf{Free}: \mathbf{Set} \rightarrow \mathbf{Grp}$.

Examples

- The fundamental group $\pi_1(X): \mathbf{hTop}_* \rightarrow \mathbf{Grp}$ is a functor.
- The free group is a functor $\mathbf{Free}: \mathbf{Set} \rightarrow \mathbf{Grp}$.
- The power set functor $P: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ which associates each function, it's preimage on the level of the power set is functorial.

- The fundamental group $\pi_1(X): \mathbf{hTop}_* \rightarrow \mathbf{Grp}$ is a functor.
- The free group is a functor $\mathbf{Free}: \mathbf{Set} \rightarrow \mathbf{Grp}$.
- The power set functor $P: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ which associates each function, it's preimage on the level of the power set is functorial.
- The hom functors $\text{hom}(A, -): \mathbf{C} \rightarrow \mathbf{Set}$ with post composition and $\text{hom}(-, A): \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ with pre composition are one of the most important ones where. For example, the dual space functor is $\text{hom}(-, \mathbb{K}): \mathbf{Vect}_{\mathbb{K}}^{\text{op}} \rightarrow \mathbf{Set}$ if we do not equip the dual space with a vector space structure.

- The fundamental group $\pi_1(X): \mathbf{hTop}_* \rightarrow \mathbf{Grp}$ is a functor.
- The free group is a functor $\mathbf{Free}: \mathbf{Set} \rightarrow \mathbf{Grp}$.
- The power set functor $P: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ which associates each function, it's preimage on the level of the power set is functorial.
- The hom functors $\text{hom}(A, -): \mathbf{C} \rightarrow \mathbf{Set}$ with post composition and $\text{hom}(-, A): \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ with pre composition are one of the most important ones where. For example, the dual space functor is $\text{hom}(-, \mathbb{K}): \mathbf{Vect}_{\mathbb{K}}^{\text{op}} \rightarrow \mathbf{Set}$ if we do not equip the dual space with a vector space structure.
- **Presheaf** on a topological space X is a functor $\text{Op}(X)^{\text{op}} \rightarrow \mathbf{Set}$.

Natural transformations

Definition

Given a categories \mathbf{C} , \mathbf{D} and pair of functors $F, G: \mathbf{C} \Rightarrow \mathbf{D}$ a natural transformation $\eta: F \Rightarrow G$ between the functors is collection of morphisms $\eta_C: F(C) \rightarrow G(C)$ for each object C in \mathbf{C} such that the following square commutes

$$\begin{array}{ccc} F(C) & \xrightarrow{\eta_C} & G(C) \\ F(f) \downarrow & & \downarrow G(f) \\ F(D) & \xrightarrow{\eta_D} & G(D) \end{array}$$

If each of the components of the natural transformation is an isomorphism we call it a natural isomorphism. Heuristically this is an isomorphism with "lack of arbitrary choices".

Natural transformations

- For any vector space $V \cong V^{**}$ naturally by evaluation. If V is finite dimensional then, $V \cong V^*$ is not natural since $(-)^*$ flips the variance.

Natural transformations

- For any vector space $V \cong V^{**}$ naturally by evaluation. If V is finite dimensional then, $V \cong V^*$ is not natural since $(-)^*$ flips the variance.
- A functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ is **representable** if there exists an object $c \in \mathbf{C}$ for which $F \cong \text{hom}(c, -)$.

Natural transformations

- For any vector space $V \cong V^{**}$ naturally by evaluation. If V is finite dimensional then, $V \cong V^*$ is not natural since $(-)^*$ flips the variance.
- A functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ is **representable** if there exists an object $c \in \mathbf{C}$ for which $F \cong \text{hom}(c, -)$.
- Hurewicz homomorphism gives that for path connected spaces, $\pi_n(-) \Rightarrow H_n(-)$ and $\text{Ab } \pi_1(-) \cong H_1(-)$.

Natural transformations

- For any vector space $V \cong V^{**}$ naturally by evaluation. If V is finite dimensional then, $V \cong V^*$ is not natural since $(-)^*$ flips the variance.
- A functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ is **representable** if there exists an object $c \in \mathbf{C}$ for which $F \cong \text{hom}(c, -)$.
- Hurewicz homomorphism gives that for path connected spaces, $\pi_n(-) \Rightarrow H_n(-)$ and $\text{Ab } \pi_1(-) \cong H_1(-)$.
- Two categories are said to be **equivalent**, if there exists pair of functors $F: \mathbf{C} \rightleftarrows \mathbf{D} : G$ with natural isomorphisms $GF \cong 1_{\mathbf{C}}$ and $FG \cong 1_{\mathbf{D}}$. If we make the symbols \cong into equalities, then we will talk about isomorphism of categories. Equivalences preserve many of essential categorical properties. The ones not preserved are called **evil**.

math before
category theory



"In a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides."

math after
category theory



- Category of locally compact Hausdorff spaces is dually equivalent to the category of commutative unital C^* -algebras.

Equivalent categories

- Category of locally compact Hausdorff spaces is dually equivalent to the category of commutative unital C^* -algebras.
- The representations of a finite group as a category is isomorphic to representations of its group algebra.

Equivalent categories

- Category of locally compact Hausdorff spaces is dually equivalent to the category of commutative unital C^* -algebras.
- The representations of a finite group as a category is isomorphic to representations of its group algebra.
- Every category is equivalent to its **skeleton** which has objects as one representative of an isomorphism class of the original category.

Equivalent categories

- Category of locally compact Hausdorff spaces is dually equivalent to the category of commutative unital C^* -algebras.
- The representations of a finite group as a category is isomorphic to representations of its group algebra.
- Every category is equivalent to its **skeleton** which has objects as one representative of an isomorphism class of the original category.
- Category of totally disconnected (singletons are the connected components) Hausdorff spaces is oppositely equivalent to the category of Boolean algebras.

Composition


There's an obvious **vertical** composition of natural transformations.

$$\begin{array}{ccccc} F(A) & \xrightarrow{\eta_A} & G(A) & \xrightarrow{\mu_B} & K(A) \\ Ff \downarrow & & Gf \downarrow & & \downarrow Kf \\ F(B) & \xrightarrow{\eta_B} & G(B) & \xrightarrow{\mu_B} & K(B) \end{array}$$

$$\begin{array}{ccc} & F & \\ \curvearrowright & \Downarrow \eta & \curvearrowleft \\ \mathbf{C} & \xrightarrow{\quad} G \xrightarrow{\quad} & \mathbf{D} \\ \curvearrowleft & \Downarrow \mu & \curvearrowright \\ & K & \end{array}$$

Composition 2

There is also something called the **horizontal** composition.



$$\begin{array}{ccccc}
 F'F(C) & \xrightarrow{\mu_{F(C)}} & G'F(C) & & \\
 F'\eta_C \downarrow & & (\mu * \eta)_C \downarrow & & G'\eta_C \downarrow \\
 F'G(C) & \xrightarrow{\mu_{G(C)}} & G'G(C) & &
 \end{array}$$

Definition

A functor is **full** if it is surjective on homsets. Similarly a functor is **faithful** if it is injective on homsets. Combining both definitions we get **fully faithful** functors i.e. bijective on level of homsets. A functor is **essentially surjective** if every object in the image has an object isomorphic to it. An **embedding** is faithful functor that acts "injectively" on objects.

Equivalences in locally small categories

Definition

A functor is **full** if it is surjective on homsets. Similarly a functor is **faithful** if it is injective on homsets. Combining both definitions we get **fully faithful** functors i.e. bijective on level of homsets. A functor is **essentially surjective** if every object in the image has an object isomorphic to it. An **embedding** is faithful functor that acts "injectively" on objects.

Theorem

A functor defining an equivalence is fully faithful and essentially surjective. Converse holds assuming the axiom of choice.

Theorem

(The Yoneda lemma) Given a functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ with \mathbf{C} locally small, there is a bijection

$$\mathrm{Nat}(\mathrm{hom}(c, -), F) \cong Fc$$

that is natural in both F (here F is viewed as an object in the functor category $\mathbf{Set}^{\mathbf{C}}$) and c given by $\eta \mapsto \eta_c(1_c)$.

Theorem

(The Yoneda lemma) Given a functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ with \mathbf{C} locally small, there is a bijection

$$\mathrm{Nat}(\mathrm{hom}(c, -), F) \cong Fc$$

that is natural in both F (here F is viewed as an object in the functor category $\mathbf{Set}^{\mathbf{C}}$) and c given by $\eta \mapsto \eta_c(1_c)$.

Corollary

(Yoneda embeddings) The bifunctor $\mathrm{hom}(-, -): \mathbf{C}^{op} \times \mathbf{C} \rightarrow \mathbf{Set}$. Defines fully faithful embeddings $\mathbf{C} \hookrightarrow \mathbf{Set}^{\mathbf{C}^{op}}$ and $\mathbf{C}^{op} \hookrightarrow \mathbf{Set}^{\mathbf{C}}$.

Consequences of the Yoneda lemma

Definition

The category of elements $\int F$ of a set valued functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ consists of pairs (C, x) with $x \in F(C)$. A morphism $(C, x) \rightarrow (C', x')$ is a morphism $f: C \rightarrow C'$ with $Ff x = x'$.

Consequences of the Yoneda lemma

Definition

The category of elements $\int F$ of a set valued functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ consists of pairs (C, x) with $x \in F(C)$. A morphism $(C, x) \rightarrow (C', x')$ is a morphism $f: C \rightarrow C'$ with $Ff x = x'$.

Theorem

A set valued functor is representable iff the category of elements has an initial object.

Consequences of the Yoneda lemma

Definition

The category of elements $\int F$ of a set valued functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ consists of pairs (C, x) with $x \in F(C)$. A morphism $(C, x) \rightarrow (C', x')$ is a morphism $f: C \rightarrow C'$ with $Ff x = x'$.

Theorem

A set valued functor is representable iff the category of elements has an initial object.

Theorem

If $\text{hom}(C, -) \cong \text{hom}(C', -)$, then $C \cong C'$.

Consequences of the Yoneda lemma

Definition

The category of elements $\int F$ of a set valued functor $F: \mathbf{C} \rightarrow \mathbf{Set}$ consists of pairs (C, x) with $x \in F(C)$. A morphism $(C, x) \rightarrow (C', x')$ is a morphism $f: C \rightarrow C'$ with $Ff x = x'$.

Theorem

A set valued functor is representable iff the category of elements has an initial object.

Theorem

If $\text{hom}(C, -) \cong \text{hom}(C', -)$, then $C \cong C'$.

Proofs of these rely on the proof of Yoneda lemma.

Example of initial object in the category of elements

Let $\text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -): \mathbf{Vect}_{\mathbb{K}} \rightarrow \mathbf{Set}$ the functor of bilinear maps. This functor acts on linear maps by precomposition.

Example of initial object in the category of elements

Let $\text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -): \mathbf{Vect}_{\mathbb{K}} \rightarrow \mathbf{Set}$ the functor of bilinear maps. This functor acts on linear maps by precomposition. Then $\int \text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -)$ consists of pairs (E, L) with E being a \mathbb{K} -vectorspace and $L: V \times W \rightarrow E$ a bilinear map. A morphism $M: (E, L) \rightarrow (E', L')$ is a linear map with

$$\begin{array}{ccc} V \times W & \xrightarrow{L} & E \\ L' \downarrow & \swarrow M & \\ E' & & \end{array}$$

Example of initial object in the category of elements

Let $\text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -): \mathbf{Vect}_{\mathbb{K}} \rightarrow \mathbf{Set}$ the functor of bilinear maps. This functor acts on linear maps by precomposition. Then $\int \text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -)$ consists of pairs (E, L) with E being a \mathbb{K} -vectorspace and $L: V \times W \rightarrow E$ a bilinear map. A morphism $M: (E, L) \rightarrow (E', L')$ is a linear map with

$$\begin{array}{ccc} V \times W & \xrightarrow{L} & E \\ L' \downarrow & \swarrow M & \\ E' & & \end{array}$$

The universal property of the tensor product says that $- \otimes -: V \times W \rightarrow V \otimes_{\mathbb{K}} W$ defines an initial object in the category of elements (hence the name universal property). Therefore $V \otimes_{\mathbb{K}} W$ also represents the functor $\text{Bilin}_{\mathbf{Vect}_{\mathbb{K}}}(V, W; -)$.

Definition


A small **diagram** of shape \mathbf{J} in \mathbf{C} , is a functor $F: \mathbf{J} \rightarrow \mathbf{C}$ with \mathbf{J} being a small category.

Definition

A small **diagram** of shape \mathbf{J} in \mathbf{C} , is a functor $F: \mathbf{J} \rightarrow \mathbf{C}$ with \mathbf{J} being a small category.


Definition

A **cone** over diagram $F: \mathbf{J} \rightarrow \mathbf{C}$ with a **summit** C is a natural transformation $\lambda: c \Rightarrow F$ (thinking of C as a constant natural transformation). So a family of morphisms $(\lambda_j: C \rightarrow Fj)_{j \in \mathbf{J}}$ that satisfies




Definition

Dually there are cones **under** or a **cocone** of a diagram, with a **nadir** C and a natural transformation $\lambda: F \Rightarrow C$ where




Both cones and cocones form a category with the morphism being m such that



Limits

Definition

A limit $\lim_{\mathbf{J}} F$ over a diagram $F: \mathbf{J} \rightarrow \mathbf{C}$ is a universal object in the category of cones. Alternative one could define limits as representing objects of so called cone functors.




We would also like to study (co)continuous functors so ones that preserve (co)limits and also functors that reflect (co)limits.

Examples of limits

- Initial and terminal objects are limits over empty diagrams.


Examples of limits

- Initial and terminal objects are limits over empty diagrams.
- A **product** is limit over diagram with **J** discrete. Examples are cartesian product product topology. Dually it is called a coproduct in **Set** this is for example the disjoint union. In many familiar categories like **Ab**, finite coproducts and products are isomorphic.




Examples of limits

- Initial and terminal objects are limits over empty diagrams.
- A **product** is limit over diagram with **J** discrete. Examples are cartesian product product topology. Dually it is called a coproduct in **Set** this is for example the disjoint union. In many familiar categories like **Ab**, finite coproducts and products are isomorphic.




- An **(co)equalizer** (example: (co)kernel is an equalizer of a homomorphism and the zero map)



Examples of limit

- A direct limit is a colimit indexed by a directed set (such as $1 \rightarrow 2 \rightarrow 3 \rightarrow \dots$). Dually inverse limits are limits over directed sets.
- **Pullback** (dually pushforward) is a limit of a diagram indexed by the category $\bullet \rightarrow \bullet \leftarrow \bullet$.



Existence of limits and continuity

A limit may not exist. For example the category of fields has no initial or terminal object. Neither does the category of manifolds have arbitrary products.

A limit may not exist. For example the category of fields has no initial or terminal object. Neither does the category of manifolds have arbitrary products.

Definition

A category is **(co)complete** if it admits all small (co)limits.

Existence of limits and continuity

A limit may not exist. For example the category of fields has no initial or terminal object. Neither does the category of manifolds have arbitrary products.

Definition

A category is **(co)complete** if it admits all small (co)limits.

Definition

A functor is **(co)continuous** if it preserves (co)limits.

Adjunctions

A fundamental fact is that equivalences preserve limits and colimits. But finding an equivalence is not always possible. Hence we have a weaker notion of equivalence called an adjunction of functors which preserve limits in some sense.

A fundamental fact is that equivalences preserve limits and colimits. But finding an equivalence is not always possible. Hence we have a weaker notion of equivalence called an adjunction of functors which preserve limits in some sense.

Definition

If for pair of functors $F: \mathbf{C} \rightleftarrows \mathbf{D} : G$ there is a natural transformation (in both C and D) $\text{hom}(F(C), D) \cong \text{hom}(C, G(D))$, then F is **left adjoint** to G . Similarly G is **right adjoint** to F . Denote this relation by $F \dashv G$.

A fundamental fact is that equivalences preserve limits and colimits. But finding an equivalence is not always possible. Hence we have a weaker notion of equivalence called an adjunction of functors which preserve limits in some sense.

Definition

If for pair of functors $F: \mathbf{C} \rightleftarrows \mathbf{D} : G$ there is a natural transformation (in both C and D) $\text{hom}(F(C), D) \cong \text{hom}(C, G(D))$, then F is **left adjoint** to G . Similarly G is **right adjoint** to F . Denote this relation by $F \dashv G$.

Theorem

Right adjunctions preserve limits, and left adjunctions preserve colimits.

Examples and non examples of adjunctions

- Tensor-Hom adjunction in **Ab** (or $\mathbb{Z}\text{-Mod}$)
 $\text{hom}(A \otimes B, C) \cong \text{hom}(A, \text{hom}(B, C)).$

Examples and non examples of adjunctions

- Tensor-Hom adjunction in **Ab** (or $\mathbb{Z}\text{-Mod}$)
 $\text{hom}(A \otimes B, C) \cong \text{hom}(A, \text{hom}(B, C)).$
- The Free \dashv Forget adjunction (free vector space, free group, free Abelian group, Free category on a directed multigraph etc... satisfy a certain universal property)

Examples and non examples of adjunctions

- Tensor-Hom adjunction in **Ab** (or $\mathbb{Z}\text{-Mod}$)
 $\text{hom}(A \otimes B, C) \cong \text{hom}(A, \text{hom}(B, C)).$
- The Free \dashv Forget adjunction (free vector space, free group, free Abelian group, Free category on a directed multigraph etc... satisfy a certain universal property)
- There's no right or left adjoint to Forget : **Field** \rightarrow **Set**, since there's no initial or terminal object in the category of fields. Therefore it does not preserve limits, meaning there's no free field.


Category theory in programming

In functional programming one deals with computational side effects by monads. These arise from the units of adjunctions. Moggi's paper "Notions of computation and monads". Functional programming 2 MOOC (which is free) at the University of Helsinki is a good introduction to monads in programming.

Something for the physics students to think about

Definition

A (2d) **TQFT** is a symmetric monoidal functor $Z: \mathbf{2COB} \rightarrow \mathbf{fdVect}_{\mathbb{K}}$. The first category has morphisms between closed 1-manifolds (disjoint union of circles) as cobordisms, composition by combining cobordism and monoidal operation of disjoint union. Think about pair of pants being a cobordism between circle and disjoint union of two circles. The target category is finite dimensional vector spaces over \mathbb{K} with tensor product.



Category theory for a working mathematician by Saunders MacLane, Category theory in context by Emily Riehl, Wikipedia.com, <https://www.dtubbenhauer.com> for definition of TQFT. Memes from facebook, diagrams screenshots from q.uiver, since tikz-cd and beamer don't like each other.