

WILDseq Library Prep v2 Protocol

Callum Malcolm

2023-11-07

Contents

Reverse Transcription	1
Overview	1
Materials	1
RT Protocol	1
PCR Step 1	2
Overview	2
Materials	2
PCR1 Protocol	2
Bead Clean Up	3
Overview	3
Materials	3
Protocol	3

Reverse Transcription

Overview

- cDNA synthesis from RNA template
- Include a -RT control if possible
 - Mix several samples and include extra H2O instead of Reverse Transcriptase
 - PCR this sample as well, checks for contamination of genomic DNA, mastermix, ect.
 - Can also use as negative control to determine product size and yield on tapestation

Materials

- For 40 samples

Component	Expected Volume/experiment	Material ID
5x SSIV Buffer	200 μ L	
SSIV RT	50 μ L	
100 mM DTT	50 μ L	
RNAse Out	50 μ L	
Thermolabile Exonuclease I	1 μ L	
NEBuffer r3.1*	2 μ L	

RT Protocol

1. In PCR strip tubes, prepare 1-5 μ g of RNA in a total volume of 10 μ l of RNAse/DNAase-free water.

2. Add 1 μ l of WS_RT_UMI_NexteraR2 primer (2 μ M stock)
3. Add 1 μ l dNTPs (10 mM each).
4. Denature at 65 C for 5 mins in the PCR machine, then straight onto ice for at least 2 mins, spin briefly to get liquid to the bottom of the tube.
5. Prepare mastermix of RT enzyme and buffers. 1.1x the number of samples.

Component	Volume	MM volume
5x SSIV Buffer	4 μ L	
SSIV RT	1 μ L	
100 mM DTT	1 μ L	
RNase Out	1 μ L	

6. Add 7 μ l of RT MM prepared above to each sample and mix, spin briefly to get liquid to the bottom of the tube.
 - Spin briefly to get liquid to bottom of the tube
7. In the PCR machine incubate at 53 C for 10 mins followed by 80 C for 10 mins.
8. Add 1 μ l Themolabile Exonuclease I (NEB M0568) to remove excess RT primer.

Component	Volume	MM volume
Thermolabile Exonuclease I	1 μ L	
NEBuffer r3.1*	2 μ L	

Most PCR buffers are compatible

9. In PCR Machine: heat at 37C for 4 mins followed by 80C for 1 min
 - Spin briefly to get liquid to bottom of the tube
10. Add 1 μ l of RNase H and incubate at 37 C for 20 mins.
11. Dilute final cDNA 1:2 with DNase/RNase-free water.

PCR Step 1

Overview

- Aim to keep number of cycles at a minimum.
- Forward primer is an equal mix of primers with different length spacers to generate diversity required for sequencing.

Materials

Component	Expected Volume/experiment	Material ID
10uM WS_RT-PCR1_Fwd	70 μ L	-
10uM WS_RT-PCR1_Rev	70 μ L	-
DNase/RNase H20	550 μ L	
Kapa Hifi HotStart Ready Mix (2X)	1200 μ L	KK2601

PCR1 Protocol

- Set up PCR reactions on ice
 - One reaction per sample plus -RT control
1. Prepare mastermix of below reagents at 1.1x the number of samples

Component	Volume
10uM WS_RT-PCR1_Fwd	1.5 μ L
10uM WS_RT-PCR1_Rev	1.5 μ L
DNase/RNase H20	12 μ L
Kapa Hifi HotStart Ready Mix (2X)	25 μ L
Total Master Mix volume	40 μL
cDNA from above	10 μ L

2. Perform PCR using the following parameters:

Step Name	Steps	Time
Initial Denaturation	Step 1: 95C	3mins
Denaturation	Step 2: 98C	20s
Annealing	Step 3: 60C	15s
Extension	Step 4: 72C	15s
Final Extension	Step 5: 72C	1 min
Hold Step	Step 6: 12C	Hold

10-25 cycles of steps 2-4

Clean up reaction using Ampure beads and check product size and yield on tapestation

Bead Clean Up

Overview

- Clean PCR reaction by removing leftover primers
- This prevents false reads, cleaner indexing in later steps, and higher quality samples

Materials

- For 40 samples

Component	Expected Volume/experiment	Material ID
Ampure Bead Kit	200 μ L	

Protocol

1. Equilibrate Ampure beads to room temp (available from supplies cold room)
2. Mix well and vortex for 30sec to ensure uniform distribution.
3. Add 90ul (1.8X volume) to PCR reaction and mix thoroughly by pipetting.
4. Incubate 5min, room temp
5. Place on magnet for 2 min and keep on magnet until final elution
6. Remove supernatant and add 200ul 70% ethanol (make fresh)
7. Incubate 30sec and remove
8. Repeat 70% ethanol wash
9. Air dry beads 5-10min room temp. Once dry the beads will go from shiny to matt in appearance. Avoid over drying (cracked appearance).
10. Add 15 ul EB buffer and resuspend beads by pipetting

11. Incubate 2min room temp
12. Place on magnet and remove eluate
13. Assess PCR product size, contamination and concentration on tapestation. Expected size = 165-172bp