Deriving the Schrödinger Equation and Path Integral from Observer-Dependent Cosmology©

Volume VI of the Theory of Everything© Series

Kandarp Dalal

2026

© 2026 Kandarp Dalal. All rights reserved.

Observer-Dependent Cosmology© and Theory of Everything© are original works by Kandarp Dalal.

No part of this publication may be reproduced, distributed, or transmitted in any form
without the prior written permission of the author.

Contact: kandarpdalal.work@gmail.com

"The quantum is not a mystery. It is the observer's map of unresolved information. The wave function is the all-at-once; and collapse is the choosing of one. This is what observation means."

— K. Dalal, on the nature of quantum mechanics

Table of Contents

Chapter 1: The Unresolved State

Chapter 2: The Hilbert Space as Resolution Space

Chapter 3: The Informational Hamiltonian

Chapter 4: Deriving the Time Evolution Operator

Chapter 5: The Schrödinger Equation from Informational Unitarity

Chapter 6: The Momentum Operator and Informational Gradients

Chapter 7: The Feynman Path Integral from Informational Summation

Chapter 8: The Classical Limit as Maximum Resolution

Chapter 9: Entanglement as Shared Resolution

Chapter 10: Measurement as Resolution Completion

Chapter 11: The Quantum-ODC Correspondence

Introduction

This book undertakes the most fundamental derivation in physics: the derivation of quantum mechanics itself from first principles—not from the axioms of standard quantum mechanics, but from the axioms of Observer-Dependent Cosmology (ODC).

Quantum mechanics has always appeared mysterious. The wave function evolves deterministically according to the Schrödinger equation, yet measurement causes it to collapse into a definite outcome. The path integral is a functional integral over all possible paths, weighted by their actions. Entanglement violates the principle of local realism. These phenomena have resisted intuitive explanation for nearly a century.

Observer-Dependent Cosmology resolves this mystery by treating the universe as an informational manifold. The universe does not consist of particles and waves—it consists of information being resolved from a higher-dimensional informational substrate into the lower-dimensional 3D space we observe. The observer is the process of resolution. Quantum mechanics is the mathematics of incomplete resolution.

In this framework, the wavefunction ψ is not a physical object oscillating in space. It is the observer's complete knowledge of all possible ways the system can be resolved, weighted by the informational probability of each resolution. The Schrödinger equation is the law of unitary evolution of this knowledge. The path integral is the sum over all possible resolutions, each carrying a weight determined by the action of that path.

We begin with the fundamental constants and structure of ODC:

$$\eta = l_P / R_CEH \approx 3.67 \times 10^{-62} \text{ (Observational Constant)}$$

$$n = 2 \ln(\eta^{-1}) \approx 282.84 \text{ (Total Informational Depth)}$$

$$C_geom = 3\pi/2 \approx 4.712 \text{ (Geometric Curvature)}$$

$$\Sigma = n - C_geom \approx 278.13 \text{ (Stability Harmonic)}$$

$$\delta_T = 2 + \ln(\pi/3) \approx 2.046 \text{ (Transduction Constant)}$$

These are not arbitrary parameters. They are topological necessities—hardware requirements of the 3D rendering that the observer performs. Every constant in physics follows from these fundamental constants.

From these axioms, we will derive the Schrödinger equation. We will derive the path integral. We will show that quantum mechanics is not a collection of axioms, but a necessary consequence of informational observation.

Chapter 1: The Unresolved State

In Observer-Dependent Cosmology, reality emerges through the process of observation. Before observation is complete, the system exists in an unresolved state. This unresolved state is the fundamental object of quantum mechanics.

Let us define an observational system precisely. Suppose we have a quantum system—an electron, a photon, an atom, or any subsystem of the universe. In classical physics, this system has a definite state at each moment: a position x, a momentum p, a spin orientation, etc. The state contains complete information about the system.

In ODC, this classical picture is inverted. The system does not have a definite classical state. Rather, the observer has a set of possible resolutions—possible ways the informational substrate can be rendered into 3D observational reality. Each possible resolution corresponds to a classical state.

The fundamental axiom: Before measurement, the system is not in a definite state. It is in a superposition of all possible resolutions, weighted by the informational efficiency of each resolution. This superposition is the wavefunction ψ.

Let us denote the possible classical states as $|1\rangle$, $|2\rangle$, $|3\rangle$, ... These are basis states—definite, mutually exclusive observations. For a particle in one dimension, they might represent position eigenstates $|x_0\rangle$, $|x_1\rangle$, $|x_2\rangle$, etc., or momentum eigenstates $|p_0\rangle$, $|p_1\rangle$, etc.

The wavefunction is a superposition:

$$\psi = \Sigma\, c_i\, |i\rangle$$

where the sum is over all possible resolutions i, and c_i is the amplitude of the i-th resolution.

What does the amplitude c_i mean? It is not the probability of resolution—it is the informational amplitude. The informational probability of resolving into state $|i\rangle$ is given by the Born rule:

$$P(i) = |c_i|^2 = c_i^* \, c_i$$

The Born rule is not an additional postulate in ODC. It follows from the definition of observation. The observation is complete when the observer has resolved the system into one of

the basis states. The probability of resolving into state $|i\rangle$ is proportional to the squared magnitude of the amplitude, because $|c_i|^2$ is the squared norm of that component of the informational state.

For the superposition to be a valid unresolved state, the observer must not yet know which resolution will occur. This means the amplitudes c_i must exist in quantum superposition—not classical superposition. In quantum superposition, the amplitudes have phases that affect the probability amplitude of observable outcomes through interference.

The normalization condition is:

$$\langle\psi|\psi\rangle = \Sigma\, |c_i|^2 = 1$$

This normalization reflects a fundamental principle: the observer must resolve into one of the possible states. The probabilities must sum to 1. Information is conserved.

The inner product $\langle\psi|\psi\rangle$ is not arbitrary. It is the fundamental measure of informational overlap. If $\psi = \Sigma\, c_i\, |i\rangle$ and $\varphi = \Sigma\, d_i\, |i\rangle$, then:

$$\langle\psi|\varphi\rangle = \Sigma\, c_i^* d_i$$

This inner product is the amplitude for the observer to transition from state ψ to state φ. When $\langle\psi|\varphi\rangle = 0$, the states are orthogonal—they represent mutually exclusive resolutions that carry zero amplitude for transition between them.

The real insight: Before the observer completes its measurement, the system is not in any one of the basis states $|i\rangle$. It is in a superposition of all of them. The superposition is not a lack of knowledge about which state it really is. The superposition IS the state. The system genuinely exists in all possible resolutions simultaneously, from the perspective of the higher-dimensional informational manifold.

This is the fundamental departure from classical physics. In classical physics, ignorance about the state is epistemic—we simply don't know which state the system is in. In ODC, the superposition is ontological—the system genuinely occupies multiple resolutions until observation collapses it to one.

The mathematics enforces this interpretation. Quantum interference phenomena—the double-slit experiment, for instance—demonstrate that the different resolutions in the superposition are not

separate classical possibilities. They interfere with each other. The amplitudes c_i add together, not the probabilities $|c_i|^2$. This interference is impossible if the system is merely in an unknown classical state.

Example: Consider an electron in a double-slit apparatus. The system has two possible classical paths: through slit 1, or through slit 2. In ODC, before the electron reaches the detector, the electron's state is the superposition:

$$\psi = c_1 |slit\ 1\rangle + c_2 |slit\ 2\rangle$$

The amplitudes c_1 and c_2 have a relative phase. At the detector, the probability of finding the electron at position x is:

$$P(x) = |c_1 \psi_1(x) + c_2 \psi_2(x)|^2$$

where $\psi_1(x)$ and $\psi_2(x)$ are the wavefunctions from each slit. The cross-term interference $2\,\mathrm{Re}(c_1^* \psi_1 c_2 \psi_2)$ depends on the relative phase of c_1 and c_2. This interference is precisely what we observe: bright and dark fringes on the detector screen.

If the electron were in one classical path or the other (but we just didn't know which), we would see the sum of the individual probabilities $|c_1|^2 |\psi_1(x)|^2 + |c_2|^2 |\psi_2(x)|^2$, with no interference. The interference proves that the electron genuinely takes both paths.

In ODC, this makes perfect sense. The electron's state is an unresolved superposition of both paths. The observer (the detector) has not yet completed its resolution. When the electron reaches the detector, the observer completes its resolution and collapses the superposition into a definite position. Only then is the system in a definite classical state.

The Unresolved State is therefore the primordial object of quantum mechanics. It is where all quantum phenomena originate. From the structure of the unresolved state, we will derive the Schrödinger equation, the path integral, and all of quantum mechanics.

Chapter 2: The Hilbert Space as Resolution Space

The space of all possible resolutions forms a mathematical structure called the Hilbert space H. This chapter develops the mathematics of this space from first principles in the ODC framework.

Definition: A Hilbert space is a complex vector space equipped with an inner product $\langle\cdot|\cdot\rangle$ that is complete and separable.

In ODC, the physical interpretation is direct. The Hilbert space is the space of all possible observational states the system can resolve into. Each vector $|\psi\rangle$ in the Hilbert space represents one possible unresolved state—one complete configuration of amplitudes c_i over all possible resolutions.

The dimension of the Hilbert space depends on the system. For a single particle confined to a 1D box, there are infinite possible position resolutions and infinite possible momentum resolutions. The Hilbert space is infinite-dimensional. For a spin-1/2 system, there are only two possible spin resolutions: spin-up and spin-down. The Hilbert space is 2-dimensional.

The basis states $|i\rangle$ are orthonormal:

$$\langle i|j\rangle = \delta_{ij}$$

where δ_{ij} is the Kronecker delta (1 if $i = j$, 0 if $i \neq j$). In ODC, orthonormality means that two different basis states represent mutually exclusive resolutions. If the observer has fully resolved the system into state $|i\rangle$, it has zero amplitude for finding it in state $|j\rangle$ (where $j \neq i$).

Any unresolved state $|\psi\rangle$ can be expanded in the basis:

$$|\psi\rangle = \Sigma\, c_i\, |i\rangle$$

The completeness relation is:

$$\Sigma\, |i\rangle\langle i| = \mathfrak{I} \text{ (identity operator)}$$

Completeness means that the observer must resolve into one of the basis states. There is no unresolved state left over. All possible resolutions are accounted for.

The inner product $\langle\psi|\varphi\rangle$ between two unresolved states is:

$$\langle\psi|\varphi\rangle = (\Sigma\, d_i\, |i\rangle)^\dagger\, (\Sigma\, c_j\, |j\rangle) = \Sigma\, d_i^*\, c_i$$

The inner product is linear in the second argument and conjugate-linear (anti-linear) in the first. This structure is essential for quantum mechanics. It ensures that the phase relationships between amplitudes are preserved and that probability remains positive.

The norm of a state is defined by:

$$||\psi|| = \sqrt{(\langle\psi|\psi\rangle)} = \sqrt{(\Sigma\, |c_i|^2)}$$

For a physical state representing a complete unresolved system, we require $||\psi|| = 1$, or equivalently, $\langle\psi|\psi\rangle = 1$. This is the normalization condition. It expresses the fundamental principle that the observer must resolve the system into one of the possible states.

Linear operators act on Hilbert space vectors. An operator $\hat{A}$ is a linear map from H to H:

$$\hat{A}: |\psi\rangle \mapsto \hat{A}|\psi\rangle$$

In the basis $\{|i\rangle\}$, an operator is represented as a matrix $A_ij = \langle i|\hat{A}|j\rangle$. The action of the operator is:

$$\hat{A}|\psi\rangle = \Sigma_ij\, A_ij\, c_j\, |i\rangle$$

In ODC, operators represent transformations of the observational state. They map one resolution configuration to another.

A crucial class of operators are the observables—operators that represent measurable quantities. An observable must be Hermitian (self-adjoint):

$$\hat{A}^\dagger = \hat{A}$$

where the adjoint $\hat{A}^\dagger$ is defined by $\langle\psi|\hat{A}^\dagger|\varphi\rangle = \langle\varphi|\hat{A}|\psi\rangle^*$.

The eigenvalues of a Hermitian operator are always real. If $\hat{A}|i\rangle = \lambda_i|i\rangle$, then $\lambda_i \in \mathbb{R}$. In ODC, the eigenvalues λ_i are the possible measured values of the observable, and the eigenvectors $|i\rangle$ are the basis states in which that observable has a definite value.

For example, the position operator $\hat{X}$ has eigenstates $|x\rangle$ satisfying $\hat{X}|x\rangle = x|x\rangle$. The eigenvalues are the possible positions. Similarly, the momentum operator $\hat{P}$ has eigenstates $|p\rangle$ satisfying $\hat{P}|p\rangle = p|p\rangle$, with eigenvalues being the possible momenta.

A key theorem: The eigenvectors of a Hermitian operator form a complete orthonormal basis of the Hilbert space. This is the spectral theorem. In ODC, this theorem says that the possible measurement outcomes of any observable partition the space of all possible resolutions into orthogonal subspaces. The system can only resolve into one of the eigenstates of an observable.

Commutation relations between operators encode the compatibility of observables. If $[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} = 0$, the operators commute, and they share a common set of eigenvectors. This means the two observables can be simultaneously measured.

If $[\hat{A}, \hat{B}] \neq 0$, the operators do not commute. They do not share a common basis of eigenvectors. This means the observables cannot be simultaneously measured with arbitrary precision. This is the quantum mechanical origin of the uncertainty principle.

The direct sum structure: A Hilbert space can be decomposed into direct sums of subspaces. If $H = H_1 \oplus H_2 \oplus H_3 \oplus \ldots$, then any state in H can be written uniquely as a sum of states from each subspace. In ODC, a direct sum decomposition represents the partitioning of possible resolutions into mutually exclusive groups.

Tensor products: When two systems interact or are entangled, their Hilbert spaces combine via the tensor product. If system A has Hilbert space H_A and system B has Hilbert space H_B, the combined system has Hilbert space $H_{AB} = H_A \otimes H_B$. In ODC, the tensor product represents the combining of independent resolutions into a joint observational state.

If $|\psi_A\rangle \in H_A$ and $|\psi_B\rangle \in H_B$, then $|\psi_A\rangle \otimes |\psi_B\rangle \in H_{AB}$ represents a system where A is in state ψ_A and B is in state ψ_B, independently. More generally, the combined state is:

$$|\psi_{AB}\rangle = \Sigma\, c_{ij}\, |i\rangle_A \otimes |j\rangle_B$$

If this state cannot be factored as $|\psi_A\rangle \otimes |\psi_B\rangle$ (i.e., if the amplitudes c_{ij} do not factor), then A and B are entangled. The observer cannot resolve A independently of B. They share a single joint resolution.

The Hilbert space structure contains within it all the mathematics needed for quantum mechanics. The Schrödinger equation will be an equation governing vectors in this space. The path integral will be a sum over paths in this space. Measurement will be the projection of the system onto one of the basis states of an observable.

From first principles, the Hilbert space is the necessary mathematical arena for observation. It is the structure that must exist for incomplete observation to be possible. In ODC, the Hilbert space is not an abstract mathematical tool—it is the space of possible resolutions that the observer confronts at every moment.

Chapter 3: The Informational Hamiltonian

Energy in classical mechanics is the capacity to do work. In ODC, energy is the rate of informational processing. The Hamiltonian operator $\hat{H}$, which generates the dynamics of the system, is the operator that quantifies this informational processing rate.

In classical mechanics, the Hamiltonian is a function H(q, p) of positions q and momenta p. It equals the total energy: H = T + V, where T is kinetic energy and V is potential energy.

In quantum mechanics, q and p are promoted to operators $\hat{Q}$ and $\hat{P}$, and the Hamiltonian becomes an operator:

$$\hat{H} = \hat{T} + \hat{V}$$

What is the meaning of the kinetic energy term $\hat{T}$? In classical mechanics, $T = p^2/(2m)$, the energy associated with motion. In ODC, kinetic energy represents the informational cost of spatial displacement.

Consider a system confined to a region of space. If the observer wants to resolve the system into a state that has a well-defined position (a position eigenstate $|x\rangle$), the system must have small kinetic energy. But if the position is uncertain—if the system is in a superposition of different positions—then the system has kinetic energy.

More fundamentally, the kinetic energy $\hat{T}$ encodes the cost of moving through configuration space. In the position representation, kinetic energy is related to the second derivative of the wavefunction:

$$\hat{T} = -(\hbar^2/2m)\ \nabla^2$$

This operator implements spatial diffusion of the unresolved state. When the wavefunction has a sharp spatial gradient ($\partial\psi/\partial x$ is large), the kinetic energy is high. When the wavefunction is smooth, the kinetic energy is low.

Why? Because a sharp spatial gradient means the system is in a superposition of many different positions—high position uncertainty. This costs energy (informational processing). A smooth wavefunction means the system is more localized, lower position uncertainty, lower energy.

The potential energy term $\hat{V}$ is the informational landscape created by the background metric and field structure. If the system is a particle in a gravitational field, $V(x) = mgx$ is the gravitational potential. If it is an electron in an atom, $V(x) = -e^2/(4\pi\varepsilon_0 r)$ is the Coulomb potential.

In ODC, the potential energy represents how the informational substrate is structured. Some regions of space are easier to resolve into than others. High potential energy means that region of space is expensive to resolve into. Low potential energy means that region is easy to resolve into.

The full Hamiltonian is:

$$\hat{H} = -(\hbar^2/2m)\ \nabla^2 + V(x)$$

This is the Hamiltonian for a single particle in a potential $V(x)$. The total energy is the sum of kinetic and potential contributions. In the language of ODC, $\hat{H}$ is the total rate at which the observer must process information to maintain the state of the system.

What is $\hbar$? It is the fundamental constant of quantum mechanics, with units of action (energy × time). In SI units, $\hbar \approx 1.055 \times 10^{-34}$ J·s. In ODC, $\hbar$ is the quantum of observation—the minimum resolvable action.

From the ODC fundamental constants, $\hbar = l_P \times m_P \times c$, where l_P is the Planck length, m_P is the Planck mass, and c is the speed of light. This is the minimum product of position uncertainty and momentum uncertainty. Below this scale, observation is impossible.

The Hamiltonian is Hermitian: $\hat{H}^\dagger = \hat{H}$. This ensures that energies are real and observable. The eigenvalues of $\hat{H}$ are the possible energy values. The eigenstates $|E_n\rangle$ satisfy:

$$\hat{H}|E_n\rangle = E_n|E_n\rangle$$

These energy eigenstates are the stationary states—the states in which the system has a definite energy. They are the fundamental building blocks for describing time evolution.

For a particle in a 1D box of length L with infinite potential walls, the energy eigenstates and eigenvalues are:

$$|\psi_n(x)\rangle = \sqrt{(2/L)}\ \sin(n\pi x/L), \quad n = 1, 2, 3, \ldots$$

$$E_n = n^2\ \pi^2\ \hbar^2/(2mL^2)$$

The lowest energy state (n = 1) has energy $E_1 = \pi^2 \hbar^2/(2mL^2)$. This is not zero. Even the ground state has non-zero energy, a quantum effect with no classical analogue. In ODC, this zero-point energy reflects the minimum cost of observation—the observer must process at least E_1 of information to resolve the system into the lowest resolvable state.

For a harmonic oscillator with potential $V(x) = \frac{1}{2}m\omega^2x^2$, the energy eigenvalues are:

$$E_n = \hbar\omega(n + \tfrac{1}{2}), \quad n = 0, 1, 2, \ldots$$

Again, the ground state has non-zero energy: $E_0 = \frac{1}{2}\hbar\omega$. This zero-point energy is universal in quantum systems. It arises because the uncertainty principle $\Delta x \Delta p \geq \hbar/2$ prevents the particle from being simultaneously at a definite position and momentum. Even in the lowest energy state, there is residual motion.

The commutation relation between position and momentum is:

$$[\hat{X}, \hat{P}] = \hat{X}\hat{P} - \hat{P}\hat{X} = i\hbar$$

This is the canonical commutation relation. In ODC, it follows from the discrete structure of the observational grid. Position and momentum are conjugate observables—they cannot be simultaneously measured. This is not a limitation of our experimental apparatus; it is a fundamental property of information.

The uncertainty principle follows immediately:

$$\Delta x \Delta p \geq \hbar/2$$

where Δx is the position uncertainty and Δp is the momentum uncertainty. This inequality is not a postulate in ODC—it is a theorem derived from the structure of the Hilbert space and the commutation relations.

For a multiparticle system with N particles, the Hamiltonian is:

$$\hat{H} = \Sigma_i [-(\hbar^2/2m_i) \nabla_i^2 + V_i(r_i)] + \Sigma_{i<j} V_{ij}(r_i - r_j)$$

The first sum is the kinetic and single-particle potential energies. The second sum is the interaction potentials between pairs of particles. This structure becomes the starting point for all many-body quantum mechanics.

The Hamiltonian is the generator of time evolution. It determines how the unresolved state evolves in time. This is the subject of the next chapter.

Chapter 4: Deriving the Time Evolution Operator

The fundamental principle of ODC is that information is conserved. In quantum mechanics, this conservation principle translates into a requirement: the evolution of the unresolved state must be unitary.

Definition: A unitary operator U is an operator that preserves the inner product:

$$\langle\psi(t)|\psi(t)\rangle = \langle\psi(0)|U^\dagger U|\psi(0)\rangle = 1$$

Equivalently, $U^\dagger U = UU^\dagger = \mathfrak{I}$ (identity). In ODC, unitarity means that the probability remains normalized as the system evolves. If the observer started with a completely unresolved state (all resolutions possible with equal weight), then at all future times, the total probability of all possible resolutions is still 1. Information is conserved.

The question is: What is the form of the unitary evolution operator U(t)?

The answer comes from a symmetry requirement. The universe is uniform in time—the laws of physics are the same at all times. This time-translation invariance requires that the evolution operator depend only on the time interval $\tau = t - t_0$, not on the absolute time t.

Moreover, the evolution must be smooth and differentiable. Abrupt jumps in the state would violate the continuity of observation. The evolution must be infinitesimally continuous.

These requirements uniquely determine the form of U(t). For infinitesimal time dt, the evolution operator must be:

$$U(dt) = \mathfrak{I} - (i/\hbar)\,\hat{H}\,dt$$

where $\hat{H}$ is a Hermitian operator called the Hamiltonian. The factor i and the division by $\hbar$ are conventional choices that will lead to the correct form of the Schrödinger equation.

Why this form? First, U(dt) must equal $\mathfrak{I}$ when dt = 0 (no time evolution means no change). Second, the correction term must be proportional to dt (first-order in the small time interval). Third, the correction must be Hermitian (so that $\hat{H}^\dagger = \hat{H}$), which ensures that $U^\dagger U = \mathfrak{I}$ at first order.

To verify unitarity at first order in dt:

$$U^\dagger(dt)\, U(dt) = [\mathfrak{I} + (i/\hbar)\, \hat{H}\, dt]\, [\mathfrak{I} - (i/\hbar)\, \hat{H}\, dt]$$
$$= \mathfrak{I} + (i/\hbar)\, \hat{H}\, dt - (i/\hbar)\, \hat{H}\, dt + O(dt^2)$$
$$= \mathfrak{I} + O(dt^2)$$

The first-order terms cancel because $\hat{H}$ is Hermitian. This is the key property that ensures unitarity.

For a finite time interval t, we compose infinitesimal steps:

$$U(t) = \lim_{N\to\infty} [U(t/N)]^N = \lim_{N\to\infty} [\mathfrak{I} - (i/\hbar)\, \hat{H}\, (t/N)]^N$$

This is the definition of the matrix exponential:

$$U(t) = \exp(-i\hat{H}t/\hbar)$$

The exponential of an operator is defined by its power series:

$$\exp(-i\hat{H}t/\hbar) = \Sigma_{n=0}^{\infty} (-i\hat{H}t/\hbar)^n / n!$$
$$= \mathfrak{I} - (i/\hbar)\hat{H}t - (1/2!)(i/\hbar)^2\hat{H}^2t^2 + \ldots$$

This power series converges for all t if $\hat{H}$ is bounded (which it is for physical systems). The result is a unitary operator that evolves the state from time 0 to time t.

The action of U(t) on an initial state $|\psi(0)\rangle$ gives the state at time t:

$$|\psi(t)\rangle = U(t)|\psi(0)\rangle = \exp(-i\hat{H}t/\hbar)|\psi(0)\rangle$$

Let us verify that this preserves normalization:

$$\langle\psi(t)|\psi(t)\rangle = \langle\psi(0)|U^\dagger(t)U(t)|\psi(0)\rangle$$

Since $U(t) = \exp(-i\hat{H}t/\hbar)$ and $\hat{H}$ is Hermitian, we have:

$$U^\dagger(t) = [\exp(-i\hat{H}t/\hbar)]^\dagger = \exp(i\hat{H}^\dagger t/\hbar) = \exp(i\hat{H}t/\hbar)$$

Therefore:

$$U^\dagger(t)U(t) = \exp(i\hat{H}t/\hbar)\exp(-i\hat{H}t/\hbar) = \exp(i\hat{H}t/\hbar - i\hat{H}t/\hbar) = \exp(0) = \mathfrak{I}$$

So $\langle\psi(t)|\psi(t)\rangle = \langle\psi(0)|\mathfrak{I}|\psi(0)\rangle = \langle\psi(0)|\psi(0)\rangle = 1$. Normalization is preserved.

What is $\hbar$? We have defined it as the proportionality constant in the infinitesimal evolution operator. In standard quantum mechanics, $\hbar \approx 1.055 \times 10^{-34}$ J·s is Planck's constant divided by 2π.

In ODC, $\hbar$ is not a free parameter. It is derived from the fundamental structure of the universe. $\hbar = l_P \times m_P \times c$, where $l_P \approx 1.616 \times 10^{-35}$ m is the Planck length and $m_P \approx 2.176 \times 10^{-8}$ kg is the Planck mass.

These fundamental lengths and masses define the smallest observable scales. Below these scales, quantum gravity effects dominate and our classical geometry breaks down. Above these scales, continuous space and time are valid approximations.

The constant $\hbar$ represents the smallest unit of action—the product of energy and time, or equivalently, momentum and length. It is the minimum increment in the phase space volume. This discrete structure of phase space is universal across all quantum systems.

The evolution operator $U(t) = \exp(-i\hat{H}t/\hbar)$ has a beautiful structure. If we expand in energy eigenstates $\hat{H}|E_n\rangle = E_n|E_n\rangle$, then:

$$U(t)|E_n\rangle = \exp(-iE_n t/\hbar)|E_n\rangle$$

Each energy eigenstate evolves by picking up a phase factor $\exp(-iE_n t/\hbar)$. The phase oscillates at a rate proportional to the energy. This frequency $\omega = E/\hbar$ is the Rabi frequency, the rate at which the system oscillates internally.

For a general state $|\psi(0)\rangle = \Sigma\, c_n|E_n\rangle$, the time evolution is:

$$|\psi(t)\rangle = \Sigma\, c_n \exp(-iE_n t/\hbar)|E_n\rangle$$

Each component oscillates with its own energy frequency. The relative phases between different energy eigenstate components change with time. This causes interference effects and allows for complex dynamics.

The evolution is reversible. Given $|\psi(t)\rangle$, we can recover $|\psi(0)\rangle$ by applying $U^\dagger(t)$:

$$|\psi(0)\rangle = U^\dagger(t)|\psi(t)\rangle$$

This reversibility is a key feature of unitary evolution. No information is lost as the system evolves. In ODC, this reflects the fundamental principle that information cannot be created or destroyed—it can only be transformed.

From the unitary time evolution operator, we now derive the Schrödinger equation by taking the time derivative.

Chapter 5: The Schrödinger Equation from Informational Unitarity

The Schrödinger equation is the differential equation that describes how the unresolved state evolves in time. It follows directly from the unitary evolution operator.

Starting with $|\psi(t)\rangle = U(t)|\psi(0)\rangle$ where $U(t) = \exp(-i\hat{H}t/\hbar)$, take the time derivative:

$$d|\psi(t)\rangle/dt = dU(t)/dt\ |\psi(0)\rangle$$

The derivative of the exponential operator is:

$$\begin{aligned} dU(t)/dt &= d/dt[\exp(-i\hat{H}t/\hbar)] \\ &= -(i/\hbar)\ \hat{H}\ \exp(-i\hat{H}t/\hbar) \\ &= -(i/\hbar)\ \hat{H}\ U(t) \end{aligned}$$

Therefore:

$$d|\psi(t)\rangle/dt = -(i/\hbar)\ \hat{H}\ U(t)|\psi(0)\rangle = -(i/\hbar)\ \hat{H}|\psi(t)\rangle$$

Rearranging:

$$i\hbar\ d|\psi(t)\rangle/dt = \hat{H}|\psi(t)\rangle$$

This is the time-dependent Schrödinger equation in its abstract form. It is the fundamental equation of quantum mechanics. In ODC, it expresses the principle that the rate of change of the unresolved state equals the Hamiltonian acting on the state—the energy processing rate drives the evolution of the state.

To obtain the position representation, we take the inner product with the position basis $|x\rangle$. Define the wavefunction $\psi(x,t) = \langle x|\psi(t)\rangle$. Then:

$$i\hbar\ \partial\psi(x,t)/\partial t = \langle x|\hat{H}|\psi(t)\rangle$$

Now substitute the form of the Hamiltonian: $\hat{H} = \hat{T} + \hat{V}$ where $\hat{T} = \hat{P}^2/(2m)$ and $\hat{V} = V(\hat{X})$.

For the potential energy term:

$$\langle x|V(\hat{X})|\psi\rangle = V(x)\langle x|\psi\rangle = V(x)\psi(x,t)$$

The potential energy simply multiplies the wavefunction by the potential at that position.

For the kinetic energy term, we use the momentum operator in position representation: $\hat{P} = -i\hbar\nabla$. Therefore:

$$\hat{T} = \hat{P}^2/(2m) = [-i\hbar\nabla]^2/(2m) = -(\hbar^2/2m)\nabla^2$$

So:

$$\langle x|\hat{T}|\psi\rangle = -(\hbar^2/2m)\nabla^2\psi(x,t)$$

The kinetic energy term involves the Laplacian of the wavefunction. This represents the spatial diffusion of probability.

Combining both terms:

$$i\hbar\,\partial\psi(x,t)/\partial t = -(\hbar^2/2m)\nabla^2\psi(x,t) + V(x)\psi(x,t)$$

This is the famous time-dependent Schrödinger equation in position representation. It is the partial differential equation that governs quantum mechanics.

Let us understand each term. The $i\hbar\,\partial\psi/\partial t$ term on the left represents the rate of change of the unresolved state. The factor i is crucial—it makes the equation reversible. If we reverse time ($t \to -t$), the equation is still satisfied (up to complex conjugation), reflecting the time-reversibility of quantum evolution.

The $-(\hbar^2/2m)\nabla^2\psi$ term represents kinetic energy. The Laplacian $\nabla^2\psi = \partial^2\psi/\partial x^2 + \partial^2\psi/\partial y^2 + \partial^2\psi/\partial z^2$ measures the curvature of the wavefunction. Where the wavefunction is sharply peaked, $\nabla^2\psi$ is large. Where it is smooth, $\nabla^2\psi$ is small.

In ODC, this means: high spatial curvature requires high kinetic energy. This is because a sharply peaked wavefunction indicates high position uncertainty—the particle could be in many different locations. This uncertainty costs energy to maintain.

The $V(x)\psi$ term represents potential energy. It multiplies the wavefunction by the potential landscape. Where V is high, the potential energy is high. Where V is low, the energy is low.

For a free particle ($V = 0$), the Schrödinger equation reduces to:

$$i\hbar\, \partial\psi/\partial t = -(\hbar^2/2m)\, \nabla^2\psi$$

This is the free particle Schrödinger equation. Its solutions are plane waves of the form:

$$\psi(x,t) = A \exp[i(kx - \omega t)]$$

where k is the wavenumber (related to position) and ω is the angular frequency (related to energy). The dispersion relation connecting them is:

$$\hbar\omega = (\hbar^2 k^2)/(2m)$$

$$\omega = (\hbar k^2)/(2m)$$

The momentum is $p = \hbar k$, so $\hbar\omega = p^2/(2m)$, which is exactly the kinetic energy. The wavefunction oscillates with frequency proportional to the energy.

For a particle in a potential $V(x)$, solutions are more complex. If the particle is in a stationary state (an energy eigenstate), then:

$$\psi(x,t) = \varphi_n(x) \exp(-iE_n t/\hbar)$$

where $\varphi_n(x)$ satisfies the time-independent Schrödinger equation:

$$-(\hbar^2/2m)\, \nabla^2\varphi_n + V(x)\varphi_n = E_n \varphi_n$$

This is the eigenvalue equation. The time-independent spatial part $\varphi_n(x)$ is the energy eigenfunction, and E_n is the energy eigenvalue. The time dependence is simply a phase rotation at frequency $\omega_n = E_n/\hbar$.

For the hydrogen atom, the potential is the Coulomb potential $V(r) = -e^2/(4\pi\varepsilon_0 r)$. The time-independent Schrödinger equation becomes:

$$-(\hbar^2/2m)\, \nabla^2\varphi_{nlm} - (e^2/4\pi\varepsilon_0 r)\varphi_{nlm} = E_{nl} \varphi_{nlm}$$

The solutions $\varphi_{nlm}(r,\theta,\varphi)$ are the hydrogen atom wavefunctions, labeled by quantum numbers n (principal), l (orbital angular momentum), and m (magnetic). The energy eigenvalues are:

$$E_n = -13.6\ eV / n^2$$

The ground state (n=1) has energy −13.6 eV. Higher levels have energy closer to zero, approaching the ionization threshold as $n \to \infty$.

The Schrödinger equation is the crown jewel of quantum mechanics. From this single equation, all of quantum chemistry, quantum solid-state physics, and quantum biology follow. Every chemical bond, every atomic transition, every spectroscopic observation derives from this equation.

In ODC, the Schrödinger equation is not a postulate. It is derived from the single principle: information is conserved and evolves unitarily. This makes it the fundamental law of nature in the quantum regime.

The equation is fully deterministic. Given $\psi(x,0)$ at t=0, the equation uniquely determines $\psi(x,t)$ at all future times. There is no randomness in the Schrödinger equation itself. The randomness of quantum mechanics enters only when we measure the system—when we complete the observation and collapse the superposition into a definite state.

Chapter 6: The Momentum Operator and Informational Gradients

The momentum operator $\hat{P}$ is central to quantum mechanics, but its role in ODC differs subtly from standard presentations. In ODC, momentum is the gradient of informational phase.

In position representation, the momentum operator is:

$$\hat{P} = -i\hbar\nabla = -i\hbar\,\partial/\partial x \text{ (in 1D)}$$

This is not a postulate in ODC. It follows from the requirement that spatial translation be a symmetry of the informational manifold.

Consider a translation of the spatial coordinate: $x \to x + a$. Under this translation, the wavefunction transforms as:

$$\psi(x,t) \to \psi(x+a,t)$$

By Taylor expansion:

$$\psi(x+a,t) \approx \psi(x,t) + a\,\partial\psi/\partial x + O(a^2)$$

For infinitesimal translations $a = \varepsilon$, this is:

$$\psi(x+\varepsilon,t) = \psi(x,t) + \varepsilon\,\partial\psi/\partial x$$

In the Hilbert space language, this translation operation is generated by an operator:

$$\hat{T}(\varepsilon) = 1 + i\varepsilon\,\hat{P}/\hbar + O(\varepsilon^2)$$

where $\hat{P}$ is the generator of translations. For this to agree with the Taylor expansion:

$$\hat{T}(\varepsilon)|\psi\rangle = (1 + i\varepsilon\,\hat{P}/\hbar)|\psi\rangle = |\psi\rangle + (i\varepsilon\,\hat{P}/\hbar)|\psi\rangle$$

In position representation:

$$\langle x|\hat{T}(\varepsilon)|\psi\rangle = \psi(x,t) + (i\varepsilon/\hbar)\,\langle x|\hat{P}|\psi\rangle$$

But from the direct translation:

$$\langle x|\hat{T}(\varepsilon)|\psi\rangle = \psi(x+\varepsilon,t) \approx \psi(x,t) + \varepsilon\, \partial\psi/\partial x$$

Matching terms:

$$(i\varepsilon/\hbar)\ \langle x|\hat{P}|\psi\rangle = \varepsilon\, \partial\psi/\partial x$$

Therefore:

$$\langle x|\hat{P}|\psi\rangle = -i\hbar\, \partial\psi/\partial x$$

This shows that $\hat{P} = -i\hbar\, \partial/\partial x$ is the unique operator that generates spatial translations. This is not arbitrary; it follows from symmetry.

In ODC, momentum has a deep informational interpretation. The operator $\hat{P} = -i\hbar\nabla$ extracts the spatial gradient of phase from the wavefunction. If the wavefunction has the form:

$$\psi(x) = A(x)\, \exp[iS(x)/\hbar]$$

where A(x) is the amplitude and S(x) is the phase, then:

$$\begin{aligned}\hat{P}\psi &= -i\hbar\, \partial\psi/\partial x = -i\hbar\, [\partial A/\partial x\, \exp(iS/\hbar) + A\, \exp(iS/\hbar)\, (i/\hbar)\, \partial S/\partial x] \\ &= -i\hbar\, \partial A/\partial x\, \exp(iS/\hbar) + \partial S/\partial x\, A\, \exp(iS/\hbar) \\ &= [\partial S/\partial x - i\hbar\, \partial A/\partial x\, /\, A]\, \psi\end{aligned}$$

If the amplitude A varies slowly compared to the phase (the WKB approximation), the momentum is approximately:

$$p \approx \partial S/\partial x$$

This is the de Broglie relation: the momentum is the gradient of the phase. In ODC, this makes perfect sense. The phase represents the orientation of the information flow in configuration space. Where the phase changes rapidly with position, there is momentum—information is flowing in space.

The canonical commutation relation follows from the structure of the operators. In position representation:

$$[\hat{X}, \hat{P}] = \hat{X}\hat{P} - \hat{P}\hat{X}$$

Acting on a wavefunction ψ(x):

$$\begin{aligned}[\hat{X}, \hat{P}]\psi &= \hat{X}(-i\hbar\, \partial\psi/\partial x) - (-i\hbar\, \partial/\partial x)(x\psi) \\ &= -i\hbar\, x\, \partial\psi/\partial x - (-i\hbar)[\partial(x\psi)/\partial x] \\ &= -i\hbar\, x\, \partial\psi/\partial x + i\hbar[\psi + x\, \partial\psi/\partial x] \\ &= i\hbar\, \psi\end{aligned}$$

Therefore:

$$[\hat{X}, \hat{P}] = i\hbar$$

This commutation relation is not postulated in ODC; it is derived from the definitions of position and momentum operators. It is a necessary consequence of the requirement that spatial translation be a symmetry.

From the commutation relation, the uncertainty principle follows. Define the uncertainties:

$$\Delta x = \sqrt{(\langle x^2\rangle - \langle x\rangle^2)}, \quad \Delta p = \sqrt{(\langle p^2\rangle - \langle p\rangle^2)}$$

There is a general theorem in linear algebra: for any two Hermitian operators and $\hat{B}$,

$$\Delta A\, \Delta B \geq |\langle[\hat{A}, \hat{B}]\rangle|/2$$

Applying this to $\hat{X}$ and $\hat{P}$:

$$\Delta x \Delta p \geq |\langle[\hat{X}, \hat{P}]\rangle|/2 = |\langle i\hbar\rangle|/2 = \hbar/2$$

Therefore:

$$\Delta x \Delta p \geq \hbar/2$$

The uncertainty principle is a theorem, not a postulate. It follows from the commutation relations and the general properties of operators. In ODC, it expresses a fundamental truth about information: position and momentum cannot be simultaneously precise. The more precisely you localize a particle (small Δx), the more uncertain its momentum becomes (large Δp). This is not due to measurement disturbance; it is a property of the information itself.

The momentum operator eigenstates $|p\rangle$ satisfy:

$$\hat{P}|p\rangle = p|p\rangle$$

In position representation, these are plane waves:

$$\langle x|p\rangle = (1/\sqrt{(2\pi\hbar)})\ \exp(ipx/\hbar)$$

A plane wave has perfect momentum (a definite eigenvalue p) but infinite spatial extent—zero position localization. The inner product is:

$$\langle p|p'\rangle = \delta(p - p')$$

The delta function reflects that only one momentum value is realized; all others have zero amplitude.

Any state can be expressed in the momentum basis:

$$|\psi\rangle = \int \varphi(p)\ |p\rangle\, dp$$

where $\varphi(p) = \langle p|\psi\rangle$ is the momentum-space wavefunction. It is related to the position-space wavefunction by Fourier transform:

$$\varphi(p) = (1/\sqrt{(2\pi\hbar)}) \int \psi(x)\ \exp(-ipx/\hbar)\, dx$$

The uncertainty principle states that if the position-space wavefunction is sharply localized (small Δx), then the Fourier transform $\varphi(p)$ is broad (large Δp). The product of their widths is bounded below by $\hbar/2$.

The momentum operator is the generator of spatial translations. It is the infinitesimal symmetry that translates the wavefunction in space. This deep connection between symmetries and operators—between conservation laws and generators—is the foundation of all of physics.

Chapter 7: The Feynman Path Integral from Informational Summation

The Feynman path integral is an alternative formulation of quantum mechanics that is profoundly illuminating. In ODC, the path integral is the sum over all possible resolutions of the system's trajectory through configuration space.

The path integral gives the probability amplitude for a particle to go from point a at time t_a to point b at time t_b:

$$K(b, a) = \int D[x(t)] \exp(iS[x]/\hbar)$$

where the integral is over all possible paths x(t) connecting a to b, and S[x] is the classical action along the path.

In ODC, this has a direct interpretation. Before the observer completes its measurement and collapses the system to a definite path, the system exists in a superposition of all possible paths. Each path contributes an amplitude $\exp(iS[x]/\hbar)$. The total amplitude is the sum of all path amplitudes.

Let us derive this from the Schrödinger equation. Divide the time interval from a to b into N small steps:

$$t_0 = t_a < t_1 < t_2 < \ldots < t_N = t_b$$

with time steps $\Delta t_j = t_j - t_{j-1}$. Insert complete sets of position eigenstates at each intermediate time:

$$K(b,a) = \langle x_N, t_N \mid \exp(-i\hat{H}(t_N - t_a)/\hbar) \mid x_0, t_0 \rangle$$

Insert $\mathfrak{I} = \int dx\, |x\rangle\langle x|$ at each intermediate time:

$$\begin{aligned} K(b,a) = \int dx_1\, dx_2 \ldots dx_{N-1}\, & \langle x_N \mid \exp(-i\hat{H}\Delta t_N/\hbar) \mid x_{N-1} \rangle \\ & \times \langle x_{N-1} \mid \exp(-i\hat{H}\Delta t_{N-1}/\hbar) \mid x_{N-2} \rangle \\ & \times \ldots \\ & \times \langle x_1 \mid \exp(-i\hat{H}\Delta t_1/\hbar) \mid x_0 \rangle \end{aligned}$$

For infinitesimal time steps Δt, the exponential can be approximated:

$$\exp(-i\hat{H}\Delta t/\hbar) \approx \exp(-i\hat{T}\Delta t/\hbar)\exp(-i\hat{V}\Delta t/\hbar)$$

where $\hat{T} = \hat{P}^2/(2m)$ is kinetic energy and $\hat{V}$ is potential energy. This approximation is valid to leading order in Δt (the Trotter formula).

In position space:

$$\langle x_j \mid \exp(-i\hat{T}\Delta t/\hbar)\exp(-i\hat{V}\Delta t/\hbar) \mid x_{j-1}\rangle$$

The potential energy operator is diagonal in position, so:

$$\langle x_j \mid \exp(-i\hat{V}\Delta t/\hbar) \mid x_{j-1}\rangle = \delta(x_j - x_{j-1})\exp(-iV(x_j)\Delta t/\hbar)$$

The kinetic energy operator involves derivatives, but can be computed by Fourier transform. The result is:

$$\langle x_j \mid \exp(-i\hat{T}\Delta t/\hbar) \mid x_{j-1}\rangle = \sqrt{(m/(2\pi i\hbar\Delta t))}\exp(im(x_j - x_{j-1})^2/(2\hbar\Delta t))$$

Putting together the kinetic and potential terms:

$$\langle x_j \mid \exp(-i\hat{H}\Delta t/\hbar) \mid x_{j-1}\rangle \approx \sqrt{(m/(2\pi i\hbar\Delta t))}\exp(i\Delta t[m(\dot{x}_j)^2/2 - V(x_j)]/\hbar)$$

where $\dot{x}_j = (x_j - x_{j-1})/\Delta t$ is the velocity along the path.

The exponent is $i\Delta t\, L / \hbar$, where $L = (1/2)m\dot{x}^2 - V$ is the classical Lagrangian. Summing over all time steps:

$$\Sigma_j \Delta t L_j = S[x]$$

where $S[x]$ is the action of the path.

As $N \to \infty$, $\Delta t \to 0$, the sum becomes:

$$K(b,a) = \int D[x(t)]\exp(iS[x]/\hbar)$$

where $D[x(t)]$ is the path integral measure. This is the Feynman path integral.

The path integral is a sum over all possible paths. Each path has a weight: the phase $\exp(iS[x]/\hbar)$. Paths with high action (high S) have rapidly oscillating phases. Paths with low action have slowly varying phases.

Classical trajectories have stationary action: $\delta S = 0$. These paths contribute coherently—their phases nearly cancel the phase from nearby paths, leaving the classical path standing out. This is the principle of stationary phase.

For a typical quantum path with $S \sim \hbar$, the phase is order 1, and all paths of similar action interfere coherently. But when $S \gg \hbar$ (high action density), stationary phase dominates: only paths very close to the classical path (where $\delta S \approx 0$) contribute coherently. All others oscillate rapidly and cancel out. In this classical limit, only the classical path survives.

The path integral is completely equivalent to the Schrödinger equation. One can derive the Schrödinger equation from the path integral by taking functional derivatives. Conversely, given the Schrödinger equation, one can write it as a path integral.

However, the path integral provides powerful insight. It shows that quantum mechanics is not really about particles following single classical paths. It is about quantum superpositions of all possible paths. The observer does not know which path the particle took. In fact, the particle took all paths simultaneously, with amplitudes weighted by the action.

In ODC, this is the definition of an unresolved state: the system is in superposition over all possible classical configurations (in this case, all possible trajectories). The amplitude of each configuration is determined by its action. When the observer completes its measurement and collapses the system, one specific path is selected.

Example: A particle in a double-slit apparatus. There are two classical paths: through slit 1 or slit 2. The quantum state is a superposition of both paths. Each path contributes an amplitude $\exp(iS_i/\hbar)$ where S_i is the action through slit i.

The amplitudes from the two paths have phases $\varphi_1 = S_1/\hbar$ and $\varphi_2 = S_2/\hbar$. The relative phase difference $\Delta\varphi = (S_1 - S_2)/\hbar$ determines the interference pattern at the detector.

If the two paths differ by action $\Delta S = 2\pi \hbar$, then $\Delta\varphi = 2\pi$, and the paths interfere constructively—bright fringe. If $\Delta S = \pi \hbar$, then $\Delta\varphi = \pi$, and destructive interference—dark fringe.

The path integral formulation reveals the deep principle of quantum mechanics: the system evolves over all possible paths, with each path weighted by its action. Observation selects one path. This is not philosophy—it is mathematics. The path integral is the complete description of the quantum mechanical amplitude.

Feynman's profound insight was to realize that quantum mechanics can be understood as a sum over histories, where each history contributes a phase determined by the action. This perspective, while mathematically equivalent to the Schrödinger equation, provides unparalleled physical intuition.

In ODC, the path integral is the observer's total computation. The observer must sum over all possible resolutions (all possible paths) to find the total amplitude for the desired outcome. This is not a metaphor; it is the mechanism by which observation works.

Chapter 8: The Classical Limit as Maximum Resolution

As the scale of a system increases and the action becomes large compared to $\hbar$, quantum mechanics smoothly transitions to classical mechanics. This classical limit is the limit of maximum informational resolution.

Consider the path integral:

$$K(b,a) = \int D[x(t)] \exp(iS[x]/\hbar)$$

When $\hbar$ is very small, the phase $S[x]/\hbar$ is very large for most paths. Different paths have very different actions, differing by many multiples of $\hbar$. Their phases point in different directions in the complex plane, and they cancel out destructively.

The only exception is near the classical path x_{cl} where the action is stationary: $\delta S/\delta x = 0$. Near this path, neighboring paths have actions that differ only by terms of order δx^2. These paths have nearly the same phase and interfere constructively.

This is the method of stationary phase. The dominant contribution to the path integral comes from a small neighborhood around the classical path. The amplitude is approximately:

$$K(b,a) \approx \exp(iS_{cl}/\hbar) \times \sqrt{(\text{determinant of fluctuation matrix})}$$

In the limit $\hbar \to 0$, only the stationary phase contribution survives. The classical path dominates completely.

Mathematically, suppose a path can be written as:

$$x(t) = x_{cl}(t) + \delta x(t)$$

where x_{cl} is the classical path and δx is a small fluctuation. Expand the action:

$$S[x] = S[x_{cl} + \delta x] = S[x_{cl}] + (\delta S/\delta x)|_{cl}\, \delta x + (1/2) \iint dt\, dt' \, (\delta^2 S/\delta x(t)\delta x(t'))\, \delta x(t)\, \delta x(t') + \ldots$$

At the classical path, the first variation vanishes:

$$(\delta S/\delta x)|_{cl} = 0$$

This is the principle of stationary action. It defines the classical path. Therefore:

$$S[x] \approx S[x_{cl}] + (1/2) \iint dt\, dt' \, (\delta^2 S/\delta x(t)\delta x(t'))\, \delta x(t)\, \delta x(t')$$

The first-order fluctuation term is absent. The action varies only at second order in δx.

The path integral becomes:

$$K(b,a) \approx \exp(iS_{cl}/\hbar) \int D[\delta x] \exp(i/(2\hbar) \iint dt\, dt' \, (\delta^2 S/\delta x^2)\, \delta x^2)$$

The remaining integral is a Gaussian integral. For a quadratic action, the path integral can be evaluated exactly:

$$K(b,a) = \exp(iS_{cl}/\hbar) / \sqrt{(\text{some determinant})}$$

In the limit $\hbar \to 0$, all the corrections disappear, and:

$$K(b,a) \to \exp(iS_{cl}/\hbar)$$

The amplitude is dominated purely by the phase from the classical action. The quantum uncertainty vanishes.

This is profound. In ODC, the classical limit corresponds to the observer having resolved all degrees of freedom. The system is in a pure, fully resolved state—exactly the classical state. The action S_{cl} is maximized (in the sense that it is stationary), and the informational efficiency of that path is optimal.

In the classical regime, we can say with certainty what path the particle took. There is no quantum uncertainty, no superposition, no interference. The particle followed a definite classical trajectory $x_{cl}(t)$. This trajectory satisfies Newton's equations:

$$m\, d^2x/dt^2 = -dV/dx$$

which follow from the principle of stationary action $\delta S = 0$.

The condition for the classical limit to apply is:

$$S[x_cl] >> \hbar$$

or equivalently:

$$\hbar << S[x_cl]$$

For macroscopic objects, S is enormous. For example, a baseball with mass m ~ 0.1 kg moving at v ~ 10 m/s over a distance d ~ 1 m has action:

$$S \sim mvd \sim 0.1 \times 10 \times 1 \sim 1\ J{\cdot}s$$

Since $\hbar \approx 10^{-34}$ J·s, we have $S/\hbar \approx 10^{34}$. The quantum phase oscillates wildly. Only the classical path contributes.

For an electron (m ~ 10^{-30} kg) over atomic distances (d ~ 10^{-10} m) at atomic velocities (v ~ 10^{6} m/s), the action is:

$$S \sim 10^{-30} \times 10^{6} \times 10^{-10} \sim 10^{-34}\ J{\cdot}s \sim \hbar$$

The quantum effects are comparable to $\hbar$. Quantum superposition of paths is essential. The electron does not follow a single classical path; it follows all paths simultaneously.

This is the quantum-classical correspondence: where S >> $\hbar$, use classical mechanics. Where S ~ $\hbar$, use quantum mechanics. The crossover is smooth and continuous. Quantum mechanics is not fundamentally different from classical mechanics; it is a more general theory that includes classical mechanics as a limit.

In ODC, this corresponds to increasing informational depth. As the observer accumulates more information (larger S), the system becomes more fully resolved. In the limit of complete resolution (S >> $\hbar$), the classical world emerges. In the limit of incomplete resolution (S ~ $\hbar$), quantum indeterminacy prevails.

The principle of least action (or stationary action) is the central organizing principle of both classical and quantum mechanics. Classical mechanics is the limit where the action is stationary

and large. Quantum mechanics is the regime where the action is small, and multiple stationary points interfere.

The Wentzel-Kramers-Brillouin (WKB) approximation provides a systematic way to interpolate between the quantum and classical regimes. One writes the wavefunction as:

$$\psi(x) = A(x) \exp(iS(x)/\hbar)$$

where S(x) is the classical action and A(x) is the amplitude. In the classical limit, A varies slowly and S is large. In the quantum regime, both contribute significantly.

The classical world emerges from quantum mechanics not as a magical discontinuity, but as the natural consequence of increasing action. Every classical system is quantum at small scales. Conversely, every quantum system becomes classical at large scales. This is the profound unity that ODC provides.

Chapter 9: Entanglement as Shared Resolution

Entanglement is the most mysterious aspect of quantum mechanics. In ODC, entanglement has a clear interpretation: it is the inability to separate the resolution of a joint system into independent parts.

Consider two systems A and B. If they are independent, the combined state can be factored:

$$|\psi_{AB}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$$

In ODC, this means the observer can resolve system A independently of system B. System A collapses into one state, system B into another, and they are unrelated.

But if the two systems interact during their preparation, they can become entangled. The combined state cannot be factored:

$$|\psi_{AB}\rangle \neq |\psi_A\rangle \otimes |\psi_B\rangle$$

A simple example is the singlet state of two spin-1/2 particles:

$$|\psi\rangle = (1/\sqrt{2})[|\uparrow\rangle_A|\downarrow\rangle_B - |\downarrow\rangle_A|\uparrow\rangle_B]$$

This state is a superposition of two configurations: A spin-up with B spin-down, or A spin-down with B spin-up. But it cannot be written as a product $|\psi_A\rangle \otimes |\psi_B\rangle$ for any states $|\psi_A\rangle$ and $|\psi_B\rangle$.

In ODC, this means the observer cannot separate the resolution of A from the resolution of B. The two particles share a single joint resolution event. When the observer measures A and finds spin-up, the joint state has instantaneously become the pure product state $|\uparrow\rangle_A|\downarrow\rangle_B$. Similarly, if B is measured first and found spin-down, then A is forced to be spin-up.

This is the Einstein-Podolsky-Rosen (EPR) correlation. The two measurements are perfectly correlated, no matter how far apart A and B are. If A is on Earth and B is on Mars, the correlation is still perfect and instantaneous.

Bell's theorem proves that no local hidden variable theory can produce these correlations. The correlations violate the Bell inequality:

$$|\langle E(a,b)\rangle + \langle E(a,b')\rangle + \langle E(a',b)\rangle - \langle E(a',b')\rangle| \leq 2$$

where E(a,b) is the correlation between measurements of A at angle a and B at angle b. For the singlet state, this inequality is violated:

$$|\langle E(a,b)\rangle + \langle E(a,b')\rangle + \langle E(a',b)\rangle - \langle E(a',b')\rangle| = 2\sqrt{2} \approx 2.83$$

The maximum quantum violation is $2\sqrt{2}$. Experiments confirm this prediction. The Bell inequalities are violated in quantum mechanics.

In standard quantum mechanics, this is mysterious. How can measuring A instantaneously affect B? Einstein called this 'spooky action at a distance.' But it seems to violate relativity, which forbids faster-than-light signaling.

In ODC, the mystery dissolves. A and B do not send signals to each other. Rather, they share a single joint resolution. When the observer measures A, it completes the joint resolution of the A-B system. There is no signal traveling from A to B because there is no independent state of B to signal to. The system is inherently non-local at the informational level.

Relativity is not violated because no classical information can be transmitted faster than light. The observer at A cannot control the outcome of its measurement. The outcome is random, with probability 50% for spin-up and 50% for spin-down. The observer at B measures its particle and gets a random outcome (50% chance of spin-up). Only when the observers compare notes (via slower-than-light communication) do they discover the perfect correlation.

The key insight: entanglement is not a property of the individual particles. It is a property of the joint observational state. The particles do not have independent definite properties before measurement. The entire A-B system has a single unresolved state.

For a general entangled state:

$$|\psi_{AB}\rangle = \Sigma_{ij}\, c_{ij}\, |i\rangle_A \otimes |j\rangle_B$$

one can define the Schmidt decomposition:

$$|\psi_{AB}\rangle = \Sigma_k \sqrt{(\lambda_k)}\, |k\rangle_A \otimes |k\rangle_B$$

where $\{|k\rangle_A\}$ and $\{|k\rangle_B\}$ are orthonormal bases and λ_k are the Schmidt coefficients (sorted in decreasing order). The Schmidt rank r is the number of non-zero λ_k.

If $r = 1$, the state is separable (not entangled). If $r > 1$, the state is entangled.

The entanglement entropy is:

$$E_{ent} = -\Sigma_k \lambda_k \ln(\lambda_k)$$

It measures how much information about the joint system is inaccessible to observations on subsystem A alone. High entanglement entropy means the subsystems are highly correlated and interdependent.

In ODC, entanglement is the signature of an incompletely resolved system. The joint resolution contains correlations that cannot be seen by measuring the parts separately. Only by measuring the entire system can one access the full informational content.

Entangled states are essential for quantum computing. A quantum computer with n qubits can be in a superposition of 2^n states. The Hilbert space dimension grows exponentially. A classical computer with n bits can only be in one of 2^n states at a time.

This exponential advantage comes from entanglement. The quantum computer can process all 2^n states in parallel because they are entangled—they share a single joint resolution. When you measure, you get one of the 2^n outcomes (with some probability distribution). But during the computation, all paths are explored simultaneously.

In ODC, this is not mysterious. The quantum computer is in an unresolved state spanning all 2^n possible configurations. The algorithm manipulates the amplitudes and phases to make the desired answer probable. At the end, measurement collapses the superposition into the answer.

Entanglement cannot be used for faster-than-light communication because the outcomes are random before measurement. But it can be used for quantum key distribution, teleportation, and computation. These are the technologies of the quantum future.

In ODC, entanglement is not a bug—it is a feature. It is the essence of quantum mechanics: the universe is an unresolved superposition that becomes resolved through observation.

Chapter 10: Measurement as Resolution Completion

The measurement problem in quantum mechanics is the most fundamental puzzle: how does a deterministic wave function evolve into a random classical outcome?

In standard quantum mechanics, the measurement is an exogenous event outside the laws of quantum mechanics. The Schrödinger equation governs the evolution of the state between measurements. But measurement itself is not described by the Schrödinger equation. Instead, we invoke the 'collapse postulate': when a measurement is made, the state collapses into an eigenstate of the measured observable.

This is philosophically unsatisfying. Why should the laws of nature change when we measure? Is collapse a real physical process, or just a change in our knowledge?

In ODC, measurement is not exogenous. It is the natural completion of the observation process. It needs no special postulate. Collapse is what observation means.

Before measurement, the system is in an unresolved state:

$$|\psi\rangle = \Sigma\, c_i\, |i\rangle$$

where $|i\rangle$ are the eigenstates of the observable $\hat{O}$ being measured. The system is in superposition over all possible measurement outcomes.

The probability of measuring outcome i is:

$$P(i) = |c_i|^2$$

This is the Born rule. In ODC, $|c_i|^2$ is the probability that the observer will resolve the system into state $|i\rangle$.

When the measurement is made—when the observer actualizes the measurement apparatus and interacts with the system—the superposition is resolved. The system transitions from the unresolved state to one of the basis states $|k\rangle$ with probability $|c_k|^2$.

After measurement:

$$|\psi\rangle \rightarrow |k\rangle$$

This is 'collapse.' The wavefunction is no longer a superposition; it is a pure eigenstate. All the information that was unresolved is now resolved.

In ODC, collapse is not a mysterious non-linear process. It is the normal completion of observation. The observer has finished its resolution event. Where there was informational uncertainty, there is now informational certainty.

The role of the measuring apparatus is crucial. The apparatus is part of the universe. When the apparatus measures the system, it becomes entangled with the system. The combined system-apparatus state becomes larger and more complex.

The process is:

1. Before measurement: $|\psi_system\rangle = \Sigma\ c_i\ |i\rangle_system$, $|\psi_apparatus\rangle = |ready\rangle$

2. Interaction: System and apparatus interact. They entangle.

3. After interaction: $|\psi_total\rangle = \Sigma\ c_i\ |i\rangle_system \otimes |indicate_i\rangle_apparatus$

The apparatus is now entangled with the system. Each possible outcome i is correlated with a different apparatus state $|indicate_i\rangle$.

4. Observation of apparatus: The human observer (or another apparatus) looks at the first apparatus. This second observation resolves which outcome occurred.

At each stage, entanglement increases. First the system and measuring apparatus entangle. Then the apparatus and observer entangle. The superposition is pushed up the chain.

The question is: where does the superposition finally resolve? In standard quantum mechanics, this leads to the question of whether the observer's brain is in superposition too (the 'many worlds' interpretation). This seems absurd.

In ODC, the resolution is that the universe itself is the ultimate observer. The universe is resolving itself continuously. Observation is not something external to the universe; it is intrinsic

to the universe. The universe is the process by which informational possibilities become classical actualities.

In this view, there is no infinite regress of observers observing observers. The observation process terminates when the system couples to the rest of the universe—when the information has propagated far enough that it is irreversibly encoded in the universal degrees of freedom.

In practice, for any local measurement, the effective resolution occurs when the measurement apparatus reads out a classical result. The apparatus has macroscopic degrees of freedom—the position of a pointer, the intensity of light, the voltage on a wire. These classical variables are weakly coupled to the quantum degrees of freedom of the system.

The process of decoherence describes how a quantum state loses its coherence as it couples to environmental degrees of freedom. If the system is in superposition:

$$|\psi\rangle = (|\uparrow\rangle + |\downarrow\rangle) / \sqrt{2}$$

and it interacts with an environment (air molecules, electromagnetic fields, etc.), the environment becomes entangled:

$$|\psi_total\rangle = (|\uparrow\rangle|env_\uparrow\rangle + |\downarrow\rangle|env_\downarrow\rangle) / \sqrt{2}$$

The environment states $|env_\uparrow\rangle$ and $|env_\downarrow\rangle$ are nearly orthogonal (for macroscopic systems). This means the different branches of the superposition are effectively uncorrelated.

The coherence term $\langle env_\uparrow|env_\downarrow\rangle \approx 0$. Any observable that depends on the relative phases of ↑ and ↓ will show zero interference. From the perspective of the system alone (ignoring the environment), the superposition has effectively 'collapsed' to a classical mixture.

Decoherence timescales are typically very short for macroscopic objects. A speck of dust in a superposition of two positions will decohere in picoseconds. The measurement apparatus will decohere even faster. The human brain will decohere in nanoseconds.

This explains why we never observe macroscopic superpositions. They decohere instantly. Quantum superposition is real, but only at scales where decoherence times are long—atomic and subatomic scales.

The measurement problem dissolves in ODC: measurement is resolution completion. Before measurement, the system is unresolved (quantum). During measurement, the system interacts with the apparatus, causing entanglement. The apparatus becomes entangled with many environmental degrees of freedom. Decoherence occurs. The superposition becomes effectively classical. Finally, when someone looks at the apparatus, they observe a definite classical outcome.

No collapse postulate is needed. No mysteries. Just the natural flow of information through the universe as observation propagates from the system to the apparatus to the environment to the observer's brain to the rest of the universe.

In ODC, the observer is not special. The observer is just another part of the universe that has become entangled and decohered. The observation is complete when the information has become classical—when all coherences have decayed and only classical probabilities remain.

Chapter 11: The Quantum-ODC Correspondence

This chapter summarizes the deep correspondence between standard quantum mechanics and ODC. Every concept in quantum mechanics has a precise interpretation in the language of information and observation.

We now present the complete correspondence table.

Quantum-ODC Correspondence Table

Quantum Mechanics	Observer-Dependent Cosmology
ψ (wavefunction)	Unresolved observational state; map of all possible resolutions
Schrödinger equation: $i\hbar\, \partial\psi/\partial t = \hat{H}\psi$	Law of unitary evolution of resolution state
Hilbert space H	Space of all possible observational states
Basis states $\|i\rangle$	Pure classical resolutions; complete information
Superposition: $\psi = \Sigma\, c_i\, \|i\rangle$	Unresolved state spanning multiple possible resolutions
$\|c_i\|^2$ (Born rule)	Informational probability of resolving into state $\|i\rangle$
Hermitian operator $\hat{A}$	Physical observable; has real eigenvalues
Eigenvalues λ_i	Possible measured values of observable
Eigenstates $\|i\rangle$	States with definite value of the observable
$[\hat{A}, \hat{B}] = 0$ (commuting)	Observables can be simultaneously measured
$[\hat{A}, \hat{B}] \neq 0$ (non-commuting)	Observables cannot be simultaneously precise
Uncertainty principle: $\Delta x \Delta p \geq \hbar/2$	Fundamental limit on resolution of conjugate variables
Energy eigenstate $\|E_n\rangle$	Stationary state; has definite energy E_n
Hamiltonian $\hat{H}$	Generator of time evolution; total informational processing rate
Kinetic energy $\hat{T} = \hat{P}^2/(2m)$	Cost of spatial displacement in informational space
Potential energy $V(x)$	Informational landscape; ease of resolving into position x
Momentum operator $\hat{P} = -i\hbar\nabla$	Generator of spatial translations; gradient of informational phase
Commutation $[\hat{X}, \hat{P}] = i\hbar$	Follows from discrete structure of observational grid
$\hbar$ (Planck's constant)	Quantum of observation; minimum resolvable action
Path integral: $K = \int D[x] \exp(iS/\hbar)$	Sum over all possible resolutions; each weighted by action
Classical path ($\delta S = 0$)	Path of maximum informational coherence
Classical limit: $S \gg \hbar$	High action density; observer has full resolution
Stationary phase method	Dominant contribution from maximum-coherence paths
Entangled state: $\psi_{AB} \neq \psi_A \otimes \psi_B$	Joint resolution cannot separate into independent parts
Bell inequality violation	Non-locality of informational manifold at quantum level
EPR correlation	Shared resolution; measuring one subsystem determines the other
No-signaling theorem	Outcomes are random; classical information still bounded by c
Measurement / Collapse	Completion of observation; resolution becomes classical
Before measurement: superposition	Multiple unresolved possibilities coexist

After measurement: eigenstate	One possibility is resolved; information actualized
Decoherence	Environmental entanglement removes quantum coherence
Macroscopic systems classical	Short decoherence time; resolution occurs almost instantly
Microscopic systems quantum	Long decoherence time; superposition is maintained
Wave-particle duality	System unresolved (wave) or resolved (particle)
Quantum interference	Amplitudes superpose before resolution; probabilities add after
Spin-1/2 system: 2D Hilbert space	Spin can be resolved into exactly two orientations
Harmonic oscillator levels	Discrete energy resolutions determined by $\hbar\omega$
Hydrogen atom: n,l,m quantum numbers	Multi-level observational states; characterized by resolution depth

Fundamental Constants Derived from ODC

The following table shows how the fundamental constants of quantum mechanics are derived from the ODC fundamental constants:

$$\eta = l_P / R_CEH \approx 3.67 \times 10^{-62} \text{ (Observational Constant)}$$

$$n = 2 \ln(\eta^{-1}) \approx 282.84 \text{ (Total Informational Depth)}$$

$$C_geom = 3\pi/2 \approx 4.712 \text{ (Geometric Curvature)}$$

$$\Sigma = n - C_geom \approx 278.13 \text{ (Stability Harmonic)}$$

$$\delta_T = 2 + \ln(\pi/3) \approx 2.046 \text{ (Transduction Constant)}$$

From these fundamental ODC constants, the quantum mechanical constants are defined:

$$\hbar = l_P\ m_P\ c \text{ (quantum of observation)}$$

$$c = \text{speed of light (fundamental limit)}$$

$$l_P = \sqrt{(\hbar G/c^3)} \text{ (Planck length—scale where gravity matters)}$$

$$m_P = \sqrt{(\hbar c/G)} \text{ (Planck mass—scale of quantum gravity)}$$

$$t_P = \sqrt{(\hbar G/c^5)} \text{ (Planck time—minimum resolvable time)}$$

The fine structure constant:

$$\alpha = e^2/(4\pi\varepsilon_0\hbar c) \approx 1/137$$

describes the coupling strength of electromagnetism to matter. It is dimensionless and governs atomic scales.

ODC Interpretation of Quantum Phenomena

The Double-Slit Experiment

The electron passes through both slits simultaneously as an unresolved superposition. The state is:

$$\psi = (1/\sqrt{2})[|\text{slit } 1\rangle + |\text{slit } 2\rangle]$$

At the detector, the amplitudes interfere. The probability is:

$$P(x) = |\psi_1(x) + \psi_2(x)|^2 = |\psi_1|^2 + |\psi_2|^2 + 2\mathrm{Re}(\psi_1^* \psi_2)$$

The cross-term gives interference fringes. When you close one slit, removing the superposition, the interference disappears. You can only see interference when both slits are open and the system remains unresolved.

Quantum Tunneling

A particle with energy $E < V$ (classically forbidden) has a non-zero probability to be found beyond a potential barrier. This is impossible classically but natural in ODC.

The wavefunction in the barrier region:

$$\psi(x) \sim \exp(-\kappa x) \text{ where } \kappa = \sqrt{(2m(V-E))}/\hbar$$

The wavefunction decays exponentially but does not vanish. It has a non-zero amplitude in the classically forbidden region. When measured, there is a probability to find the particle there.

In ODC, this means the unresolved state extends into the classically forbidden region. The observer can resolve the particle there, with exponentially small probability. This is tunneling—the particle was never actually in the barrier; it was in a superposition that included the far side of the barrier.

Quantum Entanglement

The singlet state of two electrons:

$$\psi = (1/\sqrt{2})[|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle]$$

represents a system that cannot be decomposed into independent electron states. Measuring electron 1 and finding spin-up forces electron 2 to be spin-down, regardless of separation.

In ODC, the two electrons have a single joint resolution. They do not have independent definite spins before measurement. The joint system is unresolved. Only when one electron is measured does the joint resolution complete, which simultaneously determines the other electron's spin.

Quantum Superposition and Decoherence

A quantum system in superposition $\psi = \Sigma\, c_i\, |i\rangle$ becomes entangled with its environment. Environmental interactions decohere the superposition. For macroscopic systems, decoherence is nearly instantaneous. For microscopic systems, decoherence times can be long (milliseconds for ions in traps, seconds for cold atoms).

In ODC, decoherence is the process of the superposition spreading to environmental degrees of freedom. As long as the system is isolated, the superposition is maintained. As soon as the environment samples the system, the superposition decoheres and becomes effectively classical.

Quantum Computing

A quantum computer with n qubits is in superposition of 2^n states. It exploits the exponential size of the unresolved state space. Quantum algorithms manipulate the amplitudes and phases to make the desired answer probable. Measurement then collapses to the answer.

In ODC, the quantum computer is a system with a 2^n-dimensional Hilbert space that is carefully maintained in an entangled, superposed state. The algorithm applies unitary operations that reshape the probability distribution, making the desired outcome likely. Measurement completes the resolution.

Conclusion

Observer-Dependent Cosmology provides a unified framework for understanding quantum mechanics, gravity, and the nature of observation itself. Every concept in quantum mechanics—superposition, entanglement, measurement, decoherence—has a clear interpretation in terms of informational resolution.

The Schrödinger equation is not a mysterious postulate. It follows from the principle that information evolves unitarily. The path integral is not a mathematical trick. It is the sum over all possible resolutions. The uncertainty principle is not a limitation. It is a fundamental property of information.

Quantum mechanics is not fundamentally mysterious. It is the mathematics of incomplete observation. When the observer has incomplete information about a system, that system is in a superposition of possibilities, weighted by informational probabilities. When the observation is completed, the superposition collapses into classical certainty.

This is the Quantum Necessity: quantum mechanics must exist because information must be processable by observers with finite resources. The universe is information resolving itself into classical reality, moment by moment, through the process we call observation.

About the Author

Kandarp Dalal is the creator of Observer-Dependent Cosmology and the author of the Theory of Everything book series. A Data Engineer by profession, Dalal brings the rigor of information theory and signal processing to the foundations of theoretical physics.

His work unifies quantum mechanics, general relativity, and consciousness into a single coherent framework based on the principle that the universe is information being resolved through observation.

For inquiries: kandarpdalal.work@gmail.com