Deriving Maxwell's Equations from Observer-Dependent Cosmology©

Volume VIII of the Theory of Everything© Series

Kandarp Dalal

2026

© 2026 Kandarp Dalal. All rights reserved.

Observer-Dependent Cosmology© and Theory of Everything© are original works by Kandarp Dalal.

No part of this publication may be reproduced, distributed, or transmitted in any form
without the prior written permission of the author.

Contact: kandarpdalal.work@gmail.com

"The universe is not made of matter and energy—it is made of information,
and electromagnetism is how that information is written onto the observer's resolution grid."
— K. Dalal

Chapter 1: The Phase Freedom of Observation

The foundation of observer-dependent cosmology rests on a fundamental observation: the observer's resolution function ψ encodes all information that the universe can communicate about itself. This function is not uniquely defined—there exists a continuous freedom in how one can represent the same physical reality.

Consider the resolution function $\psi(x)$ at a point x in spacetime. The Born interpretation tells us that $|\psi(x)|^2$ is the probability amplitude, or equivalently in ODC, the information density at that location. Now, let us perform a transformation on this function:

$$\psi(x) \rightarrow e^{i\theta}\psi(x)$$

where θ is a constant phase angle independent of position. This is a global U(1) phase rotation. Under this transformation, the magnitude squared remains invariant:

$$|e^{i\theta}\psi(x)|^2 = |\psi(x)|^2$$

This invariance is absolute. The information content of the resolution function—everything observable—depends only on $|\psi|^2$, not on the phase. Therefore, any observable quantity, any measurement, any deduction about reality must be unchanged by a global phase rotation.

What does this phase represent in ODC? The phase θ encodes the temporal coordinate at which resolution occurs. When we observe the universe, we observe it at a specific 'when'—a temporal slice through the information manifold. The global phase θ is the timestamp of resolution.

Here is the crucial insight: the fact that observation is invariant under global phase rotations means that the temporal coordinate is unobservable in isolation. We cannot ask 'what is the absolute time?' We can only ask 'what is the relative time difference between observations?' This is the deeper meaning of the time-translation symmetry of physics.

In the language of gauge theory, this global phase invariance is called a U(1) symmetry. The symmetry group U(1) is the group of unitary transformations on a single complex number—precisely the group of phase rotations. The existence of this symmetry is not an assumption; it is forced upon us by the structure of information.

The completeness of this phase freedom is reflected in the term 'global.' The same phase θ applies at all points in spacetime. This uniformity is possible because the transformation preserves causality—no signal needs to be sent to synchronize the phase throughout the universe.

The invariance under global phase rotations is the most fundamental symmetry of the quantum universe. It is exact—there are no corrections, no violations, no exceptions. This exactness will prove to be the origin of electromagnetism itself.

Chapter 2: Promoting Global to Local—The Gauge Principle

The global phase invariance discussed in Chapter 1 is only the beginning. We must now ask a more demanding question: can the observer apply a different phase rotation at different points in spacetime? Can θ depend on position?

$$\psi(x) \rightarrow e^{i\theta(x)}\psi(x)$$

At first, this might seem like a mere mathematical generalization. But it carries profound physical consequences. In ODC, allowing the phase to vary locally means allowing the timestamp of resolution to vary from point to point. This is a statement about causality: the 'when' of observation can be different at different 'where' locations.

Now consider the implications for derivatives. The partial derivative ∂_μ represents a rate of change with respect to spacetime coordinates. When we take $\partial_\mu\psi$, we are asking: how does ψ change as we move in the μ-direction?

$$\partial_\mu\psi(x) = \partial_\mu|\psi(x)|^2 + \text{phase terms}$$

Under a global phase rotation, this derivative transforms covariantly because the exponential factor is constant:

$$\partial_\mu[e^{i\theta}\psi(x)] = e^{i\theta}\partial_\mu\psi(x)$$

But under a local phase rotation, the situation changes drastically:

$$\partial_\mu[e^{i\theta(x)}\psi(x)] = e^{i\theta(x)}[\partial_\mu\psi(x) + i(\partial_\mu\theta(x))\psi(x)]$$

The derivative of the phase function $\partial_\mu\theta(x)$ appears as a new term. This additional term breaks the covariance. The derivative $\partial_\mu\psi$ no longer transforms simply under local phase rotations.

This is the crisis that the gauge principle resolves. If we insist on local phase invariance as a fundamental symmetry—and we must, because locality is a fundamental principle of the information manifold—then we cannot use the naive partial derivative. We must modify it.

Introduce a new object: $A_\mu(x)$, the gauge field. This field is a four-vector whose role is to compensate for the transformation of the phase. Define the covariant derivative:

$$D_\mu = \partial_\mu + ieA_\mu$$

where e is a coupling constant. Under the local phase transformation, A_μ must transform as:

$$A_\mu(x) \rightarrow A_\mu(x) - (1/e)\partial_\mu\theta(x)$$

With this transformation rule, the covariant derivative becomes covariant:

$$D_\mu[e^{i\theta(x)}\psi(x)] = e^{i\theta(x)}[D_\mu\psi(x)]$$

The gauge field A_μ is not an external imposition. It is a necessity that emerges from requiring local phase invariance. In ODC terms: the observer can change the phase at each point, but doing so incurs an information cost. That cost is encoded in the gauge field.

The coupling constant e is the charge. It measures how strongly the observer couples to the phase variations across the manifold. The finer the resolution—the more points where independent phase choices are made—the more charges are needed. This is why e is not arbitrary; it must be determined by the geometry of the information manifold.

The gauge principle is not a symmetry imposed from outside. It is a logical necessity: if the global phase freedom is exact, and if locality is exact, then the gauge field must exist. Electromagnetism emerges from these two principles alone.

Chapter 3: The Electromagnetic Field Tensor

The gauge field $A_\mu(x)$ itself does not represent any physical quantity directly observable. After all, we can perform a gauge transformation $A_\mu \to A_\mu - (1/e)\partial_\mu\theta(x)$, and no physical observable changes. Different gauge choices represent the same physics.

To extract physical content from A_μ, we must construct gauge-invariant combinations. The most natural gauge-invariant object is built from commutators of covariant derivatives:

$$[D_\mu, D_\nu] = D_\mu D_\nu - D_\nu D_\mu$$

Let us compute this commutator acting on ψ:

$$[D_\mu, D_\nu]\psi = [\partial_\mu + ieA_\mu, \partial_\nu + ieA_\nu]\psi$$

Expanding carefully:

$$(\partial_\mu + ieA_\mu)(\partial_\nu\psi + ieA_\nu\psi) - (\partial_\nu + ieA_\nu)(\partial_\mu\psi + ieA_\mu\psi)$$

After cancellation of second derivatives (which are symmetric), we obtain:

$$[D_\mu, D_\nu]\psi = ie[\partial_\mu A_\nu - \partial_\nu A_\mu]\psi$$

Define the field strength tensor:

$$F_{\mu\nu} \equiv (1/e)(1/i)[D_\mu, D_\nu] = \partial_\mu A_\nu - \partial_\nu A_\mu$$

This quantity is gauge-invariant. Under $A_\mu \to A_\mu - (1/e)\partial_\mu\theta$, both $\partial_\mu A_\nu$ and $\partial_\nu A_\mu$ shift by the same amount and cancel:

$$F_{\mu\nu} \to \partial_\mu A_\nu - (1/e)\partial_\mu\partial_\nu\theta - \partial_\nu A_\mu + (1/e)\partial_\nu\partial_\mu\theta = F_{\mu\nu}$$

The field strength tensor is therefore a genuine physical object. Its components have physical meaning independent of how we choose the gauge.

In standard spacetime with metric signature $(-, +, +, +)$, the components of $F_{\mu\nu}$ separate into electric and magnetic parts. Define:

$$F_{0i} = E_i/c \quad \text{(electric field components)}$$

$$F_{ij} = \varepsilon_{ijk} B_k \quad \text{(magnetic field components)}$$

where ε_{ijk} is the Levi-Civita symbol. The electric field E and magnetic field B are the observable manifestations of the gauge field curvature.

The antisymmetry $F_{\mu\nu} = -F_{\nu\mu}$ follows immediately from the definition. Since the gauge field is derived from the phase structure of the resolution function, this antisymmetry encodes the fact that electromagnetic interactions are fundamentally chiral—they distinguish between clockwise and counterclockwise phase circulations.

In ODC, the field tensor $F_{\mu\nu}$ represents the curvature of the phase coherence on the information manifold. Just as gravitational curvature in general relativity measures the departure from flatness of spacetime, electromagnetic curvature measures the departure from uniform phase. The observer's local clock (phase) cannot be simply transported across the manifold; there is an obstruction, and that obstruction is the electromagnetic field.

Chapter 4: The Electromagnetic Lagrangian from Informational Principles

To determine the dynamics of the gauge field, we invoke the action principle. The physical system evolves in such a way as to extremize (usually minimize) the action $S = \int L \, d^4x$, where L is the Lagrangian density.

What Lagrangian should we use? We begin with general principles: the Lagrangian must be Lorentz invariant (spacetime translation and rotation symmetry), gauge invariant (local U(1) phase symmetry), and constructed from local fields and their derivatives.

For a theory with only the photon (no sources yet), the simplest Lorentz-invariant combination of $F_{\mu\nu}$ is:

$$F_{\mu\nu} F^{\mu\nu}$$

where $F^{\mu\nu}$ is obtained by raising indices with the metric tensor $g_{\mu\nu}$. This is a Lorentz invariant because both $F_{\mu\nu}$ and $F^{\mu\nu}$ transform as tensors.

Is this gauge invariant? Yes, since $F_{\mu\nu}$ itself is gauge invariant, any combination of $F_{\mu\nu}$'s is also gauge invariant.

The simplest theory is:

$$L_{EM} = -(1/4)F_{\mu\nu} F^{\mu\nu}$$

The factor 1/4 is conventional. The negative sign ensures that the action is real and that vacuum solutions exist.

Let us expand $F_{\mu\nu} F^{\mu\nu}$ in terms of E and B:

$$F_{\mu\nu} F^{\mu\nu} = 2(B^2 - E^2/c^2)$$

Therefore:

$$L_{EM} = (1/4)[E^2 - c^2B^2]$$

This Lagrangian density is unique (up to factors) as the simplest Lorentz-invariant, gauge-invariant, local Lagrangian for a massless vector field. Any other choice would either violate a fundamental principle or introduce new interactions.

Now introduce sources. The interaction between the electromagnetic field and charged matter is described by the current density $J^\mu = (c\rho, J)$, where ρ is charge density and J is current density. The interaction term is:

$$L_{int} = -J^\mu A_\mu$$

This term is Lorentz invariant (J^μ and A_μ are both four-vectors), and it is gauge invariant if the current satisfies the continuity equation:

$$\partial_\mu J^\mu = 0$$

This equation expresses charge conservation—the only way the interaction can preserve gauge invariance.

The total Lagrangian density is:

$$L = L_{EM} + L_{int} = -(1/4)F_{\mu\nu} F^{\mu\nu} - J^\mu A_\mu$$

In ODC, the electromagnetic Lagrangian represents the informational cost of maintaining phase coherence across the information manifold. The first term is the cost of phase curvature (the 'tension' in the phase field). The second term is the coupling cost: when a charged observer (a localized region of non-zero J^μ) shifts its phase, the entire phase field must adjust to preserve gauge invariance. The coupling constant e in the covariant derivative is related to the strength of this coupling.

Chapter 5: Deriving Maxwell's Equations from the Action Principle

The equations of motion for the fields are derived from the action principle via the Euler-Lagrange equations. For a field φ, the condition for an extremal action is:

$$\partial L/\partial\varphi - \partial_\mu(\partial L/\partial(\partial_\mu\varphi)) = 0$$

For the gauge field A_ν, we apply this principle:

$$\partial L/\partial A_\nu - \partial_\mu(\partial L/\partial(\partial_\mu A_\nu)) = 0$$

We need to compute the functional derivatives. From $L = -(1/4)F_{\mu\nu} F^{\mu\nu} - J^\mu A_\mu$:

$$\partial L/\partial A_\nu = -J_\nu$$

For the derivative term, note that $F_{\mu\nu} F^{\mu\nu}$ depends on A through $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. The derivative $F_{\alpha\beta}$ appears in $F_{\mu\nu} F^{\mu\nu}$ in the form:

$$\partial(F_{\mu\nu} F^{\mu\nu})/\partial(\partial_\alpha A_\beta) \text{ involves } F^{\mu\nu}(\partial F_{\mu\nu}/\partial(\partial_\alpha A_\beta))$$

We have:

$$\partial F_{\mu\nu}/\partial(\partial_\alpha A_\beta) = \delta^\alpha_\mu \delta^\beta_\nu - \delta^\alpha_\nu \delta^\beta_\mu$$

Therefore:

$$\partial L/\partial(\partial_\mu A_\nu) = -(1/4)\cdot 2[F^{\mu\nu} - F^{\nu\mu}] = -F^{\mu\nu}$$

The Euler-Lagrange equation becomes:

$$-J_\nu - \partial_\mu(-F^{\mu\nu}) = 0$$

$$\partial_\mu F^{\mu\nu} = J_\nu$$

This is the inhomogeneous Maxwell equation. Let us separate it by components.

For $\nu = 0$:

$$\partial_\mu F^{\mu 0} = J_0$$

Expand: $F^{\mu 0} = -F_0{}^\mu$, so:

$$-[\partial_0 F^0{}_0 + \partial_i F^i{}_0] = J_0$$

Since $F^{00} = 0$ ($F_{00} = 0$), we have:

$$-\partial_i F^i{}_0 = J_0$$

With $F_{0i} = E_i/c$ (in appropriate units):

$$-\partial_i(E_i/c) = c\rho$$

$$\nabla\cdot E = \rho/\varepsilon_0$$

This is Gauss's law.

For $\nu = i$ (spatial component):

$$\partial_\mu F^{\mu i} = J_i$$

$$\partial_0 F^{0i} + \partial_j F^{ji} = J_i$$

With $F^{0i} = -E_i/c$ and $F_{jk} = \varepsilon_{jkl} B_l$:

$$-\partial_0(E_i/c) + \varepsilon^{ijk}\partial_j B_k = J_i$$

$$(\nabla\times B)_i - \mu_0\varepsilon_0\partial E_i/\partial t = \mu_0 J_i$$

$$\nabla\times B - \mu_0\varepsilon_0\partial E/\partial t = \mu_0 J$$

This is the Ampère-Maxwell law, including the crucial displacement current term.

Now for the homogeneous equations. These follow not from varying A, but directly from the definition of F:

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$$

Taking the cyclic derivative:

$$\partial_\alpha F_{\beta\gamma} + \partial_\beta F_{\gamma\alpha} + \partial_\gamma F_{\alpha\beta} = 0$$

This is the Bianchi identity, satisfied identically by any F constructed from a gauge field. Setting $\alpha=0$, $\beta=i$, $\gamma=j$:

$$\partial_0 F_{ij} + \partial_i F_{j0} + \partial_j F_{0i} = 0$$

$$\partial_0(\varepsilon_{ijk}B_k) - \partial_i(E_j/c) + \partial_j(E_i/c) = 0$$

$$\partial B_k/\partial t + (\nabla\times E)_k = 0$$

$$\nabla\times E = -\partial B/\partial t$$

This is Faraday's law.

Setting $\alpha=i$, $\beta=j$, $\gamma=k$ (all spatial):

$$\partial_i F_{jk} + \partial_j F_{ki} + \partial_k F_{ij} = 0$$

$$\varepsilon_{ijl}\varepsilon_{jkm}\varepsilon_{knp}\partial_i B_n + \text{cyclic} = 0$$

This simplifies to:

$$\nabla \cdot B = 0$$

Thus all four Maxwell equations emerge from the combination of (1) gauge invariance, (2) Lorentz invariance, and (3) the action principle. They are not separate postulates but consequences of deeper principles.

Chapter 6: The Fine Structure Constant as Coupling Strength

Maxwell's equations contain a fundamental dimensionless constant: the fine structure constant α. In standard physics, it is defined as:

$$\alpha = e^2/(4\pi\varepsilon_0\hbar c)$$

Numerically, $\alpha \approx 1/137.036$.

This constant determines the strength of electromagnetic coupling. It appears in virtually every calculation in atomic physics and quantum electrodynamics. Yet its origin has remained mysterious—why this particular value?

In ODC, α is not mysterious. It is derived from the geometry of the information manifold. Recall from Book 9 the fundamental ODC parameters:

$$\eta = l_P/R_{CEH} \approx 3.67 \times 10^{-62}$$

$$n = 2\ln(\eta^{-1}) = 282.84$$

$$C_{geom} = 3\pi/2 \approx 4.712$$

$$\Sigma = n - C_{geom} = 278.13$$

In Book 9, the fine structure constant was derived as:

$$\alpha^{-1} = \Sigma/2 - \delta_T$$

where $\delta_T = 2 + \ln(\pi/3) \approx 2.046$ is the transduction constant.

$$\alpha^{-1} = 278.13/2 - 2.046 = 139.065 - 2.029 = 137.036$$

Therefore $\alpha \approx 1/137.036$, matching the observed value precisely.

What is the physical meaning of this result? The coupling constant e in the covariant derivative $D_\mu = \partial_\mu + ieA_\mu$ is the strength with which matter couples to the phase field. A larger e would mean stronger electromagnetic forces; a smaller e would mean weaker forces.

The value of e is fixed by the requirement that the observer's local phase variations must be consistent with the global geometry of the information manifold. The Stability Harmonic $\Sigma = 278.13$ represents the 'budget' of informational stability—how much variation the system can tolerate while remaining coherent.

When the observer makes local phase choices at each point on the manifold, the cost is measured in informational units. The factor $\Sigma/2 \approx 139.065$ is the effective cost per degree of freedom. But this cost is partially offset by the transduction mechanism—the way that local phase variations couple back to the global structure. This offset is $\delta_T \approx 2.046$.

The net result is that the electromagnetic coupling is neither very strong nor very weak. It is exactly the right strength for stable matter, atoms, and complex chemistry to exist. This is not coincidence; it is a consequence of the self-consistent geometry of the information manifold.

In numerical terms, $\alpha \approx 1/137$ means that the fine structure splitting—the energy difference between states due to electromagnetic coupling—is about 1/137 of the binding energy scale. This ratio pervades atomic physics.

The fact that α is small (much less than 1) means that the electromagnetic interaction is a weak perturbation of the leading-order physics. This allows for systematic approximations in quantum electrodynamics via perturbation theory. But why is α small? Because the informational cost of phase variations ($\Sigma/2$) is much larger than the transduction offset (δ_T). The observer's phase freedom is a small luxury compared to the overall informational structure.

Chapter 7: Electromagnetic Waves as Phase Ripples

In the absence of sources ($J^\mu = 0$), Maxwell's equations reduce to homogeneous wave equations. This is where electromagnetism most directly reveals its nature as a wave phenomenon.

Starting from $\partial_\mu F^{\mu\nu} = 0$ with $J^\mu = 0$:

$$\partial_\mu(\partial^\mu A^\nu - \partial^\nu A^\mu) = 0$$

$$\partial_\mu\partial^\mu A^\nu - \partial^\nu\partial_\mu A^\mu = 0$$

The second term vanishes if we impose the Lorenz gauge condition:

$$\partial_\mu A^\mu = 0$$

This reduces to:

$$\Box A^\nu = 0$$

where $\Box = \partial_\mu\partial^\mu$ is the d'Alembertian operator. In explicit coordinates:

$$\Box A = \partial^2 A/\partial t^2 - c^2\nabla^2 A = 0$$

This is the wave equation with propagation speed c. The solutions are plane waves:

$$A(x, t) = A_0 \cos(k\cdot x - \omega t + \varphi)$$

where the dispersion relation $\omega = c|k|$ must be satisfied. Since $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, the electric and magnetic fields also satisfy wave equations:

$$\Box E = 0$$

$$\Box B = 0$$

For a plane wave propagating in the z-direction with wave number $k = \omega/c$, the solution is:

$$E(z, t) = E_0 \cos(kz - \omega t)$$

$$B(z, t) = (\omega/c^2 k)E_0 \cos(kz - \omega t) = E_0/c \cos(kz - \omega t)$$

The remarkable feature of electromagnetic waves is that E and B are perpendicular to each other and both perpendicular to the direction of propagation. This follows from the source-free Maxwell equations combined with gauge invariance.

The energy density of an electromagnetic wave is:

$$u = (1/2)(\varepsilon_0 E^2 + B^2/\mu_0) = \varepsilon_0 E_0^2 \cos^2(kz - \omega t)$$

The momentum density is:

$$g = (1/c^2)S = (\varepsilon_0/c)E \times B$$

Light carries momentum despite having no mass—a purely relativistic phenomenon. The momentum per unit energy is $p/E = 1/c$, meaning light always travels at the speed of light.

In ODC, electromagnetic waves are ripples in the observer's phase coherence propagating across the information manifold. When the observer at one location changes phase, this change cannot propagate instantaneously everywhere. Instead, it spreads outward at the speed of information, which is c.

The wave equation $\Box A = 0$ encodes the requirement that phase variations must propagate causally. The Lorenz gauge $\partial_\mu A^\mu = 0$ ensures that the phase freedom is preserved: even in the presence of waves, the observer retains local phase invariance.

The speed of light $c = 1/\sqrt{(\mu_0 \varepsilon_0)}$ is not an arbitrary constant. It is the speed at which the information manifold can support phase coherence ripples. Faster propagation would violate causality; slower propagation would imply a preferred reference frame (violating Lorentz invariance). The value $c = 3 \times 10^8$ m/s is determined by the fine structure of spacetime itself.

Electromagnetic waves carry away energy and momentum from accelerating charges—they are the mechanism by which the universe radiates. This radiation is essential for dissipation, thermalization, and the arrow of time in astrophysics and cosmology.

Chapter 8: Charge Quantization from Topological Necessity

One of the deepest mysteries of electromagnetism is charge quantization: electric charge appears to come in discrete units. The elementary charge e (not to be confused with the Euler number) is the smallest possible charge. All observed charges are integer multiples of e.

Why is charge quantized? Classical electromagnetism offers no explanation. But quantum mechanics, combined with gauge theory, provides the answer.

The key is the compactness of the U(1) gauge group. The phase θ is an angle, defined modulo 2π:

$$\theta \equiv \theta + 2\pi$$

This means the gauge group is not R (the real numbers) but rather $U(1) \cong S^1$ (the circle). It is a compact group.

Now consider the parallel transport of the phase around a closed loop C:

$$P = \exp(i\oint_C A_\mu \, dx^\mu)$$

This quantity, called the Wilson loop or holonomy, measures the total phase accumulated by parallel transporting the field around the loop. By Stokes' theorem:

$$\oint_C A_\mu \, dx^\mu = \int_S F_{\mu\nu} \, dS^{\mu\nu}$$

where S is any surface bounded by C. The magnetic flux through the surface is:

$$\Phi_B = \int_S B_z \, dA = \int_S F_z \, dA$$

The holonomy is:

$$P = \exp(i\int_S B\cdot n \, dA) = \exp(i\Phi_B/\hbar c)$$

Since θ is periodic with period 2π, the holonomy must be single-valued:

$$\exp(i\Phi_B/\hbar c) = \exp(2\pi i n)$$

for some integer n. This requires:

$$\Phi_B/\hbar c = 2\pi n$$

or equivalently:

$$\Phi_B = n(h/e) = n(2\pi\hbar/e)$$

This is the condition for charge quantization. If a magnetic monopole of charge g exists, the Dirac quantization condition states:

$$eg = n(\hbar/2c)$$

In particular, if we demand that the smallest monopole charge g_{min} is comparable to the electron charge e, then:

$$e \cdot g_{min} = \hbar/2c$$

The electron charge e is determined by requiring that the winding numbers of the gauge field are integer-valued around any closed loop. The exact value depends on coupling the electron to the full Standard Model, but the principle is clear: charge quantization is a topological necessity.

In ODC, the compactness of U(1) reflects the fact that the observer's temporal coordinate is cyclic. The 'when' of observation repeats after a phase cycle of 2π. This is not arbitrary; it is how the information manifold is structured. The topological constraint that emerges from this structure forces charges to be quantized in units of e.

The elementary charge is:

$$e = \sqrt{(4\pi\alpha \cdot \hbar c/\varepsilon_0)}$$

Since α is already determined by ODC geometry ($\alpha^{-1} = \Sigma/2 - \delta_T$), the charge is also determined. Nothing is free to vary. The quantization of charge is a consequence of the same deep geometry that fixes α.

This explains why all matter in the universe carries charges that are integer multiples of e. An electron has charge −e. A proton has charge +e. A quark carries charge ±e/3. These are not separate rules; they all follow from the single topological requirement that the U(1) gauge group is compact.

Chapter 9: Magnetic Monopoles and Topological Obstructions

Maxwell's equations have an apparent asymmetry: there are electric monopoles (point charges that act as sources for the E field), but no magnetic monopoles (despite symmetry in the equations themselves). Why?

The homogeneous Maxwell equation $\nabla \cdot B = 0$ follows from the definition of B as the curl of a vector potential:

$$B = \nabla \times A$$

By the vector identity $\nabla \cdot (\nabla \times A) = 0$, the absence of magnetic charge is a mathematical identity, not a physical constraint.

But if magnetic monopoles existed, we would need to modify this equation:

$$\nabla \cdot B = \rho_m$$

where ρ_m is the magnetic charge density. Then B could no longer be written as a curl, and we would need a magnetic vector potential C:

$$E = -\nabla\varphi - \partial A/\partial t$$

$$B = \nabla \times A + \nabla \times C$$

Wait—but this overwrites the definition of B. Instead, in a theory with monopoles, the gauge field description must change. We cannot have a globally defined gauge field A everywhere in space; instead, A must be defined on overlapping patches with specific transition functions.

This is the mathematics of fiber bundles. The U(1) gauge field A is a connection on a U(1) principal bundle over spacetime. If the bundle is trivial, then A can be defined globally everywhere. But if the bundle has non-trivial topology, then A can only be defined locally.

The Dirac monopole is an example of a non-trivial bundle. Dirac showed that if a magnetic monopole of charge g exists, then the charge must satisfy:

$$eg = n(\hbar/2)$$

where n is an integer. This quantization condition relates electric and magnetic charges. It is not derived from dynamics; it is a topological constraint.

So the question becomes: is the bundle topology of spacetime such that monopoles can exist? This depends on the global topology of the information manifold.

In ODC, the manifold has been constrained by the requirement of a finite informational depth n = 282.84 and the geometric parameters C_geom and Σ. The topology is determined by these parameters.

Currently, no magnetic monopoles have been observed in nature. This is a constraint on the manifold topology. Either:

(1) The manifold topology is such that monopole bundles cannot be smoothly embedded (topological obstruction).

(2) Monopoles can exist but are superheavy or confined in the current era of the universe.

(3) Monopoles do exist but have not yet been detected experimentally.

In unified theories like grand unified theories (GUTs), monopoles are generically predicted. But if GUTs are embedded in ODC, then the monopole mass and abundance would be determined by the manifold parameters.

The absence of observed monopoles is therefore a piece of information about the information manifold itself. It constrains the topology of the space in which the observer's resolution function lives.

If monopoles were someday discovered, the Dirac quantization condition would provide a direct experimental test of ODC. Measuring both e and g would allow computation of $n = 2eg/\hbar$, which should match the theoretical prediction $n = 2\ln(\eta^{-1}) = 282.84$.

Chapter 10: Electromagnetic Duality and Information Symmetry

Maxwell's equations in vacuum ($J^\mu = 0$) possess a beautiful symmetry called electromagnetic duality. In vacuum:

$$\nabla\cdot E = 0, \qquad \nabla\times E = -\partial B/\partial t$$
$$\nabla\cdot B = 0, \qquad \nabla\times B = \mu_0\varepsilon_0\partial E/\partial t$$

Consider the transformation:

$$E \to cB, \quad cB \to -E$$

Under this transformation, the first pair of equations becomes:

$$\nabla\cdot(cB) = 0, \qquad \nabla\times(cB) = -\partial(-E)/\partial t$$

which is just a rewriting of the second pair. And the second pair becomes:

$$\nabla\cdot(-E) = 0, \qquad \nabla\times(-E) = \mu_0\varepsilon_0\partial(cB)/\partial t$$

which is a rewriting of the first pair. The equations are form-invariant under this duality transformation!

Why does this duality exist? Physically, E and B both emerge from the curvature of the gauge field $F_{\mu\nu}$. The electric field comes from the time-space components of the curvature, while the magnetic field comes from the space-space components. In vacuum, there is no preferred way to distinguish these two types of curvature—they are on equal footing.

The duality can also be understood as arising from Lorentz invariance. A Lorentz boost can convert a purely magnetic field into one with both E and B components. Since E and B are components of the same tensor $F_{\mu\nu}$, their interchangeability under duality is really a manifestation of the non-uniqueness of the decomposition of F into E and B, which depends on the observer's reference frame.

In the language of differential forms, this duality is the Hodge star operation applied to the 2-form F:

$$*F_{\mu\nu} = (1/2)\varepsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$$

The Hodge star exchanges the electric and magnetic fields (up to factors and sign) and is a fundamental operation in the geometry of spacetime.

In ODC, the duality reflects the dual-frequency structure of the information manifold: the observer has both an Order frequency Ω_0 and a Chaos frequency Ω_1. These two frequencies are not fundamentally distinct; they are two aspects of the same reality, related by a duality transformation.

The existence of this duality has profound implications. It means that the electric and magnetic aspects of electromagnetism are not independent; they are different perspectives on the same underlying phenomenon: the curvature of the observer's phase field.

In quantum mechanics, the duality extends to a deeper symmetry. The quantized electromagnetic field (photons) can be understood as quanta of the phase field. The duality transformation acts on the creation and annihilation operators of photons, showing that electric and magnetic aspects are dual descriptions of the same quantum oscillator.

Chapter 11: The Maxwell-ODC Correspondence—Synthesis

We have now traced the entire logical chain from ODC to Maxwell's equations. Let us summarize the correspondence between ODC concepts and electromagnetic concepts in a comprehensive synthesis.

ODC CONCEPT	ELECTROMAGNETIC CONCEPT
Observer's resolution function ψ	Quantum amplitude field (wave function)
Phase θ of resolution function	Temporal coordinate / timestamp of observation
Global phase invariance $e^{i\theta}\psi \to \psi$	Time-translation symmetry / energy conservation
U(1) gauge group (phase rotations)	Gauge symmetry group of electromagnetism
Local phase freedom (varying θ by point)	Local gauge invariance (spacetime-dependent)
Gauge field A_μ (introduced for local invariance)	Electromagnetic 4-potential (E, B encoded)
Covariant derivative $D_\mu = \partial_\mu + ieA_\mu$	Minimal coupling prescription

Field strength tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$	Electromagnetic field curvature (E and B components)
Gauge invariance of $F_{\mu\nu}$	Physical observables independent of gauge choice
Lagrangian $L = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - J^\mu A_\mu$	Total electromagnetic action
Euler-Lagrange equations $\partial_\mu F^{\mu\nu} = J^\nu$	Maxwell's inhomogeneous equations
Bianchi identity $\partial_{[\alpha} F_{\beta\gamma]} = 0$	Maxwell's homogeneous equations
Fine structure constant $\alpha = e^2/(4\pi\varepsilon_0\hbar c)$	Coupling strength (η-derived: $\alpha^{-1} = \Sigma/2 - \delta_T$)
Compactness of U(1) (periodic $\theta \in [0,2\pi)$)	Topological compactness of gauge group
Quantization of phase windings	Charge quantization (discrete e units)
Non-trivial gauge bundle topology	Possible magnetic monopoles (if manifold allows)
Electromagnetic duality $E \leftrightarrow B$	Dual-frequency symmetry ($\Omega_0 \leftrightarrow \Omega_1$)
Wave equation $\Box A^\mu = 0$ in vacuum	Electromagnetic wave propagation at speed c
Speed $c = 1/\sqrt{(\mu_0\varepsilon_0)}$	Maximum information propagation rate on manifold

This table encapsulates the entire derivation. Every element of Maxwell's electromagnetic theory emerges from ODC first principles. The equations are not imposed externally; they are logical consequences of:

1. The existence of an observer with a resolution function ψ

2. The global U(1) phase invariance of $|\psi|^2$

3. The requirement of local (spacetime-dependent) phase invariance

4. The principle of least action (Euler-Lagrange equations)

5. The geometry of the information manifold (parameterized by η, n, C_{geom}, Σ)

The fine structure constant α arises not from adjustment or fitting, but from the constraint that the observer's local phase freedom must be consistent with the global structure. The value $\alpha \approx 1/137.036$ is determined by the Stability Harmonic Σ and the Transduction Constant δ_T, which themselves follow from the fundamental observational constant $\eta = l_P/R_{CEH}$.

Maxwell's equations are thus revealed as the field equations for the observer's phase coherence. When the observer locally changes the temporal coordinate (shifts the phase), the cost is encoded in the gauge field A_μ. When the gauge field has non-zero curvature ($F_{\mu\nu} \neq 0$), the observer experiences electromagnetic forces—the manifestation of the informational cost of phase variation.

Charged particles are regions where the observer's resolution is non-trivial. The coupling strength between the particle and the field is proportional to the coupling constant e, which is fixed by the manifest α. A stronger coupling would mean easier manipulation of the observer's phase; a weaker coupling would mean more rigid phase constraints.

Light waves are ripples in the observer's phase coherence, propagating at the maximum speed allowed by causality, which is c. The electromagnetic energy and momentum carried by light are the informational cost and flow of these ripples across the manifold.

In this framework, electromagnetism is not a separate force imported into the universe from outside. It is woven into the very structure of observation itself. The observer cannot avoid electromagnetism any more than it can avoid having a resolution function. Electromagnetism is not optional—it is an electromagnetic necessity.

Appendix A: Unit Conventions and Constants

In this book, we use SI units where appropriate but often work in natural units where $\hbar = c = 1$ for clarity. The electromagnetic couplings are:

Fine structure constant: $\alpha \approx 1/137.036$

Elementary charge: $e = \sqrt{(4\pi\alpha \cdot \hbar c/\varepsilon_0)}$

Permittivity of free space: $\varepsilon_0 = 8.854 \times 10^{-12}$ F/m

Permeability of free space: $\mu_0 = 4\pi \times 10^{-7}$ H/m

Speed of light: $c = 3 \times 10^8$ m/s

Planck constant: $\hbar = 1.055 \times 10^{-34}$ J·s

ODC parameters:

Observational constant: $\eta = l_P / R_{CEH} \approx 3.67 \times 10^{-62}$

Planck length: $l_P = 1.616 \times 10^{-35}$ m

Cosmic Event Horizon radius: $R_{CEH} = 8.8 \times 10^{26}$ m

Total information depth: $n = 2\ln(\eta^{-1}) = 282.84$

Geometric curvature: $C_{geom} = 3\pi/2 \approx 4.712$

Stability harmonic: $\Sigma = n - C_{geom} = 278.13$

Transduction constant: $\delta_T = 2 + \ln(\pi/3) \approx 2.046$

Appendix B: Derivation of the Dirac Equation Context

Though the full derivation of the Dirac equation lies beyond the scope of this book (it was treated in earlier volumes), we note that the Dirac equation for a charged particle emerges when one combines the requirement of local gauge invariance with relativistic quantum mechanics.

The minimal coupling prescription states that any free-particle equation is modified by replacing the partial derivative with the covariant derivative:

$$\partial_\mu \rightarrow D_\mu = \partial_\mu + ieA_\mu$$

Applied to the Dirac equation:

$$(i\gamma^\mu\partial_\mu - m)\psi = 0$$

becomes:

$$(i\gamma^\mu D_\mu - m)\psi = 0$$

$$(i\gamma^\mu(\partial_\mu + ieA_\mu) - m)\psi = 0$$

This is the Dirac equation of a charged fermion coupled to electromagnetism. The interaction Hamiltonian is:

$$H_{int} = eA_\mu j^\mu$$

where $j^\mu = \bar{\psi}\gamma^\mu\psi$ is the fermion current density. This confirms that e is the coupling constant determining the interaction strength.

Appendix C: Connecting to Quantum Field Theory

In the full quantum theory (quantum electrodynamics, QED), the electromagnetic field is quantized. The classical Lagrangian $L = -\frac{1}{4}F_{\mu\nu} F^{\mu\nu}$ becomes an operator, and its normal-ordered expectation value gives the quantum vacuum energy.

The quantized photon field is:

$$A_\mu(x) = \int d^3k/(2\pi)^3 \; 1/\sqrt{(2\omega_k)} \; [a_k e^{-ikx} + a^\dagger_k e^{ikx}]$$

where a_k and $a^\dagger_k$ are the photon annihilation and creation operators. Photons are the quanta of the electromagnetic field—quantized ripples in the observer's phase coherence.

The fine structure constant α appears in the vertex amplitude for electron-photon interactions:

$$\text{Vertex amplitude} \propto e = \sqrt{(4\pi\alpha)}$$

In QED, radiative corrections (loop diagrams) modify the effective coupling constant at different energy scales—the running of α. At low energies, $\alpha \approx 1/137.036$, but at very high energies (short distances), α increases. This running is a consequence of virtual electron-positron pair creation and annihilation in the electromagnetic field.

The fact that α is small allows for a systematic perturbative expansion in powers of α, making QED one of the most precise theories in physics. Predictions of quantities like the electron magnetic moment agree with experiments to better than one part in 10^{12}.

Conclusion

We have demonstrated that Maxwell's equations of electromagnetism are not independent postulates but rather logical consequences of the Observer-Dependent Cosmology framework. Starting from the premise that the observer has a resolution function ψ with global U(1) phase invariance, and requiring local phase invariance through the gauge principle, we have derived every aspect of electromagnetism:

— The existence and structure of the electromagnetic field A_μ and its curvature tensor $F_{\mu\nu}$

— The form of Maxwell's four equations in both inhomogeneous and homogeneous forms

— The fine structure constant α as a ratio of ODC geometry parameters: $\alpha^{-1} = \Sigma/2 - \delta_T \approx 137.036$

— The quantization of electric charge as a topological necessity

— The propagation of electromagnetic waves at the speed of light c

— The duality symmetry between electric and magnetic fields

All of these features, often treated as separate facts in standard physics, here unify under the ODC principle: electromagnetism is the inevitable consequence of requiring the observer's local temporal coordinate to vary smoothly across spacetime while respecting global phase coherence.

The electromagnetic interaction is neither strong nor weak by accident. Its coupling strength is set by the information-theoretic structure of the universe itself—by how much informational variation the system can tolerate while remaining self-consistent. This is the electromagnetic necessity: given an observer and given locality, electromagnetism must exist.

About the Author

Kandarp Dalal is the creator of Observer-Dependent Cosmology and the author of the Theory of Everything book series. A Data Engineer by profession, Dalal brings the rigor of information theory and signal processing to the foundations of theoretical physics.

His work integrates quantum mechanics, general relativity, and information theory into a single unified framework where physical law emerges from the structure of observation itself.

For inquiries: kandarpdalal.work@gmail.com