Fiscal Year:	FY 2009	**Task Last Updated:**	FY 12/09/2009
PI Name:	Shi, Yufang D.V.M., Ph.D.		
Project Title:	Apoptosis and Immune Homeostasis During Hindlimb Unloading		
Division Name:	Human Research		
Program/Discipline:	NSBRI		
Program/Discipline--Element/Subdiscipline:	NSBRI--Radiation Effects Team		
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	(1) **HHC**:Human Health Countermeasures		
Human Research Program Risks:	None		
Space Biology Element:	None		
Space Biology Cross-Element Discipline:	None		
Space Biology Special Category:	None		
PI Email:	shiyu@umdnj.edu	**Fax:**	FY 732 235 4505
PI Organization Type:	UNIVERSITY	**Phone:**	732-235-4501
Organization Name:	Robert Wood Johnson Medical School		
PI Address 1:	675 Hoes Lane		
PI Address 2:			
PI Web Page:			
City:	Piscataway	**State:**	NJ
Zip Code:	08854	**Congressional District:**	6
Comments:			
Project Type:	Ground	**Solicitation / Funding Source:**	2003 Biomedical Research & Countermeasures 03-OBPR-04
Start Date:	04/01/2005	**End Date:**	06/30/2009
No. of Post Docs:	1	**No. of PhD Degrees:**	0
No. of PhD Candidates:	1	**No. of Master' Degrees:**	0
No. of Master's Candidates:	0	**No. of Bachelor's Degrees:**	0
No. of Bachelor's Candidates:	0	**Monitoring Center:**	NSBRI
Contact Monitor:		**Contact Phone:**	
Contact Email:			
Flight Program:			
Flight Assignment:	NOTE: End date now 6/30/2009 (previously was 3/31/2009) per NSBRI data submission 12/2009.		
Key Personnel Changes/Previous PI:			
COI Name (Institution):	Roberts, Arthur (University of Med & Dentistry of NJ) Xu, Guangwu (Robert Wood Johnson Medical School)		
Grant/Contract No.:	NCC 9-58-IIH00405		
Performance Goal No.:			
Performance Goal Text:			

Task Description:

Space flight impacts the immune system, as it is often results in lymphopenia and abnormally low responses to antigenic stimulation. These changes are believed to be due to factors such as microgravity, lack of load-bearing, psychological and physical stress, and exposure to radiation, but the exact mechanisms are unknown. Hindlimb unloading (HU) in rodents, a ground-based animal model, has been used to simulate some of the factors associated with space flight. Interestingly, HU induces similar changes in the immune system of rodents as those resulting from space flight in humans. Based on our previous studies, we hypothesize that some of these changes are modulated through the induction of apoptosis in lymphocytes. We have proposed three specific aims: 1) Elucidate the role of Fas and endogenous opioids in the modulation of the immune system by HU and radiation; 2) Determine the effect of HU and radiation on lymphocyte dynamics and immune responses, and, 3) because autoimmune disease, paradoxically, is often associated with stress and radiation exposure, determine whether HU and radiation promote autoimmunity. The overall goal of these studies is to determine the mechanisms underlying apoptosis-related immune modulation using ground-based models.

PROGRESS

1. In a study at the NASA Space Radiation Laboratory at Brookhaven National Laboratory (BNL), we examined the acute effects of whole body HZE-particle irradiation. Mice were subjected to high-energy Fe-56 ions and the effects on various lymphocyte populations were determined. We found that lymphocyte numbers in the spleen and thymus were reduced in a dose-dependent manner (using doses of 200, 100 and 50 cGy), with thymocytes being more sensitive than splenocytes. Subpopulation analysis revealed that all subsets declined in number, except for CD4+CD25+ regulatory T cells. To examine the mechanism, mice deficient in apoptosis-inducing genes were tested: deficiency in Fas ligand (gld mutation) or TRAIL (TNF-related apoptosis-inducing ligand) did not prevent lymphocyte losses, indicating that these apoptosis pathways are not involved in particle radiation-induced lymphocyte losses. In contrast, gamma-radiation-induced cell depletion does require FasL and TRAIL. Thus, there are divergent pathways of apoptosis induction by particle radiation vs. gamma-radiation. Using specific inhibitors in vitro, we also found that particle radiation-induced lymphocyte apoptosis is caspase-dependent, but does not utilize nitric oxide or oxygen free radicals.

2. We have reported that endogenous granzyme B (GzmB) is a critical regulator of the balance between Th1/Th2 helper T cells. Highly expressed in Th2 cells, GzmB is required to induce apoptosis in these cells. We have found that GzmB-deficient mice develop stronger Th2-type immune responses. Since GzmB expression can be inhibited by steroid hormones, and corticosteroid levels are often elevated during stress, these findings may explain the increased levels of Th2 cytokines found in astronauts post-flight. Mice subjected to HU also have increased Th2 responses and serum glucocorticoids. We are still examining the role of GzmB in immune responses of mice subjected to HU.

3. Stress, microgravity and space radiation are well-known to induce apoptosis in lymphoid and other cells. Although apoptotic cells can induce immunosuppression, the molecular mechanisms are unknown. We found that exposure to apoptotic cells rendered dendritic cells (DC) capable of producing much greater amounts of nitric oxide in response to exogenous IFN-gamma compared to normal DC. We believe that these results have important ramifications for the understanding of space flight induced immunosuppression.

4. Mesenchymal stem cells (MSCs) are a unique population of multi-potent stem cells that are derived from adult bone marrow. These cells are immunosuppressive, and also seem to protect splenic and thymic lymphocytes from the deleterious effects of microgravity and radiation. We found that transfusing mice with MSCs immediately after gamma-irradiation or before hindlimb unloading significantly reduced splenocyte losses. In vitro, irradiated lymphocytes were protected from apoptosis when co-cultured with MSCs. These results indicate that MSCs provide survival signals that prevent lymphocyte apoptosis. In fact, we have found several such MSC-produced cytokines, which may allow for development of pharmaceutical interventions to prevent lymphocyte loss during space flight.

5. We examined the interaction between the immune and skeletal systems. Osteopontin (OPN), a bone matrix protein, has also been shown affect the immune system--as a cytokine and possibly as a transcription factor. We examined the role of OPN in HU-induced lymphocyte depletion using OPN-knockout mice, collaborating with David Denhardt and Kathryn Wang. We have reported that OPN deficiency confers resistance to HU-induced thymocyte loss. This protective effect derives from decreased corticosteroid levels in OPN-deficient mice, indicating that OPN is important in regulating the steroid response during HU. This suggests a similar role for OPN in the response to space flight conditions. Importantly, injection of a specific anti-OPN antibody (2C5) into wild-type mice ameliorates organ atrophy induced by chronic restraint; changes in corticosterone levels were also partially reversed. These studies reveal that circulating OPN plays a significant role in the regulation of the hypothalamus-pituitary-adrenal axis hormones, and that it augments chronic restraint-induced organ atrophy.

FUTURE PLANS

1. Continue ongoing studies of the molecular mechanisms that contribute to the cell survival/ radiation-protective effects of MSCs.

2. Examine the effects on the immune parameters of mice after 3 months of space flight, using control and osf-1 transgenic mice, and compare these results to our HU studies.

Rationale for HRP Directed Research:

Research Impact/Earth Benefits:

We hope to gain an understanding of the mechanisms by which stress affects immune homeostasis. With this information, it may be possible to devise dietary and lifestyle changes that would reduce the deleterious effects of stress on the immune system. Such changes may lead to an improvement in public health. We also hope to identify food supplements, such as vitamin C and seaweed extracts, that may reduce the health impact of stress, bringing benefit to the general population, especially those who suffer from high levels of occupational or life stress.
By determining the role of cellular apoptosis in mediating the deleterious effects of stress on the immune system we may be able to aid in the development of therapies for the maintenance of immune tolerance for clinical use in organ transplant or autoimmune disease.

We are studying the role of a specific subset of T lymphocytes called CD4+CD25+ regulatory T cells, which seem to be involved in the deleterious effect of stress on the immune system. Learning how these cells exert their effects may lead to potential interventions to prevent the negative health impacts of autoimmunity, cancer, and other diseases that arise from an improperly regulated immune system. It is difficult to measure the level of stress that an individual experiences

	for the purpose of allowing appropriate intervention. Identification of particular serum proteins whose concentration is related to the stress level of an individual could serve as a biomarker for the assessment of stress levels for medical or occupational purposes.
Task Progress:	1. Elucidate the role of Fas and endogenous opioids in the modulation of the immune system by hindlimb unloading (HU) and radiation: We have found that the synergistic loss of lymphocytes caused by gamma-radiation and HU is exerted through Fas. Our studies also revealed a role for TRAIL in these effects. We have now shown that osteopontin (OPN), a bone matrix protein, can modulate lymphocyte apoptosis during HU by reducing stress-induced glucocorticoid production. OPN-deficient mice were resistant to HU-induced thymocyte depletion, revealing a novel function of OPN. 2. Determine the effect of HU and radiation on lymphocyte dynamics and immune responses: We found that whole-body irradiation of mice with low-dose Fe-56 ion radiation (50, 100, 200 cGy) depletes splenic and thymic lymphocytes, with thymocytes being more sensitive than splenocytes. All lymphocyte subsets in spleen and thymus were affected, except for the CD4+CD25+ regulatory T cells, which were quite resistant. T cells and B cells were equally sensitive, while CD8+ T cells were more sensitive than CD4+ T cells. In the thymus, immature CD4+CD8+ (double-positive) thymocytes are exquisitely sensitive, mature single-positive cells were less sensitive, and immature double-negative cells were nearly completely resistant to these doses. These findings reveal for the first time the differential susceptibility of different lymphocyte subpopulations to particle radiation-induced deletion. This particle radiation-induced lymphocyte apoptosis was independent of Fas and TRAIL (TNF-related apoptosis-inducing ligand), the most prevalent receptor-induced modes of apoptosis, since deficiency in these genes did not prevent lymphocyte losses. This is striking, since lymphocyte depletion following gamma-irradiation depends on the Fas-FasL pathway. Thus, there are distinct pathways of apoptosis induction by particle radiation compared to gamma-radiation. Furthermore, we found that particle radiation-induced lymphocyte apoptosis is caspase-dependent, but does not require nitric oxide or oxygen free radicals. 3. Determine whether HU and radiation promote the development of autoimmunity: We found that unlike apoptotic cells, which promote immune tolerance, necrotic cells actually stimulate an immune response. In early studies on the development of autoimmunity following HU and concomitant irradiation, our findings as yet are inconclusive. Our molecular studies have shown that apoptotic cells cause dendritic cells (DC) to lose their ability to promote T cell production of IFN-g, while at the same time these DCs become highly sensitive to induction of nitric oxide production in response to IFN-g. The nitric oxide produced by apoptotic cell-primed DCs can in turn suppresses T cell responses. When apoptotic cells are too abundant, however, they can undergo secondary necrosis. Therefore, efficient removal of apoptotic cells is crucial in preventing autoimmunity in radiation-exposed individuals.
Bibliography Type:	Description: (Last Updated: 11/19/2010)
Articles in Peer-reviewed Journals	Das J, Ren G, Zhang L, Roberts AI, Zhao X, Bothwell AL, Van Kaer L, Shi Y, Das G. "Transforming growth factor beta is dispensable for the molecular orchestration of Th17 cell differentiation." J Exp Med. 2009 Oct 26;206(11):2407-16. PMID: 19808254 , Oct-2009
Articles in Peer-reviewed Journals	Ren G, Su J, Zhang L, Zhao X, Ling W, L'huillie A, Zhang J, Lu Y, Roberts AI, Ji W, Zhang H, Rabson AB, Shi Y. "Species variation in the mechanisms of mesenchymal stem cell-mediated immunosuppression." Stem Cells. 2009 Aug;27(8):1954-62. PMID: 19544427 , Aug-2009
Articles in Peer-reviewed Journals	Ren G, Su J, Zhao X, Zhang L, Zhang J, Roberts AI, Zhang H, Das G, Shi Y. "Apoptotic cells induce immunosuppression through dendritic cells: critical roles of IFN-gamma and nitric oxide." J Immunol. 2008 Sep 1;181(5):3277-84. PMID: 18713999 , Sep-2008
Articles in Peer-reviewed Journals	Wang KX, Shi YF, Ron Y, Kazanecki CC, Denhardt DT. "Plasma osteopontin modulates chronic restraint stress-induced thymus atrophy by regulating stress hormones: inhibition by an anti-osteopontin monoclonal antibody." J Immunol. 2009 Feb 15;182(4):2485-91. PMID: 19201904 , Feb-2009
Articles in Peer-reviewed Journals	Xu G, Zhang Y, Zhang L, Ren G, Shi Y. "Bone marrow stromal cells induce apoptosis of lymphoma cells in the presence of IFNgamma and TNF by producing nitric oxide." Biochem Biophys Res Commun. 2008 Oct 31;375(4):666-70. PMID: 18755151 , Oct-2008
Articles in Peer-reviewed Journals	Xu G, Zhang Y, Zhang L, Roberts AI, Shi Y. "C/EBPbeta mediates synergistic upregulation of gene expression by interferon-gamma and tumor necrosis factor-alpha in bone marrow-derived mesenchymal stem cells." Stem Cells. 2009 Apr;27(4):942-8 PMID: 19353522 , Apr-2009
Articles in Peer-reviewed Journals	Zhang Y, Xu G, Zhang L, Roberts AI, Shi Y. "Th17 cells undergo Fas-mediated activation-induced cell death independent of IFN-gamma." J Immunol. 2008 Jul 1;181(1):190-6. PMID: 18566384 , Jul-2008