Fiscal Year:	FY 2004	**Task Last Updated:**	FY 09/27/2004
PI Name:	Fritzsch, Bernd Ph.D.		
Project Title:	Neurobiology Of Vestibular Development In Mutant Mice		
Division Name:	Space Biology		
Program/Discipline:	SPACE BIOLOGY		
Program/Discipline--Element/Subdiscipline:	SPACE BIOLOGY--Developmental biology		
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Animal Biology: Vertebrate		
Space Biology Cross-Element Discipline:	(1) Developmental Biology (2) Neurobiology		
Space Biology Special Category:	None		
PI Email:	Fritzsch@Creighton.edu	**Fax:**	FY 402-280-5556
PI Organization Type:	UNIVERSITY	**Phone:**	402-280-2915
Organization Name:	Creighton University		
PI Address 1:	2500 California Plaza		
PI Address 2:			
PI Web Page:	http://www2.creighton.edu/		
City:	Omaha	**State:**	NE
Zip Code:	68178	**Congressional District:**	2
Comments:	NOTE (June 2016): Dr. Fritzsch is currently at University of Iowa		
Project Type:	Ground	**Solicitation / Funding Source:**	2001 Space Biology 01-OBPR-06
Start Date:	11/01/2002	**End Date:**	10/31/2005
No. of Post Docs:	1	**No. of PhD Degrees:**	
No. of PhD Candidates:	1	**No. of Master' Degrees:**	
No. of Master's Candidates:	0	**No. of Bachelor's Degrees:**	
No. of Bachelor's Candidates:	1	**Monitoring Center:**	NASA ARC
Contact Monitor:		**Contact Phone:**	
Contact Email:			
Flight Program:			
Flight Assignment:			
Key Personnel Changes/Previous PI:			
COI Name (Institution):	de Caprona, Dominique (Creighton University) Beisel, Kirk (Creighton University) Nichols, Dave (Creighton University)		
Grant/Contract No.:	NAG2-1595		
Performance Goal No.:			
Performance Goal Text:			
Task Description:	To investigate the effect of normal and hypergravity on vestibular development in mice by investigating the effect of absence of otoconia on gravistatic behavior, neuronal connections and sensory epithelia gene expression. Using otoconia mutant mice we can investigate the absence of linear acceleration perception such as gravity on the behavior and connectional development of mice raised in 1g. Such genetically controlled manipulations allow for a more standardized analysis of a greater number of animals. By understanding the effect of absence of linear acceleration perception (gravity) on the functional development of ear input into the behavior and neuronal organization, we will be able to provide a more molecular understanding of the anticipated changes in the absence of gravity.		

Rationale for HRP Directed Research:	
Research Impact/Earth Benefits:	Our results from a study of embryonic rats developing in microgravity as well as several studies using hypergravity exposure suggest that exposure to altered gravity alters the development of gravistatic vestibular connections and causes long lasting behavioral changes. We now propose to examine how vestibular connections are affected by hypergravity and whether there is a critical age during which gravistatic vestibular connections must be exposed to 1g in order to develop normal connections. The connections of specific inner ear receptors to their central nuclei will be analyzed using the lipophilic dye, DiI, to trace fibers in aldehyde fixed tissue and to analyze their connections to vestibular neurons both light and electron microscopically. In a first, descriptive approach, we will examine the normal sequence of development and how it is altered by exposure to hypergravity by exposing developing rat embryos and pups to hypergravity (2g) for variable periods of time before and after birth. Subsequently we will seek a possible critical age during which the formation of abnormal connections between the ear and brain caused by hypergravity can be appropriately reorganized by intermittent exposure to 1g. For this we will expose rat pups to hypergravity with intermittent exposures to 1g to counteract the hypergravity-induced effects. It is hoped that this program will ultimately help to identify the ages during which biologically relevant gravity stimuli are essential for normal vestibular development. Thus it may be possible to minimize long-term effects of microgravity on the development of vestibular connections during long-term flights by using a 1g centrifuge at appropriate intervals.
Task Progress:	Progress on Aim1: Behavioral analysis of vestibular mutant mice. We have finalized data collection and have a paper in press on our behavioral data. This paper shows that mice without otoconia can not only learn behavioral balance tasks but they do with a steeper learning curve than their control littermates owing to the more profound handicap they are starting with. This paper refutes previous claims in the literature that removal of the ear eliminates spatial learning, likely because the sensory projection of the tlt-mutant mice is at least grossly normal whereas it is absent in animals that underwent ear removal. Title: Partial behavioral compensation is revealed in balance tasked mutant mice lacking otoconia. Authors: D. Crapon de Caprona, K.W. Beisel, D.H. Nichols & B.Fritzsch. Brain Research Bulletin, in press. 2004 Abstract: We describe for the first time behavioral tests which show that mammals with congenital absence of otoconia can learn a motor task that normally relies on gravity perception. The mouse mutation tilted (tlt) occurs in the otopetrin 1 gene (Otop1tlt/tlt) and eliminates an essential component necessary for the formation of otoconia. Our data show that even in the absence of otoconia, tlt mutant mice, like normal mice, learn to cross a bar suspended between two boxes and, with practice, improve their speed of crossing. Despite this learned compensatory skills, tlt mutant mice show balance impairments, such as falling from the bar, not observed in wild type or heterozygous Otop1+/tlt littermates. The tlt mutant mice also use their tail as additional support, a behavior that is rarely exhibited in the control littermates. Interestingly, the Otop1+/tlt heterozygous littermates show in many aspects an intermediate phenotype between wild type and tlt mutant mice, suggestive of a gene dosage effect. Overall, these data support the notion that mammals can use other otic and extraotic receptors such as semicircular canals and limb proprioreceptors, respectively, to compensate for the absence of otoconia mediated gravity perception in a balance task. Progress in Aim 2: Development of connections of vestibular end organs to the brainstem and cerebellum. We have shown that the overall projection of tlt mutant mice is at least grossly normal (see the above cited paper) and are currently investigating the details of the saccular projections in neonates compared to the posterior crista. We concentrate on three areas, the projection to the cell group Y (which appears to be even more prominent in tlt mutant mice), the projection to the dorsal part of the medial vestibular nucleus (which appears to be reduced in tlt mutant mice) and the projection the uvuvla in the cerebellum (which appears to be less segregated than in control mice). We are currently analyzing more of these animals with longer diffusion times. To achieve better segregation of fiber staining we developed in collaboration with PTI Research modified lipophilic dyes. A paper describing some aspect of this new technique is currently in press and a second, technical paper is being submitted within the next few weeks. Progress in Aim 3: Alterations in ion channel expression in vestibular organs of mutants. We have generated, in collaboration with Dr. B. Soarez, Iowa City, a microchip that allows easy analysis of differences in expression profiles of a large number of ion channels. We also generated probes for a number of ion channels that can now be used for quantitative assessment of ion channel expression (real time PCR) as well as qualitative assessment (in situ hybridization of utricular sensory epithelia). We have therefore now generated all the necessary tools for a high throughput analysis of vestibular endorgans of tilt and otogelin mutant mice and control littermates in the next two years.
Bibliography Type:	Description: (Last Updated: 03/26/2008)
Articles in Peer-reviewed Journals	Gu, C.; Rodriguez, E.R.; Reimert, D.V.; Shu, T.; Fritzsch, B.; Richards, L.J.; Kolodkin, A.L.; Ginty, D.D. "Neuropilin-1 conveys semaphorin and VEGF signaling during neural and cardiovascular development." Developmental Cell 5(1):45-57, Jul 2003. , Jul-2003
Articles in Peer-reviewed Journals	Fritzsch, B. "The ear of Latimeria chalumnae revisited." Zoology 106(3):243-248, 2003. , Oct-2003
Articles in Peer-reviewed Journals	Fritzsch, B.; Beisel, K.W. "Molecular conservation and novelties in vertebrate ear development." Current Topics in Developmental Biology 57:1-44, Dec 2003. , Dec-2003
Articles in Peer-reviewed Journals	Fritzsch, B. "Molecular developmental neurobiology of formation, guidance and survival of primary vestibular neurons." Advances in Space Research 32(8):1495-1500, Oct 2003. , Oct-2003
Articles in Peer-reviewed Journals	Fritzsch, B.; Tessarollo, L.; Coppola, E.; Reichardt, L.F. "Neurotrophins in the ear: their roles in sensory neuron survival and fiber guidance." Progress in Brain Research 146:265-278, 2004. , Jan-2004
Articles in Peer-reviewed Journals	Tessarollo, L.; Coppola, V.; Fritzsch, B. "NT-3 replacement with brain-derived neurotrophic factor redirects vestibular nerve fibers to the cochlea." Journal of Neuroscience 24(10):2575-2584, Mar 2004. , Mar-2004
Articles in Peer-reviewed Journals	Fritzsch, B.; Beisel, K.W. "Keeping sensory cells and evolving neurons to connect them to the brain: Molecular conservation and novelties in vertebrate ear development." Brain, Behavior and Evolution 64(3):182-197, Sep 2004. , Sep-2004