Fiscal Year:	FY 2021	**Task Last Updated:**	FY 02/16/2021
PI Name:	Hyde, Truell Ph.D.		
Project Title:	PK-4: Self-Ordering of Interacting Complex Plasma Particles in Microgravity		
Division Name:	Physical Sciences		
Program/Discipline:			
Program/Discipline--Element/Subdiscipline:	FUNDAMENTAL PHYSICS--Fundamental physics		
Joint Agency Name:	National Science Foundation	**TechPort:**	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	None		
Space Biology Cross-Element Discipline:	None		
Space Biology Special Category:	None		
PI Email:	Truell_Hyde@baylor.edu	**Fax:**	FY
PI Organization Type:	UNIVERSITY	**Phone:**	254-710-6717
Organization Name:	Baylor University		
PI Address 1:	Center for Astrophysics, Space Physics, and Engineering Research (CASPER)		
PI Address 2:	1 Bear Pl Unit 97283		
PI Web Page:			
City:	Waco	**State:**	TX
Zip Code:	76798-7283	**Congressional District:**	17
Comments:			
Project Type:	Flight	**Solicitation / Funding Source:**	2016 Physical Sciences NNH16ZTT002N: Fundamental Physics – NASA/NSF Partnership on Science of Dusty Plasmas: Utilizing the PK-4 Facility on board the ISS
Start Date:	04/19/2017	**End Date:**	12/31/2023
No. of Post Docs:	0	**No. of PhD Degrees:**	2
No. of PhD Candidates:	4	**No. of Master' Degrees:**	1
No. of Master's Candidates:	1	**No. of Bachelor's Degrees:**	2
No. of Bachelor's Candidates:	6	**Monitoring Center:**	NASA JPL
Contact Monitor:	Callas, John	**Contact Phone:**	
Contact Email:	john.l.callas@jpl.nasa.gov		
Flight Program:	ISS		
Flight Assignment:	ISS NOTE: End date changed to 12/31/2023 per U. Israelsson/JPL (Ed., 1/6/22) NOTE: End date changed to 8/31/2022 per U. Israelsson/JPL (Ed., 8/12/21)		
Key Personnel Changes/Previous PI:	None.		
COI Name (Institution):	Matthews, Lorin Ph.D. (Baylor University) Hartmann, Peter Ph.D. (Baylor University) Petrov, Oleg Ph.D. (Joint Institute for High Temperatures, Russian Academy of Sciences) Rosenberg, Marlene Ph.D. (University of California, San Diego)		
Grant/Contract No.:	JPL 1571701		
Performance Goal No.:			
Performance Goal Text:			

Task Description:

Abstract --
Complex plasma consists of ions, electrons, neutral gas, and solid microparticles referred to as dust. Dust grains immersed in plasma become negatively charged and interact with one another as well as plasma components through electromagnetic interactions. In particular, dust grains experience a plasma drag force, altering the trajectories of ions in the plasma, and participate in collective plasma interactions, such as waves and instabilities. An interesting aspect of complex plasma is its ability to self-organize into dust liquids, 2D and 3D dust crystal lattices, and 1D dust chains, which are aligned with electric fields in the plasma. This feature provides researchers the ability to examine classical collective phenomena on size and time scales allowing for direct observation (at the kinetic level) of fundamental microscopic and macroscopic processes, as the dust can easily be imaged with high-speed cameras.

In many environments, positive ions in the plasma have a directed flow with respect to the negatively charged dust grains. The resulting interaction between the dust and flowing plasma creates an ion wakefield downstream from the dust particles, with the positive space region modifying the interaction between the charged grains and contributing to the observed dynamics and particle structure. Since the ion wakefield is weak, the effects are often masked by gravity for terrestrial experiments. This international research collaboration employs the Plasma Krystall-4 (PK-4) on the International Space Station (ISS) to conduct experiments in microgravity which investigate the underlying physics behind self-ordering of interacting complex plasma dust particles into field-aligned chains.

Scientific Goals / Objectives of the Research --

The goal of the proposed research project is to investigate the underlying physics behind self-ordering of interacting complex plasma dust particles into field-aligned chains in a microgravity environment. This goal will be accomplished by pursuing the following specific objectives: 1. Establishing the complex plasma parameters required for the formation and stability of extended field-aligned dust particle chains, 2. Measuring the grain-grain potential along and perpendicular to the ion flow direction, and 3. Studying interchain interactions and their role in the structural properties of an ordered complex plasma system.

Significance of the Research --

This research will be used to develop comprehensive theories on interparticle potential and grain-plasma interactions which determine the microscopic and macroscopic properties of complex plasma, including external forces acting on the particles, the fundamental interactions between the particles, thermodynamics, and self-organization of the system. The results will directly impact research in astrophysics and planetary science (dust in protoplanetary disks; charged dust on the surface of airless bodies, such as Mars, the moon, and asteroids; planetary ring systems), atmospheric physics (noctilucent clouds), fusion research (dust contamination, operating stability), materials physics (soft matter, metastable states, complex fluids, granular matter), and advanced manufacturing (semiconductor, etching, and deposition systems).

Rationale for HRP Directed Research:

Impact on Principle Discipline –
Experimental (Baylor & Wigner Research Centre for Physics, Budapest) – Over the past four years, the research analog to the PK-4 (ISS), the PK-4 BU, has been constructed, calibrated, and commissioned. The PK-4 BU is currently the only device of this sort in the world which allows researchers to identify new parameter regimes or develop experimental plans which cannot be developed (or tested) on current PK-4 ground units. Specifically, over the past two years (2019 - 2020), the PK-4 BU was calibrated against data collected by both the PK-4 ISS and the PK-4 ground unit at DLR (German Aerospace Center) Munich. Additionally, the PK-4 BU was calibrated against the initial ground test data collected during the construction of the PK-4 ISS (before launch) to verify that both PK-4 BU Langmuir probe data and LIF (logical interface) data were in agreement with the final calibration data produced before the PK-4 ISS was installed on board the ISS. In both cases this data was collected with the help of our collaborators, with the LIF data collected in collaboration with Caltech and the Langmuir probe data collected in calibration with the Institute of Space Systems (IRS) at the University of Stuttgart.

The above has resulted in the PK-4 BU now being employed successfully for identification of experimental parameter spaces for future experiments aboard the space station and analysis of previous PK-4 ISS data showing anomalies. This process culminated in the summer of 2019 with the PK-4 BU being used by researchers from Baylor, Auburn, and DLR to establish specific parameter spaces for two future experimental campaigns on the PK-4 ISS. These campaigns (Campaign 7 and Campaign 9) were accepted for flight and flown on the PK-4 ISS in November of 2019 and February of 2020. The data from Campaign 7 is under analysis with the first of several papers soon to be submitted. Unfortunately, due to the global pandemic, Campaign 9 data has not yet been delivered to the researchers. Hopefully, this will occur before funding runs out in April of 2021.

Numerical (Baylor) – Along with the development of the PK-4 BU, an entirely new code, DRIAD (Dynamic Response of Ions and Dust) has been written to model the ion-dust interactions within the PK-4 BU discharge. This allows both dust charging and dynamics to be self-consistently calculated, including the effects of the ion wakefield on the dust-dust interactions and charging. The code is written in C++ with CUDA to make use of the parallel processing power of Graphical Processing Units (GPUs). The code makes use of the boundary conditions for the plasma provided by a PIC-MCC code (developed by our collaborators at the Wigner Research Centre for Physics in Hungary) used to model the PK-4 DC discharge. DRIAD is a MD simulation of ion and dust dynamics, each resolved on their own timescale. The charging of the dust is calculated self-consistently within the ion wake. The dynamics of the ions include calculations of ion-ion accelerations, ion-dust accelerations, ion-neutral collisions, ion insertions (assuming a shifted Maxwellian velocity distribution incident on the simulation boundary), and ion deletions due to leaving the simulation boundary or impacting the dust particle. Dust dynamics include calculations of charge, ion-dust accelerations, dust-dust accelerations, external electric fields present in the DC plasma, additional confining electric fields, gas drag, and a thermal gas bath.

The code has been tested using boundary conditions provided by the aforementioned PIC-MCC code, showing the effects of the plasma striations (which occur on the microsecond timescale) on the dust dynamics (which occur on the millisecond time scale).

While many features implemented within the code are specific to the PK4 discharge, the code is robust and can be applied to other dusty plasma experiments by changing the boundary conditions governing the ion flow and external

Research Impact/Earth Benefits:

confining potentials. This allows users of the code to explore the parameter space of experimental conditions to determine which are most favorable for a particular dust configuration, such as one-dimensional strings. A second version of the code is under development which takes advantage of capabilities in C++ to allow rapid implementation of new forces, boundary conditions, elements, etc. This version of the code will also be adaptable beyond dusty plasma physics.

Numerical (Institute for Solid State Physics and Optics, Wigner Research Centre for Physics) -- The results of the PIC/MCC simulations of the PK-4 gas discharge predicting the presence of fast ionization waves during both stationary DC discharge and during polarity switching have now been experimentally verified at Baylor employing the PK-4 (BU). These rapid features remained undetected in the PK-4 (ISS) experiment due to lack of fast diagnostics but are apparently of key importance for the formation of the observed dust particle chains. (See Hartmann, P., M. Rosenberg, Z. Juhasz, L. Matthews, D. Sanford, K. Vermillion, J. Reyes, T. Hyde, Ionization waves in the PK-4 direct current neon discharge, Plasma Sources Science and Technology, 29(11), 115014, November 2020.) These findings provide the necessary foundation for design of future experimental campaigns.

Impact on Scientific Goals / Objectives of the Research

Objective 1: Establishing the plasma parameters required for the formation and stability of dust particle chains.

Experimentally, the PK-4 BU has now been employed successfully for identification of experimental parameter spaces for both current and future experiments aboard the space station and analysis of previous PK-4 ISS data showing anomalies. This process culminated in the summer of 2019 with the PK-4 BU being used by researchers from Baylor, Auburn, and DLR to establish specific parameter spaces for two experimental campaigns on the PK-4 ISS. These campaigns (Campaign 7 and Campaign 9) were accepted for flight and flown on the PK-4 ISS in November of 2019 and February of 2020.

Numerically, the MD model has now been used to show that there is a difference in dust behavior when time-averaged values of the plasma parameters (i.e., electric field, ion and electron density and temperature) are used, compared to the evolving plasma parameters, even when these occur on a microsecond time scale. An important part of this is the extinction/reignition of the plasma during the DC polarity switching of the PK4 electrodes. The average ion concentration around the dust particles is found to be much stronger resulting in ‘deeper wakes’ in the electric potential when the time-evolution of the plasma conditions is resolved.

Objective 2: Measuring the grain-grain potential along and perpendicular to the ion flow direction.

Experimentally / Numerically, the grain-grain potential along and perpendicular to the ion flow direction has been shown to differ significantly from previous analytical models used to explain the formation of dust chains.

Objective 3: Studying interchain interactions and their role in the structural properties of an ordered complex plasma system.

Experimentally, the PK-4 BU has now been employed to identify and reproduce multiple parameter sets favorable to dust string formation and modification of the internal structural properties of these systems. Numerically, preliminary models have been run using up to 1000 dust grains to examine the interaction of multiple chains. One interesting discovery was that particles form nested cylinders arranged in a hexagonal mesh wrapped around each cylindrical surface. Axial cuts through these cylinders reveal ordered linear structures, or chains.

Significant Results

Using data collected on the PK4-BU and time-resolved boundary conditions provided by the PIC-MCC model, the DRIAD code has shown that the ionization waves in the DC column lead to large variations in particle charge on the microsecond time scale. The PIC-MCC model also reveals that the plasma is completed extinguished and reignites over a time span of approximately 200 μs each DC switching cycle. This causes the dust to discharge and then recharge over the same time span. These large variations in the dust charge are most likely the main contributors to the anomalous heating observed for dust particles in experiments.

Impact on Other Disciplines

A second version of the code, RAPID (Rapid Accelerates Physics sImulation Development) is being developed which takes advantage of capabilities in C++ to allow rapid implementation of new forces, boundary conditions, elements, etc. This version of the code will be adaptable beyond dusty plasma physics and be broadly applicable to many types of problems in physics.

Since the field of complex plasmas is now considered the plasma state of Soft Matter, these results will also directly impact research in this field.

Impact on the Development of Human Resources

Baylor –Over the past year, three undergraduate students, five graduate students, and one Assistant Research Professor were directly involved in the experimental portion and five undergraduate and three graduate students were directly involved in the numerical portion of the PK-4 project at Baylor. Four of the five undergraduate students mentioned are members of under-represented minorities in physics and three of the graduate students are female and now using the code in their graduate research. As part of this project, each of these students gained experience in either experimental or numerical skills as well as practice in analyzing data, writing, and presenting papers.

The experimental students also learned required safety skills (i.e., laser safety, basic operating skills in HV, lasers, vacuum systems, and diagnostics) while the numerical students gained experience in testing and developing code, learned how to program using CUDA, and sharpened their skills in C++ and Matlab. (Each of these programming languages is listed among the top twenty programming languages for 2017 by Institute of Electrical and Electronics Engineers (IEEE).) In addition, all of the students listed are also gaining experience in both technical writing and speaking.

Impact on Professional Development –

CASPER (Baylor) – Over the past year, three undergraduate students and five graduate students were directly involved in the experimental portion and five undergraduate and three graduate students were directly involved in the numerical

portion of the PK-4 project at Baylor. Experimental students learned required safety skills (i.e., laser safety, basic operating skills in HV, lasers, vacuum systems, and diagnostics) while numerical students gained experience in testing and developing code, learned how to program using CUDA, and sharpened their skills in C++ and Matlab. One undergraduate student (Dustin Sanford) and three graduate students (Katrina Vermillion, Sina Rostami, Khandaker Sharmin Ashrafi) were involved in the development and testing of the computer code over the past year. The students learned how to program using CUDA in addition to sharpening their skills programming in C++ and Matlab. In addition, three other undergraduate students (Martin Lechuga, Emerson Gehr, and Evelyn Guay) learned to use ImageJ and Matlab to analyze data from the PK4 experiments on the Space Station as well as numerical data proved by DRIAD.

Undergraduates Emerson Gehr and Evelyn Guay presented their work at the 62nd Annual Meeting of the American Physical Society-Division of Plasma Physics (APS-DPP) (virtual), October, 2020. The students also presented their results as part of the Baylor Physics Department colloquium series in September and December 2020, and will make poster presentations during Universities Space Research Association (USRA) Scholars Week, March 2021.

Graduate students Katrina Vermillion and Zhiyue Ding made presentations on their work at the 62nd Annual Meeting of the APS-DPP (virtual), October, 2020. Ms. Vermillion also presented at the APS Texas Section Meeting, October 2020.

Graduate student Zhiyue Ding defended his dissertation on Nonlinear Responses in Dusty Plasma and received the Ph.D. degree in December, 2020. Graduate student Khandaker Sharmin Ashrafi defended her dissertation on numerical modeling of dust and ions and received the Ph.D. degree in December, 2020.

All of the students listed gained experience in both technical writing and speaking with their research presented at the Annual Meetings of the APS-DPP. Both Mr. Ding and Ms. Ashrafi presented their results as part of the Virtual International Conference IOSDP during the fall of 2020. Additionally, undergraduate and graduate students presented their results as part of the Baylor Physics Department colloquium series through poster presentations during USRA Scholars Week.

Additionally, over the past year CASPER students (both graduate and undergraduate) have had the opportunity to attend virtual meetings with the PI and our Co-Is from all over the world (Peter Hartmann, Hubertus Thomas, Vladimir Nosenko, Marlene Rosenberg).

Impact on Physical / Numerical Resources –

Experimental (CASPER, Baylor & Institute for Solid State Physics and Optics, Wigner Research Centre for Physics) --

The PK-4 BU is a new tool that continues to prove itself useful for the design and testing of potential experiments employing the PK-4 ISS. The PK-4 BU is now being used by multiple US research groups (including graduate students) and should prove useful for years to come. Motivated by new insights provided by both numerical simulation of DC gas discharge plasma (Wigner Research Centre for Physics in Budapest, Hungary) and the PK-4 BU device (CASPER HIDPL laboratory at Baylor University) experimental verification of fundamental calibration data has now been completed. This has proven invaluable to the identification, design, and planning of future experiments on the PK-4 facility ISS with two experimental campaigns (Campaign 7 and Campaign 9) accepted for 2019 and 2020.

Numerical (Baylor) -- The IonWake code and the DRIAD (Dynamic Response of Ions and Dust) code are both new tools which will prove useful for many years to come. In the PI's experience, computer codes which provide new capabilities are used for ten to twenty years (or more) and provide a research platform for many future graduate and undergraduate students. As an example, previously developed codes have been used by about 20 undergraduate students and 3-5 graduate students over a period of ten years.

Impact on Society –

The impact of this project on graduate students, undergraduate students and K-12 students continues to impact the overall science literacy of the voting public, making an impact beyond the bounds of science and engineering alone.

Collaborator Meetings –

APS DPP / IEEE PPPS - CASPER Personnel (PI, Co-I, Research Faculty, Graduate Students) with Ed Thomas, Jeremiah Williams, Paul Bellan, Peter Hartmann & Marlene Rosenberg (Virtual). CASPER Personnel (PI) with DLR Personnel (All) (Virtual)

New Technology: Installation and alignment of additional video equipment; LIF assessment of plasma across multiple regimes; Langmuir probes developed and installed with data compared to original source data.

Experimental (Baylor & Wigner Research Centre for Physics, Budapest) – Over the past four years, the research analog to the PK-4 (ISS), the PK-4 BU, has been constructed, calibrated, and commissioned. The PK-4 BU is currently the only device of this sort in the world that allows researchers to identify new parameter regimes or develop experimental plans which cannot be developed (or tested) on current PK-4 ground units. Specifically, over the past two years (2019 - 2020), the PK-4 BU was calibrated against data collected by both the PK-4 ISS and the PK-4 ground unit at DLR Munich. Additionally, the PK-4 BU was calibrated against initial ground test data collected during the construction of the PK-4 ISS (before launch) to verify that both PK-4 BU Langmuir probe data and LIF data were in agreement with the final calibration data produced before the PK-4 ISS was installed on board the ISS. In both cases this data was collected with the help of our collaborators, with the LIF data collected in collaboration with Caltech and the Langmuir probe data collected in calibration with the IRS at the University of Stuttgart.
The above has resulted in the PK-4 BU being employed successfully for identification of experimental parameter spaces for PK-4 ISS campaigns and analysis of previous PK-4 ISS data showing anomalies. This process culminated in the summers of 2019 and 2020 with the PK-4 BU being used by researchers from Baylor, Auburn, and DLR to establish specific parameter spaces for two experimental campaigns on the PK-4 ISS. These campaigns (Campaign 7 and Campaign 9) were accepted for flight and flown on the PK-4 ISS in November of 2019 and February of 2020. The data from Campaign 7 is under analysis with the first of several papers soon to be submitted. Unfortunately, due to the global pandemic, Campaign 9 data has not yet been delivered to the researchers. Hopefully, this will occur before funding runs out in May of 2021.

Numerical (Baylor) – Along with the development of the PK-4 BU, an entirely new code, DRIAD (Dynamic Response

of Ions and Dust) was written to model the ion-dust interactions within the PK-4 BU discharge. This allows both dust charging and dynamics to be self-consistently calculated, including the effects of the ion wakefield on the dust-dust interactions and charging. The code is written in C++ with CUDA to make use of the parallel processing power of Graphical Processing Units (GPUs). The code makes use of the boundary conditions for the plasma provided by a PIC-MCC code (developed by our collaborators at the Wigner Research Centre for Physics in Hungary) used to model the PK-4 DC discharge. DRIAD is a MD simulation of ion and dust dynamics, each resolved on their own timescale. The charging of the dust is calculated self-consistently within the ion wake. The dynamics of the ion includes calculations of ion-ion accelerations, ion-dust accelerations, ion-neutral collisions, ions insertions (assuming a shifted Maxwellian velocity distribution incident on the simulation boundary), and ion deletions due to leaving the simulation boundary or impacting the dust particle. The dust dynamics includes calculations of charge, ion-dust accelerations, dust-dust accelerations, external electric fields present in the DC plasma, additional confining electric fields, gas drag, and a thermal gas bath.

The code has been tested using boundary conditions provided by the aforementioned PIC-MCC code, showing the effects of the plasma striations (which occur on the microsecond timescale) on the dust dynamics (which occur on the millisecond time scale).

While many features implemented within the code are specific to the PK4 discharge, the code is robust and can be applied to other dusty plasma experiments by changing the boundary conditions governing the ion flow and external confining potentials. This allows users of the code to explore the parameter space of experimental conditions to determine which are most favorable for a particular dust configuration, such as one-dimensional strings. A second version of the code is being developed which takes advantage of capabilities in C++ to allow rapid implementation of new forces, boundary conditions, elements, etc. This version of the code will also be adaptable beyond dusty plasma physics

Numerical (Institute for Solid State Physics and Optics, Wigner Research Centre for Physics) -- The results of the PIC/MCC simulations of the PK-4 gas discharge predicting the presence of fast ionization waves during both stationary DC discharge and during polarity switching have now been experimentally verified employing the PK-4 (BU). These rapid features remained undetected in the PK-4 (ISS) experiment due to lack of fast diagnostics but are apparently of key importance for the formation of the observed dust particle chains. These findings should provide the necessary foundation for design of future experimental campaigns.

Task Progress:

Theoretical (University of California, San Diego) – M. Rosenberg, Major Activities

A. PK-4 discharge modeling. The UCSD-University of California, San Diego PI collaborated on a theoretical study of the PK-4 cylindrical discharge. The study was geared toward determining plasma parameters to be used for studies of dust dynamics in this device. The numerical investigation used a two-dimensional, electro- static, particle-in-cell with Monte Carlo collisions discharge simulation code. The simulations showed that the positive column is inhomogeneous on short timescales with rapidly moving, high amplitude ionization waves dominating the structure. As a result, the local plasma parameters fluctuate at frequencies of the order of 10 KHz and with amplitudes up to 10 times the time-averaged values. The oscillatory behavior of the plasma environment is expected to affect the charging and chain formation of dust particles in PK-4 experiments. For example, the ion drift speed is a crucial parameter in the electrorheological model of chain formation, and the electric field of the ionization waves can significantly affect the ion drift speed. A paper was published: P. Hartmann, M. Rosenberg, Z. Juhasz, L. S. Matthews, D. L. Sanford, K. Vermillion, J. Carmona-Reyes and T. W. Hyde, 2020, Ionization waves in the PK-4 direct current neon discharge, Plasma Sources Sci. Technol. 29, 115014.

B. Chain formation in PK-4. The UCSD PI has been collaborating with the Baylor and WRC groups on studies of how fluctuations in local plasma parameters due to ionization waves in PK-4 can affect the formation of chains of dust particles. Papers are in preparation.

C. Waves in electrorheological dusty plasmas. The UCSD PI worked on theoretical studies of waves in a one-dimensional dust chain in an electrorheological dusty plasma, where the dust particles interact via combined Yukawa and dipole-dipole potentials. The configuration of the dust particles is given by a pair correlation function, and the chain is confined in the perpendicular direction by a parabolic potential. We are using pair correlation functions and confinement potentials provided by simulations of chain formation by the Baylor group. Both longitudinal and transverse waves are being considered, and dust-neutral collisions are taken into account. Work is continuing.

D. Ion-dust streaming instability. Dust acoustic, or dust density, waves can be excited by the flow of ions via an ion-dust streaming instability. Dust acoustic, or dust density, waves have been observed in numerous ground-based experiments, and also in PK-4 experiments (e.g., Jaiswal et al., 2018, Phys. Plasmas 25, 083705; Goree et al., 2020, Phys. Plasmas 27, 123701). The UCSD PI, together with colleagues at UCSD, began to study the nonlinear development of the ion-dust streaming instability in parameter regimes where the ion and dust collision frequencies are a significant fraction of their respective plasma frequencies. This regime may be relevant to ground-based or microgravity experiments with high neutral pressure. The study used a one-dimensional particle-in-cell electrostatic simulation code that included thermalizing collisions with the background neutrals. Using a particular set of nominal parameters with an ion thermal Mach speed of 3, the simulations found cnodial-like dust density waveforms (i.e, waveforms with broad troughs and sharp crests) in the nonlinear phase. The study may have relevance for high pressure environments of PK-4, but further work is needed to model the ion-dust instability using representative PK-4 parameters.

A paper was published: K. Quest, M. Rosenberg and A. Levine, Simulations of ion- dust streaming instability in a highly collisional plasma, 2020, J. Plasma Phys. 86, 905860605.

E. Possible use of Mie-resonant microparticles. Recently there has been interest in the use of dielectric particles with high refractive index n in novel optical and photonic devices (e.g., Yang et al. 2017, Phys. Reports 701, 1-50). The UCSD PI considered theoretically the scattering of far-infrared radiation in a dusty plasma composed of spherical, high-n dielectric particles of radii ~ 4 - 5 μm. In particular, the scattering of radiation in the 30 - 40 μm range from aluminum oxide or magnesium oxide particles was studied. Using Mie theory, it was found that a large scattering efficiency can occur at magnetic dipole resonance wavelengths, and there can be directional scattering of electromagnetic waves (e.g., for a fixed particle size, the scattering can be predominantly backward in certain wavelength intervals). Possible applications were considered, including partial mirror behavior, enhanced extinction, and particle size analysis. Possible complications were also considered, including issues related to particle heating which leads to constraints on the background gas pressure as well as the intensity of the incident radiation.

	A paper is in preparation: M. Rosenberg, “On scattering in a dusty plasma composed of Mie-resonant dielectric particles.”
Bibliography Type:	Description: (Last Updated: 08/08/2024)
Articles in Other Journals or Periodicals	Qiao K, Ding Z, Carmona J, Matthews LS, Hyde TW. "Ion wake induced mode coupling in a horizontal chain in complex plasmas." Submitted to Physical Review E, August, 2020. bioRxiv preprint server: https://arxiv.org/abs/2008.01165 , Aug-2020
Articles in Peer-reviewed Journals	Quest K, Rosenberg M, Levine A. "Simulations of ion-dust streaming instability in a highly collisional plasma." Journal of Plasma Physics. 2020 Dec;86(6):905860605. https://doi.org/10.1017/S002237782000135X , Dec-2020
Articles in Peer-reviewed Journals	Hartmann P, Rosenberg M, Juhasz Z, Matthews LS, Sanford D, Vermillion K, Reyes JC, Hyde TW. "Ionization waves in the PK-4 direct current neon discharge." Plasma Sources Science and Technology. 2020 Nov;29:115014. https://doi.org/10.1088/1361-6595/abb955 , Nov-2020
Articles in Peer-reviewed Journals	Kostadinova EG, Padgett J, Liaw C, Matthews LS, Hyde TW. "Numerical study of anomalous diffusion of light in semicrystalline polymer structures." Physical Review Research. 2020 Dec;2(4):043375. https://doi.org/10.1103/PhysRevResearch.2.043375 , Dec-2020
Articles in Peer-reviewed Journals	Sharmin Ashrafi K, Yousefi R, Chen M, Matthews LS, Hyde TW. "Dust as probes: determining confinement and interaction forces." Physical Review E. 2020 Oct;102(4):043210. https://doi.org/10.1103/PhysRevE.102.043210 , Oct-2020
Articles in Peer-reviewed Journals	Xiang C, Carbadillo A, Matthews LS, Hyde TW. "The initial structure of chondrule dust II: charged grains." Icarus. 2021 Jan 15;354:114053. https://doi.org/10.1016/j.icarus.2020.114053 , Jan-2021
Articles in Peer-reviewed Journals	Schmidt J, Hyde TW. "Discharge parameters of PlasmaKristall-4BU: A modifiable dusty plasma experiment." Rev Sci Instrum. 2020 Aug 1;91(8):083506. https://doi.org/10.1063/5.0005325 ; PMID: 32872968 , Aug-2020
Articles in Peer-reviewed Journals	Lisin EA, Petrov OF, Sametov EA, Vaulina OS, Statsenko KB, Vasiliev MM, Carmona-Reyes J, Hyde TW. "Experimental study of the nonreciprocal effective interactions between microparticles in an anisotropic plasma." Sci Rep. 2020 Aug 12;10(1):13653. https://doi.org/10.1038/s41598-020-70441-z ; PMID: 32788600; PMCID: PMC7423964 , Aug-2020
Articles in Peer-reviewed Journals	Xiang C, Carbadillo A, Matthews LS, Hyde TW. "Detailed model of the growth of fluffy dust aggregates in a protoplanetary disk: effects of nebular conditions." Astrophysical Journal. 2020 Jul 20;897(2):182. https://doi.org/10.3847/1538-4357/ab96c2 , Jul-2020
Articles in Peer-reviewed Journals	Padgett JL, Kostadinova EG, Liaw CD, Busse K, Matthews LS, Hyde TW. "Anomalous diffusion in one-dimensional disordered systems: a discrete fractional Laplacian method." Journal of Physics A: Mathematical and Theoretical. 2020 Apr 3;53(13):135205. https://doi.org/10.1088/1751-8121/ab7499 , Apr-2020
Articles in Peer-reviewed Journals	Matthews LS, Sanford D, Kostadinova EG, Ashrafi K, Guay E, Hyde TW. "Dust charging in dynamic ion wakes." Physics of Plasmas. 2020 Feb;27(2):023703. https://doi.org/10.1063/1.5124246 , Feb-2020
Articles in Peer-reviewed Journals	Schmidt J, Laufer R, Hyde TW, Herdrich G. "Operational behaviour of the inductively-heated plasma generator IPG6-B for scientific applications." Vacuum. 2020 Jun;176:109338. https://doi.org/10.1016/j.vacuum.2020.109338 , Jun-2020
Dissertations and Theses	Ding Z. "Nonlinear Responses in Dusty Plasma." Dissertation, Baylor University, December 2020. , Dec-2020
Papers from Meeting Proceedings	Schmidt J, Carballido A, Laufer R, Herdrich G, Hyde TW. "Behavior of Dust in an Inductively-Heated Plasma Jet with Application to Planetary Science." Presented at 51st Lunar and Planetary Science Conference, The Woodlands, Texas, March 16-20, 2020. Proceedings of the Lunar Planetary Science Conference, 2020. 51st Lunar and Planetary Science Conference, March, 2020 at The Woodlands, Texas. LPI Contribution No. 2326, id.1839. , Mar-2020