Fiscal Year:	FY 2005	**Task Last Updated:**	FY 12/16/2004
PI Name:	Bikle, Daniel David M.D., Ph.D.		
Project Title:	The Effect of Skeletal Unloading on Bone Formation		
Division Name:	Space Biology		
Program/Discipline:	SPACE BIOLOGY		
Program/Discipline--Element/Subdiscipline:	SPACE BIOLOGY--Organismal biology		
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Cell & Molecular Biology (2) Animal Biology: Vertebrate		
Space Biology Cross-Element Discipline:	(1) Developmental Biology (2) Musculoskeletal Biology		
Space Biology Special Category:	(1) Cell Culture		
PI Email:	doctor@itsa.ucsf.edu	**Fax:**	FY 415-750-6929
PI Organization Type:	GOVERNMENT	**Phone:**	415-221-4810 3338
Organization Name:	Veterans Affairs Medical Center - University of California		
PI Address 1:	4150 Clement Street (111N)		
PI Address 2:			
PI Web Page:			
City:	San Francisco	**State:**	CA
Zip Code:	94121	**Congressional District:**	8
Comments:			
Project Type:	Ground	**Solicitation / Funding Source:**	2003 Space Biology 03-OBPR-03
Start Date:	02/01/2004	**End Date:**	09/30/2007
No. of Post Docs:	1	**No. of PhD Degrees:**	
No. of PhD Candidates:	0	**No. of Master' Degrees:**	
No. of Master's Candidates:	0	**No. of Bachelor's Degrees:**	
No. of Bachelor's Candidates:	0	**Monitoring Center:**	NASA ARC
Contact Monitor:		**Contact Phone:**	
Contact Email:			
Flight Program:			
Flight Assignment:	NCE to 9/30/2007, per A. Chu 7/2006.		
Key Personnel Changes/Previous PI:	n/a		
COI Name (Institution):			
Grant/Contract No.:	NNA04CC67G		
Performance Goal No.:			
Performance Goal Text:			
	Skeletal unloading results in an inhibition of bone formation associated with a decrease in osteoblast number, impaired mineralization of bone, and altered proliferation and differentiation of osteoprogenitor cells. Although such changes are likely to be mediated by multiple factors, resistance to the skeletal growth promoting action of insulin like growth factor-I (IGF-I) appears to play an important role. IGF-I mRNA and protein levels rise during skeletal unloading, but the ability of IGF-I to stimulate proliferation of bone cells in vivo and in vitro when taken from unloaded bone is markedly impaired. This resistance to IGF-I is associated with a failure of IGF-I to activate its receptor and the subsequent IGF-I signaling pathways. BMSC from unloaded bone show a reduction in integrin expression, and inhibition of integrin signaling reproduces the findings of IGF-I resistance. Therefore, we propose to test the following hypothesis: Skeletal unloading inhibits the ability of IGF-I to stimulate osteoprogenitor cell proliferation by impairing, via an		

Task Description:	integrin-mediated mechanism, IGF-I signal transduction. This hypothesis will be tested by achieving three specific aims: 1) Determine the points at which skeletal unloading blocks IGF-I stimulated bone formation. This will be accomplished by determining whether unloading alters stem cell commitment to the osteoblast lineage, IGF-I receptor activation, and/or the mRNA and protein levels and/or activation of the components of the IGF-I signaling pathway(s). 2) Determine the ability of skeletal unloading to block the response of osteoprogenitor cells to other growth factors such as EGF, PDGF, and FGF with respect to proliferation and activation of the MAPK and/or PI3K pathway. Those growth factors found to affected similarly to IGF-I by skeletal unloading will be further explored with respect to mechanism. 3) Determine the interaction between IGF-I and integrin signaling in the regulation of osteoprogenitor cell proliferation. This will be accomplished by determining the effect of skeletal unloading on the mRNA and protein levels of the various integrins in osteoprogenitor cells, determining which integrins potentiate IGF-I responsiveness, whether the effect of integrins on IGF-I signaling is direct or indirect, and whether correction of integrin levels, if altered by skeletal unloading, will restore IGF-I responsiveness. We anticipate that these studies will increase our understanding of the mechanism(s) for bone loss in weightlessness, and lead to measures that will prevent or reverse such losses. Specific Aims: 1. Determine the point(s) at which skeletal unloading blocks IGF-I stimulated bone formation. a. Determine whether skeletal unloading alters the mesenchymal pool committed to osteoblast differentiation. b. Determine whether skeletal unloading alters the ability of IGF-I to bind to and activate the IGF-I receptor (IGF-IR) in those cells committed to osteoblast differentiation c. Determine the relative contributions of the ras/MAPK and PI3K/Akt pathways to the proliferative response to IGF-I. d. Determine whether skeletal unloading alters the protein levels and/or activation of the components of the pathway(s) mediating the proliferative response to IGF-I. e. Determine whether skeletal unloading alters the nuclear translocation of ERK, PI3K, and Akt and cellular levels of cyclin D1, p21cip, and p27kip 2. Determine the ability of skeletal unloading to block the response of BMSC to other growth factors such as EGF, PDGF and FGF with respect to: a. Proliferation b. Growth factor receptor number, binding, and activation c. Activation of the ras/MAPK and/or PI3K/Akt pathways 3. Determine the interaction between IGF-I and integrin signaling as affected by skeletal unloading in the regulation of osteoprogenitor cell proliferation. a. Determine the mRNA and protein levels of the various integrins in osteoprogenitor cells from normally loaded and unloaded bone b. Determine which integrins potentiate IGF-I responsiveness with respect to proliferation and activation of the IGF-IR and the ras/MAPK and PI3K/Akt pathways c. Determine whether the effect of integrins on IGF-IR activation requires direct interaction with the IGF-IR d. Determine whether correction of integrin levels, if altered by skeletal unloading, will restore IGF-I responsiveness with respect to proliferation and activation of IGF-I signaling pathways. The bone loss associated with skeletal unloading is of great importance to individuals subjected to prolonged periods of skeletal unloading whether this be in microgravity, in a cast, or following neural injury or stroke. Determining the mechanism by which such bone loss occurs is the first step toward effective prevention and treatment. Our project is directed at understanding this mechanism.
Rationale for HRP Directed Research:	
Research Impact/Earth Benefits:	This proposal seeks to understand the mechanism(s) for bone loss during skeletal unloading. This is a major problem clinically for immobilized and aging patients. Bone is a modeling and remodeling tissue meaning that it is undergoing continual development. We expect such research to enable us to understand whether, and how, the development of cells and systems (in this case bone) depend on mechanical forces, the role of cellular structures (bone cells within their own matrix) in response to mechanical forces, how the bone and bone cells sense and use such forces, and how physical and chemical signals interact to determine biological structure and function (bone shape and strength). Such research will contribute to our knowledge of how osteoporosis develops whether it be by aging, by immobilization, or by drugs. As our understanding of the processes by which bone maintains its integrity is increased, we anticipate that our ability to develop preventive measures and treatment for osteoporosis will improve.
Task Progress:	We have made good progress toward achieving these aims of the task. 1. Determine the points at which skeletal unloading blocks IGF-I stimulation bone formation. In this aim we have demonstrated that skeletal unloading does not alter the binding of IGF-I to its receptor. We have used an analog of IGF-I, namely des IGF-I, which does not bind to IGF-bind proteins and demonstrated that like IGF-I itself binding to the bone marrow stromal cells (BMSC) from loaded and unloaded bones is equivalent. We have shown that activation of the IGF-I receptor via phosphorylation is the critical step that is impaired by skeletal unloading. The down stream events such as activation of the MAPK pathway and the subsequent proliferative response to IGF-I are impaired in BMSC from unloaded bones, but this follows the failure to activate the IGF-I receptor. 2. Determine the ability of skeletal unloading to block the response of BMSC to other growth factors. We have examined in depth the effect of skeletal unloading on the resistance to PDGF. To our surprise, BMSC from unloaded bones respond quite well to PDGF with respect to activation of the PDGF receptor (via phosphorylation) and proliferation. Furthermore, the integrin antagonist echistatin, which blocks the response of BMSC to IGF-I, fails to block the response of BMSC to PDGF. Thus, the resistance to IGF-I caused by skeletal unloading appears to be specific to IGF-I and is not generalized to all other growth factors. 3. Determine the interaction between IGF-I and integrin signaling as affected by skeletal unloading. We have found that the expression of a number of integrins is reduced in the BMSC from unloaded bones. We are having some difficulty showing that the protein levels are reduced because of the lack of good antibodies to rat integrins. We are currently developing the methods to selectively knockout specific integrins to determine which if any are critical for BMSC response to IGF-I. These experiments are being performed in both rat BMSC and a human osteoblast cell line for which better integrin antibodies are available. Using this latter cell line, we have had preliminary success in showing that at least one of the integrins of interest binds directly to the IGF-I receptor, as shown by co-immunoprecipitation.
Bibliography Type:	Description: (Last Updated: 04/15/2008)
Articles in Peer-reviewed Journals	Bikle DD, Sakata T, Halloran BP "The Impact of Skeletal Unloading on Bone Formation. " Grav Space Biol Bull 16:45-54 , Jun-2003
Articles in Peer-reviewed Journals	Nauman EA, Sakata T, Keaveny TM, Halloran BP, Bikle DD "bFGF administration lowers the phosphate threshold for mineralization in bone marrow stromal cells." Calc Tiss Internl 73:147-152 , May-2003
Articles in Peer-reviewed Journals	Sakata T, Halloran BP, Elalieh HZ, Munson SJ, Rudner L, Venton L, Ginzinger D, Rosen CJ, Bikle DD "Skeletal unloading induces resistance to insulin-like growth factor-I on bone formation." Bone 32:669-680 , Oct-2003

Articles in Peer-reviewed Journals	Sakata T, Wang Y, Halloran BP, ElAlieh HZ, Cao J, Bikle DD. "Skeletal unloading induces resistance to insulin-like growth factor-I (IGF-I) by inhibiting activation of the IGF-I signaling pathways." J Bone Miner Res. 2004 Mar;19(3):436-46. PMID: 15040832 , Mar-2004
Articles in Peer-reviewed Journals	Bikle DD "Impact of Skeletal Unloading on Bone Formation: New Understanding of Molecular Mechanisms. " J Ortho Sci , Dec-2004