| | | | |
|---|---|---|---|
| **Fiscal Year:** | FY 2018 | **Task Last Updated:** | FY 05/07/2018 |
| **PI Name:** | Brandizzi, Federica Ph.D. | | |
| **Project Title:** | Mechanisms for Plant Adaptation to Space Environment | | |
| **Division Name:** | Space Biology | | |
| **Program/Discipline:** | SPACE BIOLOGY | | |
| **Program/Discipline--Element/Subdiscipline:** | SPACE BIOLOGY--Cellular and molecular biology | | |
| **Joint Agency Name:** | | **TechPort:** | No |
| **Human Research Program Elements:** | None | | |
| **Human Research Program Risks:** | None | | |
| **Space Biology Element:** | (1) Cell & Molecular Biology<br>(2) Plant Biology | | |
| **Space Biology Cross-Element Discipline:** | None | | |
| **Space Biology Special Category:** | (1) Bioregenerative Life Support | | |
| **PI Email:** | brandizz@msu.edu | **Fax:** | FY |
| **PI Organization Type:** | UNIVERSITY | **Phone:** | 517-353-7872 |
| **Organization Name:** | Michigan State University | | |
| **PI Address 1:** | 106 Plant Biology | | |
| **PI Address 2:** | MSU-DOE Plant Research Laboratory | | |
| **PI Web Page:** | | | |
| **City:** | East Lansing | **State:** | MI |
| **Zip Code:** | 48824-1046 | **Congressional District:** | 8 |
| **Comments:** | | | |
| **Project Type:** | Flight | **Solicitation / Funding Source:** | 2011 Space Biology NNH11ZTT002N |
| **Start Date:** | 07/27/2012 | **End Date:** | 12/31/2019 |
| **No. of Post Docs:** | 0 | **No. of PhD Degrees:** | 0 |
| **No. of PhD Candidates:** | 1 | **No. of Master' Degrees:** | 0 |
| **No. of Master's Candidates:** | 0 | **No. of Bachelor's Degrees:** | 0 |
| **No. of Bachelor's Candidates:** | 0 | **Monitoring Center:** | NASA KSC |
| **Contact Monitor:** | Levine, Howard | **Contact Phone:** | 321-861-3502 |
| **Contact Email:** | howard.g.levine@nasa.gov | | |
| **Flight Program:** | ISS | | |
| **Flight Assignment:** | ISS<br>NOTE: Extended to 12/31/2019 per NSSC information (Ed., 12/28/18)<br>NOTE: Extended to 12/31/2018 per PI (Ed., 12/10/15)<br>NOTE: Extended to 12/31/2015 per NSSC information (Ed., 12/16/14)<br>NOTE: Extended to 12/31/2014 per NSSC information; original end date was 1/31/13 (Ed., 1/14/14)<br>NOTE: Extended to 12/31/2013 per NSSC information; original end date was 1/31/13 (Ed., 1/25/13) | | |
| **Key Personnel Changes/Previous PI:** | May 2018 report: No changes. | | |
| **COI Name (Institution):** | | | |
| **Grant/Contract No.:** | NNX12AN71G | | |
| **Performance Goal No.:** | | | |
| **Performance Goal Text:** | | | |

**Task Description:**

Successful plant growth in closed-loop life support conditions is a difficult challenge for the realization of long-term habitation of spacecraft and other extraterrestrial environments. In such environments, plants can undergo stress induced by a number of factors including changes in gravity, radiations, vibration, limited exchange of gases, and suboptimal growth conditions (temperature, light, nutrients). These sources of stress are often associated with reprogramming of gene expression and can cause limited plant growth, development, and yield. To facilitate plant life in space, it is crucial to acquire a better understanding of the genetic changes that enable plant cells to respond to spaceflight stress. To do so, one goal of this proposal is to define the underlying mechanisms of plant adaptation to the spaceflight environment at a transcriptional level. Through research supported by NASA, we have successfully identified a protein named AtIRE1 as a master regulator of transcription in conditions of stress responses to abiotic, biotic stress, and to gravity changes in plants. However, the downstream signaling pathways controlled by AtIRE1 are not well defined, especially in conditions of altered gravity. A better understanding of the signaling pathways controlled by AtIRE1 is important to understand how plants can grow in conditions of stress. We propose to use in-flight and ground resources along with genomics and transcriptomics analyses in the model plant Arabidopsis thaliana to understand the regulatory role of AtIRE1 in adaptation to spaceflight stress. Development of our research will continue to contribute to the understanding of basic signaling pathways that are in place to ensure stress survival in hostile environments, thus making possible the design and growth of plants that are resistant to space stress. To contribute further to the successful realization of habitation in space, we also aim to develop plants that can function as bioindicators of stress during in-flight situations. To do so, we will engineer plants with an AtIre1 substrate that is activated specifically in conditions of stress and that we will adapt to function as a visual stress reporter. The availability of real-time stress bioindicators will provide scientists and astronauts with direct read-outs of stress in the space environment. The results gathered in our research will contribute to NASA's strategic plans for the realization of long-term habitation of space and planetary surfaces. Because of the conservation of stress sensing and response mechanisms across multicellular organisms, we expect that our results will also have important implications for the general knowledge of stress and in the design of solutions for space stress management in multicellular organisms, including humans.

**Rationale for HRP Directed Research:**

**Research Impact/Earth Benefits:**

The molecular study of human health conditions that develop in response to long term habitation in spaceflight conditions is inherently limited by the necessity to insure safety of study participants. As such, the cellular and molecular pathology of these disease states and potential remediations are unknown. Comparative biology, or the study of similar situations in different types of organisms, presents a unique opportunity to explore these human conditions with minimal associated risks. The study of conserved cellular stress responsive pathways between humans and plants provides this benefit and many others. In particular, plants are the primary production source for metabolic energy and oxygen in our terrestrial environment, and proper application in extraterrestrial and spaceflight conditions could potentially reduce the cost of recreating human habitable environments in extremely harsh conditions.
Plants, like animals, suffer from stress linked to a variety of challenges. For example, certain adverse environmental conditions, such as changes in temperature, reduced nutrients, and disease, as well as physiological situations, such as growth, are known to generate an imbalance between demand and capacity of protein synthesis in the endoplasmic reticulum (ER), a condition that is also known as ER stress (1, 2). The ER, a conserved structure in human and plant cells, initiates the production of all secretory proteins. Secretory proteins constitute one third of the cellular proteome and allow cells to communicate with the external environment. The ER can sense stress due to accumulation of unfolded proteins in its lumen, and restores homeostasis by invoking a largely conserved cytoprotective signaling pathway, called the unfolded protein response (UPR). Compromised UPR leads to serious conditions and even death in animals and plants (3-5). In spaceflight conditions a number of factors including microgravity, radiations, vibration, limited exchange of gases, and suboptimal growth conditions (temperature, light, nutrients) reduces overall plant growth, impairs development, and reduces yield.

In animals and plants, several sensors and transducers initiate the UPR; however, IRE1, an ER-associated protein kinase and ribonuclease, functions as a major ER stress sensor and plays a key role in the UPR signaling by controlling UPR gene expression (4, 6, 7). Activities in our lab and others have demonstrated that plant IRE1 is needed to respond to a diversified portfolio of stimuli linked to abiotic and biotic stress and cell growth and differentiation, as well as to modulate responsive outputs (4, 5).

In all eukaryotes, IRE1 activates the UPR by an unconventional splicing of a bZIP transcription factor, namely Hac1 in yeast8-10 and Xbp1 in metazoans (11, 12). Recent efforts in our lab and others have shown that in the model species Arabidopsis thaliana, the functional equivalent of Hac1 and Xbp1 is bZIP60 (5, 13, 14). The bZIP60 mRNA splicing is followed by the synthesis of the mature form of bZIP60, which can modulate transcription of UPR genes in the nucleus. Thus, synthesis of the transcriptionally active bZIP60 occurs only in conditions of stress. In our research we will adopt the stress-dependent activation of bZIP60 to generate plants that can function as real-time stress bioreporters that can be easily used to report on the onset of stress in plants.

We believe that we will learn the mechanisms that enable plants to respond to ER stress. This is highly significant for the understanding of basic defense mechanisms in plants, and for engineering plants that can cope with stress on Earth.

We also expect that our research will have an important impact in biology with implications well beyond plants. Because the signaling pathways between plant and mammalian UPR are largely conserved, it is likely that understanding the UPR in a tractable multicellular model system, such as Arabidopsis, will allow building approaches with a strong potential to contribute to the solution of the problems in humans confronting cellular stress.

We also expect that we will be able to develop transplantable bioindicators of stress without the need of sophisticated instrumentation. These reporters will potentially facilitate the optimization of plant cultivation in hostile environments, such as marginal lands. In addition, because of the conservation of UPR mechanisms in plants and humans, real-time stress bioindicators will likely reflect the perception of stress in other organisms besides plants and thus they may be useful to develop practices to improve quality of life for humans.

In conclusion, not only will our research improve our nation's scientific competitiveness, it will also benefit education and the quality of life on Earth. Life on Earth depends on plants. With sharply growing environmental concerns (global warming, pollution, necessity to expand agriculture to marginal lands, etc.), understanding how plants can resist stress in hostile environments and adopting bioindicators help research to expand agriculture practices and thus ensure continued survival of the ever-growing human population.

REFERENCES

1. Ron D, Walter P. Signal integration in the endoplasmic reticulum unfolded protein response. Nat Rev Mol Cell Biol 2007; 8 (7):519-529.

2. Liu JX, Howell SH. Endoplasmic reticulum protein quality control and its relationship to environmental stress responses in plants. Plant Cell 2010; 22 (9):2930-2942.

3. Iwawaki T, Akai R, Yamanaka S, Kohno K. Function of IRE1 alpha in the placenta is essential for placental development and embryonic viability. Proc Natl Acad Sci U S A 2009; 106(39):16657-16662.

4. Chen Y, Brandizzi F. AtIRE1 and AGB1 independently control two essential unfolded protein response pathways in plants. Plant J 2012; 69: 266-277.

5. Deng Y, Humbert S, Liu JX et al. Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in Arabidopsis. Proc Natl Acad Sci U S A 2011; 108 (17):7247-7252.

6. Tirasophon W, Lee K, Callaghan B, Welihinda A, Kaufman RJ. The endoribonuclease activity of mammalian IRE1 autoregulates its mRNA and is required for the unfolded protein response. Genes Dev 2000; 14 (21):2725-2736.

7. Urano F, Bertolotti A, Ron D. IRE1 and efferent signaling from the endoplasmic reticulum. J Cell Sci 2000; 113 Pt 21:3697-3702.

8. Sidrauski C, Walter P. The transmembrane kinase Ire1p is a site-specific endonuclease that initiates mRNA splicing in the unfolded protein response. Cell 1997; 90 (6):1031-1039.

9. Cox JS, Chapman RE, Walter P. The unfolded protein response coordinates the production of endoplasmic reticulum protein and endoplasmic reticulum membrane. Mol Biol Cell 1997; 8 (9):1805-1814.

10. Kawahara T, Yanagi H, Yura T, Mori K. Endoplasmic reticulum stress-induced mRNA splicing permits synthesis of transcription factor Hac1p/Ern4p that activates the unfolded protein response. Mol Biol Cell 1997; 8 (10):1845-1862.

11. Plongthongkum N, Kullawong N, Panyim S, Tirasophon W. Ire1 regulated XBP1 mRNA splicing is essential for the unfolded protein response (UPR) in Drosophila melanogaster. Biochem Biophys Res Commun 2007; 354 (3):789-794.

12. Shen X, Ellis RE, Lee K et al. Complementary signaling pathways regulate the unfolded protein response and are required for C. elegans development. Cell 2001; 107 (7):893-903.

13. Nagashima Y, Mishiba K, Suzuki E et al. Arabidopsis IRE1 catalyses unconventional splicing of bZIP60 mRNA to produce the active transcription factor. Scientific Reports 2011; 1 (29):DOI: 10.1038/srep00029.

14. Moreno AA, Mukhtar MS, Blanco F et al. IRE1/bZIP60-Mediated Unfolded Protein Response Plays Distinct Roles in Plant Immunity and Abiotic Stress Responses. PLoS One 2012; 7:e31944.

**Task Progress:**

In the previous BRIC-18-2 (Biological Research in Canisters) experiment, we have established differential gene regulation in wild type but also in mutant Arabidopsis with defects in pro-survival management strategies. These findings open new areas of research on plant management of the different stresses that are found in space.
We are currently expanding upon this work through exploration of a larger set of Arabidopsis mutants that have been flown to the International Space Station (ISS) in the BRIC-22 experiment. These mutants are missing genes that are critical in regulating the response to stresses observed previously, but have different molecular functions. We successfully completed both of the preflight tests (Science and Experiment Verification Tests--SVT and EVT) to validate the exact experimental protocols necessary to prepare and analyze the samples returned from BRIC-22.

On August 14, 2017 our experiment was launched on board the SpaceX-12 ISS resupply mission and was successfully executed by ISS astronauts. The samples were preserved on orbit and underwent atmospheric reentry on board the same Dragon capsule in September of 2017. These were then returned to the Brandizzi lab still preserved at -80 degrees C and the samples were processed yielding experimental material that was determined to be in excellent condition. The RNA extracted from these plants is currently undergoing high-throughput sequencing, which will enable us to examine the types of molecular responses activated by plants under spaceflight associated stress, in WT (wild type) and in mutant lines to determine how the mutated genes participate in maintaining the health of the plant under these conditions. We expect to receive the raw sequences from the sequencing facility at Michigan State University (MSU) in two weeks.

Additionally, further research on the genes missing from the Arabidopsis mutants to be tested in BRIC-22 is also being completed in ground-based studies. Through a better understanding of gene function at a cellular and organism level in terrestrial growth conditions, we will be better positioned to explore what role these genes have in spaceflight conditions. This information will be adapted to help produce Arabidopsis lines resistant to these stresses.

**Bibliography Type:**

Description: (Last Updated: 12/08/2025)

**Abstracts for Journals and Proceedings**

Brandizzi F. "Role of the plant unfolded protein response to space stress." 33rd Annual Meeting of the American Society for Gravitational and Space Research, Seattle, WA, October 25-28, 2017.
33rd Annual Meeting of the American Society for Gravitational and Space Research, Seattle, WA, October 25-28, 2017. , Oct-2017

**Articles in Peer-reviewed Journals**

Angelos E, Ruberti C, Kim SJ, Brandizzi F. "Maintaining the factory: the roles of the unfolded protein response in cellular homeostasis in plants." Plant J. 2017 May;90(4):671-82. Epub 2017 Mar 10. https://doi.org/10.1111/tpj.13449 ; PubMed PMID: 27943485; PubMed Central PMCID: PMC5415411 , May-2017