Fiscal Year:	FY 2005	**Task Last Updated:**	FY 01/26/2005
PI Name:	Dickman, J. David Ph.D.		
Project Title:	Development of the otolith system in altered gravity environments		
Division Name:	Space Biology		
Program/Discipline:	SPACE BIOLOGY		
Program/Discipline--Element/Subdiscipline:	SPACE BIOLOGY--Developmental biology		
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Animal Biology: Vertebrate		
Space Biology Cross-Element Discipline:	(1) Reproductive Biology (2) Developmental Biology (3) Neurobiology		
Space Biology Special Category:	None		
PI Email:	ddickman@cns.bcm.edu	**Fax:**	FY
PI Organization Type:	UNIVERSITY	**Phone:**	713-798-4589
Organization Name:	Baylor University/Baylor College of Medicine		
PI Address 1:	Department of Neuroscience		
PI Address 2:	One Baylor Plaza, MS: 295		
PI Web Page:			
City:	Houston	**State:**	TX
Zip Code:	77030-3411	**Congressional District:**	7
Comments:	NOTE PI moved to Baylor in Fall 2011; previously at Washington University, St. Louis, MO (Ed., 12/20/2011)		
Project Type:	Ground	**Solicitation / Funding Source:**	2003 Space Biology 03-OBPR-03
Start Date:	02/01/2004	**End Date:**	09/30/2006
No. of Post Docs:	1	**No. of PhD Degrees:**	
No. of PhD Candidates:	2	**No. of Master' Degrees:**	
No. of Master's Candidates:	0	**No. of Bachelor's Degrees:**	
No. of Bachelor's Candidates:	0	**Monitoring Center:**	NASA ARC
Contact Monitor:		**Contact Phone:**	
Contact Email:			
Flight Program:			
Flight Assignment:			
Key Personnel Changes/Previous PI:			
COI Name (Institution):			
Grant/Contract No.:	NNA04CC52G		
Performance Goal No.:			
Performance Goal Text:			
	In all vertebrates, the otolith system functions to detect head position and movement relative to gravity and to modulate or control gravity depending neuromotor responses. How does gravity influence the development of the otolith receptor cells, their polairzation topography, and their innervating afferents during embryogenesis? The avian otolith system offers an excellent model with which to study the effects of gravity upon development, due to the short maturation period following fertilization and due to the broad existing knowledge of otolith system structure and function in birds. The present project is examining the effects of gravity upon the structure of the avian otolith receptor system. Japanese quails, due to their short incubation time of 15 days are being exposed to different levels of gravity during embryogenesis and their vestibular receptor systems examined using a number of different quantitative anatomical measures. Quail raised from fertilization in either normal gravity, hypergravity (2 - 3G),or microgravity (spaceflight)		

Task Description:	conditions will be compared. The anatomical development of the receptor cell organization and the afferent innervation patterns of 4day, 7 day, and 12 day qauil embryos are being collected for comparison across the different gravity conditions and developmental stages. The primary goal is to define whether gravity is a determinant trophic factor in development of the otolith system during embryogenesis. In order for manned long duration missions to succeed, the effects of altered gravity environments upon nervous system structure and function must be determined. The primary objective of the present proposal is to determine the effects of gravity upon development of the peripheral vestibular system. It is known that even moderate exposure to spaceflight conditions in adult animals can produce modifications in the anatomy, physiology, and neuromotor responses related to the vestibular system. In addition, more limited evidence shows that altered gravity environments can directly affect the development of vestibular receptors. Within the vestibular system, the otolith organs comprise the inertial sensors that respond to linear acceleration or tilts of the head relative to gravity. During spaceflight or centrifugation, the acceleration constant due to normal Earth gravity is eliminated or changed, thereby altering the inertial reference upon which vestibular system responses are based. Does gravity serve as a regulatory factor in the development of the vestibular system and its proper neuromotor function? This project serves as a ground-based investigation that will examine vestibular receptor organ development in quails completely reared in either Earth normal or altered hypergravity environments. The current application will build upon the isolated findings of our recent spaceflight investigation on STS-108, where embryogenesis in microgravity was found to affect vestibular otoconia, afferent innervation, and synapse formation in quails raised to embryonic day 12. Current experiments are designed to specifically examine how otoconial biosynthesis, receptor morphology, afferent innervation, and synaptogenesis are affected by exposure to constant hypergravity conditions during embryogenesis and early post-hatch development in Japanese quails. Objective 1. Determine how altered gravity affects the morphology and molecular biosynthesis of otoconia in birds. Otoconia are dense biomineral particles in the otolith organs that serve as mechanical inertial transducers for the detection of linear acceleration and subsequent receptor cell activation. It is known that in invertebrates and fish, development of the homologous statolith is highly sensitive to altered gravity conditions. We hypothesize that the regulation of otoconial biosynthesis is affected by gravity. In Objective 1, the otoconial morphology and the molecular parameters of otoconial biosynthesis during quail embryogenesis and early post-hatch development will be examined using both normal gravity and hypergravity conditions. Objective 2. Determine how gravity affects the development of otolith receptor cells, their afferent innervation, and synaptogenesis. Two types of receptor cells, three afferent types, and two major forms of hair cell-afferent synapses are known to exist in the otolith sensory epithelium of all amniote vertebrates. These morphological characteristics develop at different times during embryogenesis and post-hatch maturation. We hypothesize that the development of receptor cell distribution, afferent innervation, and synaptogenesis in the otolith system is regulated by gravity. We predict that the rate of maturation, number of cells, number and type of synapses, and afferent terminal innervations will all increase proportionally with the force of gravity. In Objective 2 we will examine these morphological characteristics at different developmental ages in quails raised in either normal or hypergravity environments. In all vertebrate animals, the vestibular system forms an essential component in the production of movement related responses that are critical for the daily function and survival of the animal. During manned spaceflight, vestibular disturbances frequently occur, with approximately 80 – 90 % of current flight crew personnel experiencing some symptoms space motion sickness, such as disorientation, nausea, and emetic attacks during the first 48 - 72 hours of weightlessness (Davis et al., 1988; Homick et al., 1984; Reschke et al., 1998; Thorton et al., 1987). These symptoms can best be described as acute, although they often return after the flight immediately upon reintroduction to the 1g environment. However, more serious vestibular related effects that occur during long duration spaceflight include changes in compensatory eye movements (Clarke et al., 1993; Clement and Berthoz, 1990; Cohen et al., 1992; Kornilova et al., 1993; Yakovleva et al., 1982), muscle tone and reflex (Edgerton and Roy, 1995; Edgerton et al., 1995), orthostatic intolerance (Yates and Kerman, 1998), and postural stability (Anderson et al., 1986; McDonald et al., 1994; Paloski et al., 1992). Although a number of investigators have postulated that the lack of gravity as a constant stimulus during spaceflight produces profound changes in vestibular system function, the basic mechanisms that are affected by changes in gravity are only beginning to be addressed. Several reports have indicated that the structure and function of the vestibular system is affected by exposure to altered gravity conditions. For example, changes in size of otoconia (Fermin et al., 1996; Ross and Donovan, 1986; Wiederhold et al., 2000), in the receptor-afferent morphology (Ross, 1993; 2000; Gaboyard et al., 2002; Raymond et al., 2000), in hair cell conductance (Chabbert et al., 2003), in vestibular afferent responsiveness (Boyle et al., 2001; Bracchi et al., 1975; Correia et al., 1992; Jones et al., 1993), in vestibular central pathways (Anken et al., 2002; Bruce and Fritzsch, 1997; Kozlovskaya et al., 1984; Pomeiano et al., 2002), and in vestibular related neuromotor responses (Dai et al., 1994; Sebastian, 1996; Takabayashi et al., 1987; von Baumgarten et al., 1975) have all been observed in both adult and developing animals exposed for brief periods in either microgravity or hypergravity. The current project seeks to determine how gravity affects the morphology of the vestibular otolith receptor system during embryogenesis and post-hatchling development. Before long duration missions such as a manned presence on the Moon or a Mars flight can be manifested in Spirals 4 and 5, basic knowledge regarding the nature and extent of developmental or adaptational changes in sensorimotor systems that occur as a consequence of exposure to altered gravity conditions must be acquired.
Rationale for HRP Directed Research:	
Research Impact/Earth Benefits:	Many sensory systems are known to enter a critical period of development, during which stimulation of modality specific receptors is required for normal neural function. For example, in the visual system, if deprived of light during the early critical period, normal sight is impaired or eliminated during maturation. In the vestibular system which controls normal postural, eye and head movement responses during motion, no effort to study the effects of gravity upon development have been extensively performed. Thus, it is currently unknown whether altered gravity environments affect vestibular system development and subsequent reflex motor behavior. The current project is examining the trophic influence of gravity upon the development of receptors and their innervating afferent neurons in qauil embryos. Both hypergravity as induced by chronic centrifugation and microgravity as induced during spaceflight are being used to determine the morphological development of vestibular organs at different stages of development. The research should provide basic answers into questions related to vestibular receptor pathology that is known to occur with aging humans and in some disease cases.

Task Progress:

Centrifuge construction: In order to examine the nature of gravity as an epigenetic factor in vestibular system development, we initiated the construction of a hypergravity centrifuge for our laboratory. The centrifuge consists of an 8 foot diameter arm, with a unique Avian Hatchling Habitat (SHOT, Inc.) attached to one end. The hatchling habitat provides stable environments for quail hatchling development during hypergravity exposure at levels ranging between 1 and 3 g chronic loads. In the center of the centrifuge resides a control habitat for hatchlings, where rotational forces are imparted but the gravity load is essentially normal (1.008 g). The centrifuge will be operational by February ’05, so that hypergravity exposures upon quail hatchlings can be performed.
Morphology and biosynthesis of otoconia: Our first project objective was to investigate the morphology of otoconia in quails developed from fertilization in normal and hypergravity environments. Japanese quails have a 15 day incubation time. The otoconia of Japanese quails developed in Earth 1g environments were first examined at each embryonic stage from E4 – E15 using scanning electron microscopy (SEM) and sectioned tissue histology. At developmental stage E4, otoconia were not yet detectable, but at stage E5 immature otoconia were beginning to form in the saccule but not the utricle or lagena. At E6, otoconia began to form in the utricle and lagena. Most of the newly developing otoconia in quails were small with double fluted ends. A quantitative analysis of the number and size of otoconia for animals raised in 1g conditions was performed, with measures taken at each embryonic day and at each of the first four post-hatch weeks. Otoconia from different regions of the maculae were compared, as it is known that otoconia vary in size depending upon macular location in adults. We implemented two quantification methods for examining otoconial growth. The first, provided measurements of the length and width of individual otoconia. Values for over were obtained from each of 3 standard locations (i.e., lateral striola, central, medial periphery) in each otolith organ, counting only otoconia that lay approximately perpendicular to the visualization plane. Over 13,000 otoconia were quantified for the utricle and saccule, at the different developmental stages. For example, at E10 a total of 976 otoconia from 4 utricles were measured; where sizes ranged between 0.9 – 19.7 ?m in length, 0.8 – 8.2 ?m in width, and mean length values (?SD) were 8.0 (±3.6) for striolar, 6.6 (?2.8) for central, and 6.6 (?2.6) for medial peripheral regions. In adult quails, the otoconia were significantly larger. Comparable values for 596 otoconia from adult (>6 week) quails were: length ranged between 1.6 – 45.5 µm, width ranged between 1.2 – 19.7 µm, and mean length values of 18.0 (±9.6) µm for striola, 10.2 (±4.8) µm for central, and 9.1 (±3.3) µm for medial peripheral regions. Mature individual otoconium were characterized by a barrel shape with two trihedral faceted ends. However, initial formation of otoconia at E5 (saccular) and E6 (utricular) maculae was characterized by a double fluted morphology that consisted of an hourglass shape with extended fins forming trihedral angles of 120º. Double fluted otoconia rapidly filled, so that by embryonic day 8 mature otoconia dominated the maculae for the remainder of development through adulthood. Thus, a progression from double fluted to mature forms was noted. Mature utricular otoconia in adult quails averaged 11µm in length and 5µm in width, with length/width ratios of approximately 2.5:1, for all size ranges. Saccular otoconia were smaller, having about 70% the size of utricular otoconia in both length and width. During development, the average size and range of individual otoconia increased nearly linearly for both otolith organs. In the utricular macula, large otoconia were concentrated in the lateral regions of the epithelium. In contrast, otoconia of various sizes were distributed uniformly across the surface of the saccular macula. We have also initiated studies to examine the biosynthesis of otoconia during development. Using an in-situ probe developed in our laboratory, we found that the major protein contained in mammalian and avian otoconia, termed otoconin 90 (OC90) is expressed in quail. We examined Oc90 mRNA expression from stage E4 through post-hatch day P7. Oc90 was expressed in the thin, nondifferentiated epithelia on the roof of saccule, utricle, and lagena but was absent from the sensory neuroepithelia. Interestingly, this unique epithelial expression pattern of Oc90 mRNA in the quail is virtually identical to that of mammals. We observed strong expression of Oc90 at E4 – E8, with declining expression at E9 – E12. By E13 – P7, no expression was observed. These findings suggest that otoconial formation occurs as an interaction of secreted OC90 with other local factors in the gelatinous membrane (Wang et al., 1998; Thalmann et al., 2001). In support of that hypothesis, a novel gene, Otopetrin (Otop1) has been identified as the point mutation source for the naturally occurring mutant tilted (tlt) and mergulhador (mlh) mice that lack otoconia (Hurle et al., 2003). In wild type mice, Otop1 mRNA was found using in-situ hybridization to be expressed in the otolith receptor epithelium, most likely in the support cells, and OTOP1 was identified as a protein found in the acellular matrix of the otoconial membrane in the utricle and saccule. Recently, three separate highly conserved regions of the murine Otop1 cDNA have been identified in expressed sequence tag (EST) clones of chicks, strongly suggesting similar mRNA and protein are also expressed in quails. We are currently constructing in-situ probes for chick Otop1 to be used in the quail. The expression, timing, and localization of OC90 and Otop1 will be directly examined during embryogenesis and post-hatch development for both normal and hypergravity reared animals in the next budget year (Objective 1).

Morphology of avian otolith receptors, afferent innervation, and synaptogenesis: In Objective 2, we hypothesize that the development of vestibular receptor cell distribution, afferent innervation, and synaptogenesis in the otolith organs are dependent upon gravity. We have begun examination of the afferent innervation patterns of quail utricular fibers in embryos exposed to normal and 2g environments. Although incomplete, our initial results suggest that E12 quails exposed to 1g normal or 2g hypergravity conditions show different afferent morphologies and innervation patterns. For example, the axon diameter for fibers penetrating the macula basement membrane of the utricle were different depending upon gravity condition experienced during embryogenesis. Utricular afferents (n> 180 for each gravity condition) of quails raised in 1g were smaller than those of 2g embryo afferents. In terms of innervation pattern, the appearance and size of calyx afferents were delayed in hypergravity as compared to 1g controls. The calyceal terminals were less well developed, with many digiform projections and proto cup shapes. E12 dimorph and bouton afferents both had fewer branches, smaller terminal fields, and less terminals when raised in 1g as compared to 2g embryos. All fibers had larger axon diameters and much increased terminal fields when raised in 2g hypergravity as compared to either 1g control. Since no type I hair cell marker has been available to date, it has been difficult to assess when these cells develop and become innervated by calyceal terminals. However, recent monoclonal antibodies for two proteins may prove useful for identifying the appearance of type I hair cells and the formation of calyceal terminals around them. TUJ1, detects a neuron-specific class III ß tubulin (Lee et al., 1990) and EphA5 detects a receptor tyrosine kinase involved in neuron attraction and repulsion guidance (Knoll and Drescher, 2002). We have utilized these antibodies and observed that TUJ1 detection in adult quail utricles encompasses the outline of calyceal terminals, while EphA5 is expressed at the base of putative type I hair cells only. We believe the EphA5 serves as an attractant, attachment, or signaling receptor for calyceal formation and maintenance, since no EphA5 antibody was observed in type II hair cells. If true, we can expect to utilize EphA5 as a marker for type I hair cell formation before, during and after calyceal formation during development. We will confirm the specificity for EphA5 using TEM of a secondary antibody with an electron dense conjugate (e.g., HRP). More complete data is required to determine the effects of different gravity exposures upon afferent innervation. For example, it remains unclear as to whether hypergravity accelerated and accentuated afferent growth subsides in later post-hatch development. We will further examine these issues during budget year 2 studies.

Bibliography Type:	Description: (Last Updated: 08/06/2012)
Abstracts for Journals and Proceedings	Dickman JD, Huss D, Warchol M. "Development of vestibular otolith receptors in Japanese quails." Poster, Society for Neuroscience annual meeting, November 2004. Society for Neuroscience Abstracts, November 2004. , Nov-2004
Abstracts for Journals and Proceedings	Dickman JD, Lysakowski A, Huss D, Price S. "Vestibular Otolith Development In Spaceflight and Hypergravity." Barany Society meeting, July 2004. Barany Society abstracts, July 2004. , Jul-2004
Abstracts for Journals and Proceedings	Haque A, Ohlemiller K, Angelaki DE, Dickman JD. "Eye and head contributions to avian gaze stabilization." Poster, Society for Neuroscience annual meeting, November 2004. Society for Neuroscience Abstracts, November 2004. , Nov-2004
Articles in Peer-reviewed Journals	Haque A, Dickman JD. "Vestibular gaze stabilization: different behavioral strategies for arboreal and terrestrial birds." J Neurophysiol. 2005 Mar;93(3):1165-73. Epub 2004 Nov 3. PMID: 15525803 , Mar-2005
Articles in Peer-reviewed Journals	Hughes I, Blasiole B, Huss D, Warchol ME, Rath NP, Hurle B, Ignatova E, Dickman JD, Thalmann R, Levenson R, Ornitz DM. "Otopetrin is required for otolith formation in the zebra fish Danio rerio." Dev Biol. 2004 Dec 15;276(2):391-402. PMID: 15581873 , Dec-2004
Articles in Peer-reviewed Journals	Dickman JD, Huss D, Lowe M. "Morphometry of otoconia in the utricle and saccule of developing Japanese quail." Hear Res. 2004 Feb;188(1-2):89-103. PMID: 14759573 , Feb-2004