Fiscal Year:	FY 2007	**Task Last Updated:**	FY 02/21/2007
PI Name:	Peterson, Kevin Ph.D.		
Project Title:	Using Molecular Clocks And Regulatory Networks To Test Origin-Of-Bilateria Scenarios		
Division Name:	Space Biology		
Program/Discipline:	SPACE BIOLOGY		
Program/Discipline--Element/Subdiscipline:	SPACE BIOLOGY--Evolutionary biology		
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Cell & Molecular Biology (2) Animal Biology: Invertebrate		
Space Biology Cross-Element Discipline:	(1) Developmental Biology (2) Musculoskeletal Biology		
Space Biology Special Category:	None		
PI Email:	kevin.peterson@dartmouth.edu	**Fax:**	FY 603-646-1347
PI Organization Type:	UNIVERSITY	**Phone:**	603-646-0215
Organization Name:	Dartmouth College		
PI Address 1:	212 Gilman Hall		
PI Address 2:	North college Street		
PI Web Page:	http://www.dartmouth.edu/ http://dms.dartmouth.edu/		
City:	Hanover	**State:**	NH
Zip Code:	03755	**Congressional District:**	2
Comments:			
Project Type:	Ground	**Solicitation / Funding Source:**	2001 Space Biology 01-OBPR-06
Start Date:	11/01/2002	**End Date:**	10/31/2006
No. of Post Docs:	1	**No. of PhD Degrees:**	0
No. of PhD Candidates:	0	**No. of Master' Degrees:**	0
No. of Master's Candidates:	0	**No. of Bachelor's Degrees:**	1
No. of Bachelor's Candidates:	1	**Monitoring Center:**	NASA ARC
Contact Monitor:		**Contact Phone:**	
Contact Email:			
Flight Program:			
Flight Assignment:			
Key Personnel Changes/Previous PI:			
COI Name (Institution):			
Grant/Contract No.:	NAG2-1584		
Performance Goal No.:			
Performance Goal Text:			
	Because of the wealth of both phylogenetic and developmental data, there is now a tremendous interest in the origin of bilaterian metazoans and the associated paleontological events such as the Cambrian explosion. It is now generally accepted that the latest common ancestor of bilaterians was a fairly complex animal complete with anterior-posterior and dorsal-ventral axes, and a central nervous system. Nonetheless, it is still unclear if the origin of this latest common ancestor shortly preceded the Cambrian explosion, as suggested by the fossil record, or if there was a significant but cryptic period of bilaterian evolution before their first appearance in the fossil record, as suggested by most molecular-clock studies. Understanding which stage of the bilaterian life cycle is primitive for Bilateria, the larval stage or the adult stage, is crucial for deciphering the temporal incongruence between the fossil record and these molecular-clock analyses. To test for homology of larval types, we cloned a gene in the purple sea urchin		

Task Description:	Strongylocentrotus purpuratus called NK2.1, a member of the NK-2 class of homeodomain transcription factors. NK2.1 is expressed at the apical pole of the sea urchin, and transcripts are eventually restricted to the larval apical organ. Because the apical organ is restricted to primary larval forms, homology of the apical organ implies homology of the larva. In order to test this homology, we want to test for the conservation of gene batteries underlying apical pole development in sea urchins and in gastropod molluscs. We have already identified several important upstream factors in the sea urchin, and are in the process of examining these same genes in the mollusc. Determining the polarity of the larval vs. adult bilaterian body plan will then allow for a better understanding of major events in the history of life on Earth including the Cambrian explosion. Our primary objective was completed this past year with the elucidiation of the regulatory network underlying apical organ development in the sea urchin, and examination of this network in the gastropod. Not unexpectedly, this network is not wired the same in the gastropod suggesting that the two different larval forms arose convergently, and the last common ancestor of bilaterians was not constructed like a bilaterian larvae. This was published this year in Evolution and Development (2007, vol. 9(1):10-24). The significance is that these data are consistenet with zooplankton evolving multiple times indepdenently in the earliest Cambrian, and feeding larvae evolving in the earliest Ordovician, both of which important components of many Cambrian explosion scenarios.
Rationale for HRP Directed Research:	
Research Impact/Earth Benefits:	Ascertaining the primitive life cycle of bilaterians will shed tremendous insight into the ecological ramifications of surviving catastrophic environmental perturbations such as "snowball earth". Indeed, current discussions among snowball-earth workers have focused, in part, on the inferred ecology of the animals surviving the glaciation(s) (see Runnegar 2000, and discussion by Schrag & Hoffman 2001 and Hyde et al. 2001). Thus, a proper understanding of the earliest stages of bilaterian evolution is necessary in order to properly understand the geological and paleontological events of the Late Neoproterozoic. Finally, given that bilaterians may have been present for ~ 100 Ma before they made their first appearance in the fossil record, small but potentially complex metazoans may have thrived and diversified without leaving a trace on our own planet in a relatively recent time frame. Discussions centered on discovering complex life throughout the universe should keep this most interesting possibility in mind.
Task Progress:	We have completed our study, and the final data-driven publication was published earlier this year (Dunn et al. 2007 Evolution and Development 9(1):10-24). We show that the the two larval forms develop their respective ciliary tufts using different gene regulatory mechanisms, strongly indicative of evolutionary convergence. This is entirely consistent with our earlier suggestions that the independent evolution of larvae may have been an important driving factor in the Cambrian explosion and the Ordovician diversification. Finally, these data do not support the notion that a larval-like form is indicative of the earliest bilaterians in the late Precambrian.
Bibliography Type:	Description: (Last Updated: 02/23/2007)
Articles in Peer-reviewed Journals	Dunn EF, Moy VN, Angerer LM, Angerer RC, Morris RL, Peterson KJ. "Molecular paleoecology: using gene regulatory analysis to address the origins of complex life cycles in the late Precambrian." Evol Dev. 2007 Jan-Feb;9(1):10-24. PMID: 17227363 , Jan-2007
Articles in Peer-reviewed Journals	Arnone MI, Rizzo F, Annunciata R, Cameron RA, Peterson KJ, Martinez P. "Genetic organization and embryonic expression of the ParaHox genes in the sea urchin S. purpuratus: insights into the relationship between clustering and" Dev Biol. 2006 Dec 1;300(1):63-73. Epub 2006 Aug 4. PMID: 16959236 , Dec-2006
Articles in Peer-reviewed Journals	Bottjer DJ, Davidson EH, Peterson KJ, Cameron RA. "Paleogenomics of echinoderms." Science. 2006 Nov 10;314(5801):956-60. Review. PMID: 17095693 , Nov-2006
Articles in Peer-reviewed Journals	Sea Urchin Genome Sequencing Consortium. "The genome of the sea urchin Strongylocentrotus purpuratus." Science. 2006 Nov 10;314(5801):941-52. PMID: 17095691 , Nov-2006
Articles in Peer-reviewed Journals	Sperling EA, Peterson KJ. "Poriferan paraphyly and its implications for Precambrian paleobiology." Journal of Geological Society of London, in press 2007. , Jan-2007
Articles in Peer-reviewed Journals	Sempere LF, Cole CN, McPeek MA, Peterson KJ. "The phylogenetic distribution of metazoan microRNAs: insights into evolutionary complexity and constraint." J Exp Zoolog B Mol Dev Evol. 2006 Nov 15;306(6):575-88. PMID: 16838302 , Nov-2006