Fiscal Year:	FY 2026	**Task Last Updated:**	FY 12/18/2025
PI Name:	Luderer, Ulrike M.D., Ph.D.		
Project Title:	Ovarian Cancer and Space Radiation		
Division Name:	Human Research		
Program/Discipline:			
Program/Discipline--Element/Subdiscipline:			
Joint Agency Name:		**TechPort:**	No
Human Research Program Elements:	(1) **SR**:Space Radiation		
Human Research Program Risks:	(1) **Cancer**:Risk of Radiation Carcinogenesis		
Space Biology Element:	None		
Space Biology Cross-Element Discipline:	None		
Space Biology Special Category:	None		
PI Email:	uluderer@uci.edu	**Fax:**	FY
PI Organization Type:	UNIVERSITY	**Phone:**	949-824-8081
Organization Name:	University of California - Irvine		
PI Address 1:	Center for Occupational and Environmental Health		
PI Address 2:	856 Health Sciences Rd, Suite 3200, Zotcode 1830		
PI Web Page:			
City:	Irvine	**State:**	CA
Zip Code:	92697	**Congressional District:**	45
Comments:			
Project Type:	Ground	**Solicitation / Funding Source:**	2018 HERO 80JSC018N0001-Crew Health and Performance (FLAGSHIP, OMNIBUS). Appendix A-Flagship, Appendix B-Omnibus
Start Date:	08/20/2019	**End Date:**	08/19/2025
No. of Post Docs:	0	**No. of PhD Degrees:**	0
No. of PhD Candidates:	1	**No. of Master' Degrees:**	3
No. of Master's Candidates:	3	**No. of Bachelor's Degrees:**	2
No. of Bachelor's Candidates:	4	**Monitoring Center:**	NASA JSC
Contact Monitor:	Zawaski, Janice	**Contact Phone:**	
Contact Email:	janice.zawaski@nasa.gov		
Flight Program:			
Flight Assignment:	NOTE: End date changed to 8/19/2025 per NSSC information (Ed., 9/10/24) NOTE: End date changed to 8/19/2024 per NSSC information (Ed., 7/6/23) NOTE: End date changed to 8/19/2023 per NSSC information (Ed., 4/3/23) NOTE: End date changed to 8/19/2022 per HRP Space Radiation (Ed., 8/3/21)		
Key Personnel Changes/Previous PI:	June 2022 report: Eleanor Blakely, Ph.D. (Biosciences Area, Lawrence Berkeley Lab) and Polly Chang, Ph.D. (Biosciences Division, SRI International) were added as CoInvestigators (Ed., 8/10/22).		
COI Name (Institution):	Blakely, Eleanor (Biosciences Area, Lawrence Berkeley Lab) Chang, Polly (Biosciences Division, SRI International)		
Grant/Contract No.:	80NSSC19K1620		
Performance Goal No.:			
Performance Goal Text:			

Task Description:	Thirty percent of astronauts are women, but the risks of space radiation to women's reproductive health and risks of gynecological cancers remain poorly understood. Radiation treatment for cancer is known to cause temporary infertility and premature menopause. Premature menopause increases women's risks for cardiovascular disease, osteoporosis, and Alzheimer's disease. In addition, animal studies and studies of atomic bomb survivors have shown that radiation exposure increases the risk for ovarian cancer. Ovarian cancer has a high mortality rate and is the leading cause of gynecological cancer deaths in women. To best protect the health of women astronauts, it is important to understand whether space radiation has similar effects on the ovary as the types of radiation exposure that are common on Earth. Our prior pilot study showed that the ovary is highly sensitive to follicle destruction by charged particle radiation, typical of exposures in space. Exposure to charged iron and oxygen particles resulted in dose-dependent follicle depletion and premature ovarian failure. Exposure to charged iron particles induced epithelial ovarian tumors later in life; ovarian tissues from oxygen charged particle irradiated mice of two strains and charged iron irradiated mice of the second mouse strain were archived for future analysis for tumor endpoints. We propose to leverage these stored tissue and blood samples, together with ovaries from gamma-irradiated mice from the NASA tissue archive to 1) compare ovarian tumor prevalence and molecular characteristics after low dose charged particle irradiation (oxygen and iron ions) with gamma irradiation in adult female mice; 2) examine the persistence and types of ovarian oxidative damage after irradiation and evaluate serum concentrations of a clinically utilized biomarker of ovarian reserve, Anti-Müllerian Hormone (AMH), as a potential early biomarker of ovarian tumorigenesis. We will quantify the effects of charged particles on numbers of ovarian follicles and ovarian tumor number and size. We will use in situ methods to assess oxidative damage and to molecularly characterize the ovarian tumors. Our analyses will provide critical insights into whether preneoplastic changes in ovarian follicle numbers, serum AMH, as well as ovarian oxidative damage caused by exposure to charged particles demonstrate similar dose-response as ovarian tumor induction. The analyses will also examine the relative biological effectiveness of gamma versus charged particle irradiation for these endpoints. These studies will help to fill important gaps in our understanding of the effects of space radiation on ovarian function and ovarian carcinogenesis and will lead to better ways to prevent ovarian cancer and protect reproductive health in women astronauts.
Rationale for HRP Directed Research:	
Research Impact/Earth Benefits:	The research increases understanding of the effects of low dose gamma- and charged particle radiation on the ovary. Both gamma and charged particle radiation are used for cancer therapy. Therefore, it is important to understand side-effects of radiation therapy that may impact cancer survivors' quality of life.
Task Progress:	Aim 1: Utilize archived ovaries to compare ovarian tumor induction by irradiation with a mixed heavy ion beam of silicon, titanium, and iron ions, oxygen ions only, or silicon charged particles only. Fixed ovaries from 50 mice sacrificed at 16 months after irradiation with 0.3 or 0.6 Gy mixed heavy ion beam or control, unirradiated mice were embedded in paraffin, serially sectioned, and every 5th or 10th section was stained with hematoxylin and eosin. All ovaries were reviewed by a board-certified veterinary pathologist. There was a dose-dependent, highly statistically significant increase in ovarian tubular adenomas, with 91% of mice in the 0.6 Gy mixed beam group having unilateral or bilateral tumors, 8% having a unilateral tumor in the 0.3 Gy mixed beam group, and no ovarian tumors found in the control mice. There was also a highly statistically significant increase in hyperplasia and fibrosis of the ovarian surface epithelium in the 0.3 Gy group. Hyperplasia of the ovarian surface epithelium is believed the be a precursor to ovarian tubular adenomas, so these results suggest that these mice would have eventually developed tumors. Tubular adenomas are epithelial ovarian tumors; and positive immunostaining of cells lining the tubular structures for epithelial markers, using a pancytokeratin antibody and a keratin 19 antibody, confirmed that the tumors are epithelial. Interestingly, the cells between the tubular structures of these tumors stained positively for FOXL2, a granulosa cell marker, and/or CYP17A1, a theca cell marker, indicating that these tumors are of mixed cellular origin. The tumors had very few dividing cells, determined by immunostaining for the mitosis marker Ki67, consistent with the non-malignant nature of tubular adenomas. As expected in mice of advanced age (19 months), there were relatively few follicles in the ovaries of the control mice. Nonetheless, counts of ovarian follicles in these ovaries demonstrated statistically significantly fewer ovarian follicles in the irradiated groups compared to the control group. Unilateral ovaries from 30 mice per group irradiated with 0, 0.04, 0.08, 0.12, and 0.32 Gy silicon charged particles were similarly processed. 0.32 Gy silicon ovaries were reviewed by a board-certified pathologist, and only one ovarian tumor was found in the 0.32 Gy group. Therefore, the ovaries from mice irradiated with lower doses of silicon ions were not evaluated for tumors. Ovarian follicle counts on ovaries from 0 and 0.32 Gy dose groups showed that 0.32 Gy silicon ions statistically significantly decreased ovarian follicle numbers compared to 0 Gy controls. Ovaries from mice of two different strains, collected 15 months after irradiation, with 0 or 50 cGy oxygen ions, were also analyzed for follicle numbers and tumors. Follicle numbers were significantly decreased, and the percentage of mice with ovarian tumors was significantly increased 15 mos after irradiation with 0.50 Gy oxygen ions. Tumors were positive for cytokeratin in the cells of the tubules, positive for the granulosa cell marker FOXL2 and theca cell marker CYCP17A1 in the cells between the tubules, and negative for alpha smooth muscle actin. Ovaries from the mixed heavy ion beam-irradiated and control mice were also examined for markers of oxidative damage, inflammation, and fibrosis. We did not find increased ovarian oxidative damage (measured using a marker of oxidative lipid damage), or inflammation (assessed by measuring the area of the ovary positive for immune cells called macrophages), or increased age-related accumulation of proteins called lipofuscin, or increased fibrosis (assessed by measuring the area of the ovaries containing two types of collagen) in ovaries of irradiated compared to control mice. Aim 2: Utilize archived ovaries harvested at various time points after irradiation with low doses of oxygen or iron-charged particles to examine the persistence of ovarian oxidative lipid, protein, and DNA damage, archived serum to measure a biomarker of ovarian reserve, and evaluate these as potential early biomarkers of ovarian tumorigenesis. Irradiate mice with low doses of gamma-radiation and harvest ovaries at 1 week after irradiation to compare ovarian follicle depletion by charged iron or oxygen particles with gamma-radiation. Ovarian follicle counts in the gamma-irradiated mice have been completed. There were statistically significant effects of dose on primordial, primary, and secondary follicle numbers, with fewer primordial, primary, and secondary follicles in the 0.50 Gy group compared to the 0 Gy group. The decreases at the lower doses of 0.05 and 0.15 Gy were less than we previously observed at one week after irradiation with similar doses of charged oxygen particles, supporting that the latter are more damaging to the ovaries than gamma-rays.

	Serum Anti-Müllerian Hormone (AMH) concentrations at one week after irradiation did not vary significantly with dose of gamma- or 56Fe-radiation. There was a statistically significant effect of 16O irradiation dose (P=0.040), with dose-dependent decrease in serum AMH concentrations at 0.30 Gy compared to 0 Gy, but no difference in concentrations of AMH between the 0 compared to 0.05 and 0.50 Gy groups. Serum AMH was significantly decreased in ovaries harvested 8 weeks after 0.50 Gy 56Fe irradiation. The lack of dose-dependent decrease in serum AMH contrasts with the pronounced dose-dependent decrease in ovarian follicle numbers at one week after irradiation in the same mice exposed to 56Fe or 16O in our prior studies or the mice exposed to 0.5 Gy gamma-radiation described above. Primordial, primary, and secondary ovarian follicle numbers were also significantly decreased in the ovaries of mice irradiated with 0.50 Gy 16O and sacrificed 16 months after irradiation. As noted above, we did not find increased oxidative damage to proteins, lipids, or DNA at 15 months after irradiation with a mixed heavy ion beam. Conclusions We conclude that mixed heavy ion irradiation at 0.6 Gy total dose potently induces ovarian tumors, with 91% of the mice having ovarian tumors at 16 months after irradiation, while the 0.3 Gy total dose was much less effective at inducing ovarian tumors. We further conclude that irradiation with a beam of 0.5 Gy oxygen ions also potently induced ovarian tumors with 52% of the irradiated B6C3F1 mice and 63% of the irradiated C57BL/6J mice having unilateral or bilateral ovarian tumors. We also conclude that serum AMH concentration at one week after irradiation does not correlate with primordial follicle numbers and therefore is not a useful biomarker of ovarian reserve at this time point after irradiation.
Bibliography Type:	Description: (Last Updated: 02/12/2026)
Abstracts for Journals and Proceedings	Pore A, Leon Parada KN, Lawson G, Luderer U. "Analyzing the effects of charged oxygen particle radiation on the ovaries of female mice." UC Irvine Undergraduate Research Opportunities Program Symposium, Irvine, California, May 12-16, 2025. Abstracts. UC Irvine Undergraduate Research Opportunities Program Symposium, Irvine, California, May 12-16, 2025. , May-2025
Articles in Peer-reviewed Journals	Leon Parada KN, Lawson G, Chang P, Blakely EA, Bajwa L, Gramajo-Aponte K, Getze S, Luderer U. "Irradiation with a mixed heavy ion beam induces ovarian follicle loss and dose-dependent mixed ovarian tumor development." Biol Reprod. 2026 Jan 21:ioag019. https://doi.org/10.1093/biolre/ioag019 ; PMID: 41562572 , Jan-2026