

PROBABILISTIC MACHINE LEARNING

LECTURE 23

EM


Philipp Hennig

20 July 2023

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



FACULTY OF SCIENCE
DEPARTMENT OF COMPUTER SCIENCE
CHAIR FOR THE METHODS OF MACHINE LEARNING



A special option: The EM algorithm

Maximizing model Evidence

The general recipe for hyperparameter inference:

Consider a model with **parameters** θ , **observed data** y and **latent variables** z

- Ideally, we would like to maximize the **marginal (log-) likelihood (evidence)**

$$\log p(y \mid \theta) = \log \left(\int p(y, z \mid \theta) dz \right) \quad (\star)$$

- if we can not do this integral, we can try **Laplace**. This is nearly always *possible* (if $\log p(y \mid z)$ is twice differentiable), but it is fundamentally an approximation.
- however, *in some cases*, we may be able to compute the **Expectation** of the “complete data” log **likelihood** (for a fixed value θ_*)

$$q(\theta, \theta_*) = \int p(z \mid y, \theta_*) \log p(y, z \mid \theta) dz$$

and then **Maximize** $q(\theta, \theta_*)$ **with respect to** θ . This can be easier than (\star) because the log “simplifies things” (e.g. turns products into sums, thus factors into components).

Definition: The Expectation Maximization (EM) algorithm:

Consider a model with parameters θ , observed data y and latent variables z .

while not converged, do:

E compute the Expected complete data log-likelihood

$$q(\theta, \theta_t) = \int p(z | y, \theta_t) \log p(y, z | \theta) dz$$

M Set θ_{t+1} to Maximize $\theta_{t+1} = \arg \max_{\theta} q(\theta, \theta_{t+1})$.

EM is an attempt to maximize the evidence $p(y | \theta)$. Why does it work?

An Observation

maximizing a lower bound

- We constructed an approximate distribution $q(z) = p(z | x, \theta)$ for our latent quantity
- For *any* such approximation $q(z)$ (if $q(z) > 0$ wherever $p(x, z | \theta) > 0$):

$$\begin{aligned} \log p(x | \theta) &= \log \int p(x, z | \theta) dz &= \log \int q(z) \frac{p(x, z | \theta)}{q(z)} dz \\ &\geq \int q(z) \log \frac{p(x, z | \theta)}{q(z)} dz &=: \mathcal{L}(q) \end{aligned}$$

Theorem (Jensen's (1906) inequality)

Let $(\Omega, \mathcal{A}, \mu)$ be a probability space, g be a real-valued, μ -integrable function and ϕ be a convex function on the real line. Then

$$\phi \left(\int_{\Omega} g d\mu \right) \leq \int_{\Omega} \phi \circ g d\mu.$$



- ▶ We constructed an approximate distribution $q(z) = p(z | x, \theta)$ for our latent quantity
- ▶ For *any* such approximation $q(z)$ (if $q(z) > 0$ wherever $p(x, z | \theta) > 0$):

$$\begin{aligned}\log p(x | \theta) &= \log \int p(x, z | \theta) dz &&= \log \int q(z) \frac{p(x, z | \theta)}{q(z)} dz \\ &\geq \int q(z) \log \frac{p(x, z | \theta)}{q(z)} dz &&=: \mathcal{L}(q)\end{aligned}$$

- ▶ Thus, by maximizing the RHS in θ in the M-step, we increase a **lower bound on the Evidence**
- ▶ $\mathcal{L}(q)$ is thus called the **Evidence Lower Bound (ELBO)**
- ▶ But can we be sure that this increases the Evidence? To show that this is the case, we will now establish that the E-step makes the bound *tight* at the local θ .



$$\begin{aligned}\mathcal{L}(q) &= \int q(z) \log \frac{p(x, z | \theta)}{q(z)} dz \\ &= \int q(z) \log \frac{p(z | x, \theta) \cdot p(x | \theta)}{q(z)} dz \\ &= \int q(z) \log \frac{p(z | x, \theta)}{q(z)} dz + \log p(x | \theta) \int q(z) dz\end{aligned}$$

thus

$$\begin{aligned}\log p(x | \theta) &= \mathcal{L}(q) - \int q(z) \log \frac{p(z | x, \theta)}{q(z)} dz \\ &= \mathcal{L}(q) - D_{\text{KL}}(q \| p(z | x, \theta))\end{aligned}$$

Richard A. Leibler, 1914–2003

The Kullback-Leibler divergence satisfies $D_{\text{KL}}(q \| p) \geq 0$ with
 $D_{\text{KL}}(q \| p) = 0 \iff q \equiv p$

Solomon Kullback, 1907–1994



EM maximizes the ELBO / minimizes Free Energy


a more general view

exposition based on C.M. Bishop, 2006 §9.4

$$\log p(x | \theta) = \mathcal{L}(q, \theta) + D_{\text{KL}}(q \| p(z | x, \theta))$$

$$\mathcal{L}(q, \theta) = \int q(z) \log \left(\frac{p(x, z | \theta)}{q(z)} \right) dz$$

$$D_{\text{KL}}(q \| p(z | x, \theta)) = - \int q(z) \log \left(\frac{p(z | x, \theta)}{q(z)} \right) dz$$



EM maximizes the ELBO / minimizes Free Energy

a more general view


exposition based on C.M. Bishop, 2006 §9.4

$$\log p(x | \theta) = \mathcal{L}(q, \theta) + D_{\text{KL}}(q \| p(z | x, \theta))$$

$$\mathcal{L}(q, \theta) = \int q(z) \log \left(\frac{p(x, z | \theta)}{q(z)} \right) dz$$

$$D_{\text{KL}}(q \| p(z | x, \theta)) = - \int q(z) \log \left(\frac{p(z | x, \theta)}{q(z)} \right) dz$$

E -step: $q(z) = p(z | x, \theta_{\text{old}})$, thus $D_{\text{KL}}(q \| p(z | x, \theta_i)) = 0$



EM maximizes the ELBO / minimizes Free Energy

a more general view

exposition based on C.M. Bishop, 2006 §9.4

$$\log p(x | \theta) = \mathcal{L}(q, \theta) + D_{\text{KL}}(q \| p(z | x, \theta))$$


$$\mathcal{L}(q, \theta) = \int q(z) \log \left(\frac{p(x, z | \theta)}{q(z)} \right) dz$$

$$D_{\text{KL}}(q \| p(z | x, \theta)) = - \int q(z) \log \left(\frac{p(z | x, \theta)}{q(z)} \right) dz$$

E-step: $q(z) = p(z | x, \theta_{\text{old}})$, thus $D_{\text{KL}}(q \| p(z | x, \theta_i)) = 0$

M-step: **Maximize ELBO**

$$\begin{aligned}
 \theta_{\text{new}} &= \arg \max_{\theta} \int q(z) \log p(x, z | \theta) dz \\
 &= \arg \max_{\theta} \mathcal{L}(q, \theta) + \int q(z) \log q(z) dz
 \end{aligned}$$





Setting: Want to find *maximum likelihood* (or MAP) estimate for a model involving a **latent** variable

$$\theta_* = \arg \max_{\theta} [\log p(x | \theta)] = \arg \max_{\theta} \left[\log \left(\int p(x, z | \theta) dz \right) \right]$$

Algorithm: Initialize θ_0 , then iterate:

1. Compute $q(z) = p(z | x, \theta_{\text{old}})$, **thereby setting** $D_{\text{KL}}(q || p(z | x, \theta)) = 0$
2. Set θ_{new} to the **Maximize the Evidence Lower Bound**

$$\theta_{\text{new}} = \arg \max_{\theta} \mathcal{L}(q, \theta) = \arg \max_{\theta} \int q(z) \log \left(\frac{p(x, z | \theta)}{q(z)} \right) dz$$

3. Check for convergence of either the log likelihood, or θ .

dealing with the data is the main challenge, adding a prior is easy

- It is straightforward to extend EM to maximize a **posterior** instead of a likelihood. Just add a log prior for θ :

Initialize θ_0 , then iterate between

1. Compute $q(z) = p(z \mid x, \theta_{\text{old}})$, thereby setting $D_{\text{KL}}(q \parallel p(z \mid x, \theta)) = 0$
2. Set θ_{new} to the **Maximize the Evidence Lower Bound**

$$\theta_{\text{new}} = \arg \max_{\theta} \int q(z) \log \left(\frac{p(x, z \mid \theta) p(\theta)}{q(z)} \right) dz = \arg \max_{\theta} \mathcal{L}(q, \theta) + \log p(\theta)$$

3. Check for convergence of either the log likelihood, or θ .

This maximizes

$$\begin{aligned} \log p(x \mid \theta) + \log p(\theta) &\geq \mathcal{L}(q, \theta) + \log p(\theta) \\ &\triangleq \log p(\theta \mid x) \end{aligned}$$

why is it even useful to build an iterative update in this way?

If $p(x, z \mid \theta)$ is an **exponential family** with θ as the natural parameters, then

$$p(x, z) = \exp(\phi(x, z)^\top \theta - \log Z(\theta))$$


$$\mathcal{L}(q(z), \theta) = \mathbb{E}_{q(z)}(\phi(x, z)^\top \theta - \log Z(\theta)) = \mathbb{E}_{q(z)}[\phi(x, z)]^\top \theta - \log Z(\theta)$$

$$\nabla_{\theta} \mathcal{L}(q(z), \theta) = 0 \quad \Rightarrow \quad \nabla_{\theta} \log Z(\theta) = \mathbb{E}_{p(x, z)}[\phi(x, z)] = \mathbb{E}_{q(z)}[\phi(x, z)]$$

and optimization may be analytic (example below: Gaussian Mixture Models).

Example

Gaussian Mixture Models




Example

Gaussian Mixture Models



$$p(\mathbf{x} \mid \pi, \mu, \Sigma) = \prod_{n=1}^N \sum_{k=1}^K \pi_k \mathcal{N}(x_n \mid \mu_k, \Sigma_k),$$
$$\pi_k \in [0, 1], \sum_k \pi_k = 1$$
$$p(\mathbf{x}, \mathbf{z} \mid \pi, \mu, \Sigma) = \prod_{n=1}^N \prod_{k=1}^K \pi_k^{z_{nk}} \mathcal{N}(x_n \mid \mu_k, \Sigma_k)^{z_{nk}}$$






EM for Gaussian Mixtures

introducing a class membership variable z factorizes the likelihood

- Want to maximize, as function of $\theta := (\pi_k, \mu_k, \Sigma_k)_{k=1, \dots, K}$

$$\log p(x \mid \pi, \mu, \Sigma) = \sum_{n=1}^N \log \left(\sum_{k=1}^K \pi_k \mathcal{N}(x_n; \mu_k, \Sigma_k) \right)$$



EM for Gaussian Mixtures


introducing a class membership variable z factorizes the likelihood

- Want to maximize, as function of $\theta := (\pi_k, \mu_k, \Sigma_k)_{k=1,\dots,K}$

$$\log p(x \mid \pi, \mu, \Sigma) = \sum_{n=1}^N \log \left(\sum_{k=1}^K \pi_k \mathcal{N}(x_n; \mu_k, \Sigma_k) \right)$$

- Instead, maximizing the “complete data” likelihood is easier:

$$\begin{aligned}
 \log p(x, z \mid \pi, \mu, \Sigma) &= \log \prod_{n=1}^N \prod_{k=1}^K \pi_k^{z_{nk}} \mathcal{N}(x_n; \mu_k, \Sigma_k)^{z_{nk}} \\
 &= \sum_{n=1}^N \sum_{k=1}^K z_{nk} \underbrace{(\log \pi_k + \log \mathcal{N}(x_n; \mu_k, \Sigma_k))}_{\text{easy to optimize (exponential families!)}}
 \end{aligned}$$



E-Step: Compute $p(z \mid x, \theta)$:

$$\begin{aligned} p(z_{nk} = 1 \mid x_n, \pi, \mu, \Sigma) &= \frac{p(z_{nk} = 1)p(x_n \mid z_{nk} = 1)}{\sum_{k'=1}^K p(z_{nk'} = 1)p(x_n \mid z_{nk'} = 1)} \\ &= \frac{\pi_k \mathcal{N}(x_n; \mu_k, \Sigma_k)}{\sum_{k'} \pi_{k'} \mathcal{N}(x_n; \mu_{k'}, \Sigma_{k'})} =: r_{nk} \end{aligned}$$

Note that discrete distributions $q(z_{nk} = 1) = r_{nk}$ have expectation $\mathbb{E}_q[z_{nk}] = r_{nk}$

M-Step: Maximize ELBO

$$\mathbb{E}_{p(z|x,\theta)} (\log p(\mathbf{x}, \mathbf{z} \mid \theta)) = \sum_{n=1}^N \sum_{k=1}^K r_{nk} (\log \pi_k + \log \mathcal{N}(x_n; \mu_k, \Sigma_k))$$

$$\mathbb{E}_{p(z|x, \theta)} (\log p(x, z | \theta)) = \sum_{n=1}^N \sum_{k=1}^K r_{nk} (\log \pi_k + \log \mathcal{N}(x_n; \mu_k, \Sigma_k))$$

To maximize w.r.t. μ set gradient of ELBO to 0:

$$\begin{aligned} \nabla_{\mu_\ell} \mathbb{E}_{p(z|x, \theta)} (\log p(x, z | \theta)) &= - \sum_{n=1}^N r_{n\ell} \Sigma_\ell^{-1} (x_n - \mu_\ell) \stackrel{!}{=} 0 \\ \Rightarrow \quad \mu_\ell &= \frac{1}{R_\ell} \sum_{n=1}^N r_{n\ell} x_n \quad R_j := \sum_{n=1}^N r_{nj} \end{aligned}$$

$$\mathbb{E}_{p(\mathbf{z}|\mathbf{x},\theta)} (\log p(\mathbf{x}, \mathbf{z} \mid \theta)) = \sum_{n=1}^N \sum_{k=1}^K r_{nk} (\log \pi_k + \log \mathcal{N}(x_n; \mu_k, \Sigma_k))$$

To maximize w.r.t. Σ set gradient of ELBO to 0
(note $\partial|\Sigma|^{-1/2}/\partial\Sigma = -\frac{1}{2}|\Sigma|^{-3/2}|\Sigma|\Sigma^{-1}$ and $\partial(v^\top\Sigma^{-1}v)/\partial\Sigma = -\Sigma^{-1}vv^\top\Sigma^{-1}$):

$$\begin{aligned} \nabla_{\Sigma_\ell} \mathbb{E}_{p(\mathbf{z}|\mathbf{x},\theta)} (\log p(\mathbf{x}, \mathbf{z} \mid \theta)) &= -\frac{1}{2} \sum_{n=1}^N r_{n\ell} \left(\Sigma_\ell^{-1} (x_n - \mu_\ell)(x_n - \mu_\ell)^\top \Sigma_\ell^{-1} - \Sigma_\ell^{-1} \right) \\ \Rightarrow \quad \Sigma_\ell &= \frac{1}{R_\ell} \sum_{n=1}^N r_{n\ell} (x_n - \mu_\ell)(x_n - \mu_\ell)^\top \quad R_\ell := \sum_{n=1}^N r_{n\ell} \end{aligned}$$

$$\mathbb{E}_{p(\mathbf{z}|\mathbf{x},\theta)} (\log p(\mathbf{x}, \mathbf{z} \mid \theta)) = \sum_{n=1}^N \sum_{k=1}^K r_{nk} (\log \pi_k + \log \mathcal{N}(\mathbf{x}_n; \mu_k, \Sigma_k))$$


To maximize w.r.t. π , enforce $\sum_k \pi_k = 1$ by introducing Lagrange multiplier λ and optimize

$$\nabla_{\pi_\ell} \mathbb{E}_{p(\mathbf{z}|\mathbf{x},\theta)} (\log p(\mathbf{x}, \mathbf{z} \mid \theta)) + \lambda \left(\sum_{k=1}^K \pi_k - 1 \right) = \sum_{n=1}^N r_{n\ell} \frac{1}{\pi_\ell} + \lambda \stackrel{!}{=} 0$$
$$\pi_\ell = -\frac{1}{\lambda} \sum_{n=1}^N r_{n\ell} = -\frac{1}{\lambda} R_\ell$$

$$\sum_{k=1}^K \pi_k = 1 \Rightarrow \lambda = -N \quad \text{and} \quad \pi_\ell = \frac{R_\ell}{N}$$


EM for Gaussian Mixtures

example run




EM for Gaussian Mixtures

example run




EM for Gaussian Mixtures

example run




EM for Gaussian Mixtures

example run



EM for Gaussian Mixtures

example run



The EM algorithm:

- ▶ to find *maximum likelihood* (or MAP) estimate for a model involving a **latent** variable

$$\boldsymbol{\theta}_* = \arg \max_{\boldsymbol{\theta}} [\log p(\mathbf{x} \mid \boldsymbol{\theta}) p(\boldsymbol{\theta})] = \arg \max_{\boldsymbol{\theta}} \left[\log \left(\sum_z p(\mathbf{x}, z \mid \boldsymbol{\theta}) p(\boldsymbol{\theta}) \right) \right]$$

- ▶ Initialize $\boldsymbol{\theta}_0$, then iterate (checking convergence of either the log likelihood, or $\boldsymbol{\theta}$)

E Compute $p(z \mid \mathbf{x}, \boldsymbol{\theta}_{\text{old}})$, thereby setting $D_{\text{KL}}(q \parallel p(z \mid \mathbf{x}, \boldsymbol{\theta})) = 0$

M Set $\boldsymbol{\theta}_{\text{new}}$ to the **Maximize the Expectation Lower Bound**

$$\boldsymbol{\theta}_{\text{new}} = \arg \max_{\boldsymbol{\theta}} \mathcal{L}(q, \boldsymbol{\theta}) = \arg \max_{\boldsymbol{\theta}} \sum_z q(z) \log \left(\frac{p(\mathbf{x}, z \mid \boldsymbol{\theta}) p(\boldsymbol{\theta})}{q(z)} \right)$$

Please cite this course, as

```
@techreport{Tuebingen_ProbML23,
  title =
    {Probabilistic Machine Learning},
  author = {Hennig, Philipp},
  series = {Lecture Notes
    in Machine Learning},
  year = {2023},
  institution = {Tübingen AI Center}}
```

