

Program 7 : Knowledge Base Resolution

Code:

```
def tell(kb, rule):
    kb.append(rule)

combinations = [(True, True, True), (True, True, False),
                (True, False, True), (True, False, False),
                (False, True, True), (False, True, False),
                (False, False, True), (False, False, False)]

def ask(kb, q):
    for c in combinations:
        s = all(rule(c) for rule in kb)
        f = q(c)
        print(s, f)
        if s != f and s != False:
            return 'Does not entail'
    return 'Entails'

kb = []

# Get user input for Rule 1
rule_str = input("Enter Rule 1 as a lambda function (e.g., lambda x: x[0] or x[1] and (x[0] and x[1]): ")
r1 = eval(rule_str)
tell(kb, r1)

# Get user input for Query
query_str = input("Enter Query as a lambda function (e.g., lambda x: x[0] and x[1] and (x[0] or x[1]): ")
q = eval(query_str)

# Ask KB Query
result = ask(kb, q)
print(result)
```


Observation:

Date : 29/12/23

Knowledge Based Resolution

```
def negate_literal(literal)
  if literal[0] == '~'
    return literal[1:]
  else
    return '~' + literal
```

```
def resolve(c1, c2):
    resolved_clause = set(c1) | set(c2)
    for literal in c1:
        if negate_literal(literal) in c2:
            resolved_clause.remove(literal)
            (negate_literal(literal))
    return tuple(resolved_clause)
```

$$LB: (A \vee \sim B) \wedge (B \vee \sim C) \wedge C \wedge \sim A$$


Output:

Kanjika Singh-1BM21CS086

Step	Clause	Derivation

1.	$R \vee \sim P$	Given.
2.	$R \vee \sim Q$	Given.
3.	$\sim R \vee P$	Given.
4.	$\sim R \vee Q$	Given.
5.	$\sim R$	Negated conclusion.
6.		Resolved $R \vee \sim P$ and $\sim R \vee P$ to $R \vee \sim R$, which is in turn null.

A contradiction is found when $\sim R$ is assumed as true. Hence, R is true.

Kanjika Singh-1BM21CS086

Step	Clause	Derivation

1.	$P \vee Q$	Given.
2.	$\sim P \vee R$	Given.
3.	$\sim Q \vee R$	Given.
4.	$\sim R$	Negated conclusion.
5.	$Q \vee R$	Resolved from $P \vee Q$ and $\sim P \vee R$.
6.	$P \vee R$	Resolved from $P \vee Q$ and $\sim Q \vee R$.
7.	$\sim P$	Resolved from $\sim P \vee R$ and $\sim R$.
8.	$\sim Q$	Resolved from $\sim Q \vee R$ and $\sim R$.
9.	Q	Resolved from $\sim R$ and $Q \vee R$.
10.	P	Resolved from $\sim R$ and $P \vee R$.
11.	R	Resolved from $Q \vee R$ and $\sim Q$.
12.		Resolved R and $\sim R$ to $R \vee \sim R$, which is in turn null.

A contradiction is found when $\sim R$ is assumed as true. Hence, R is true.

Kanjika Singh-1BM21CS086

Step	Clause	Derivation
------	--------	------------

1.	$P \vee Q$	Given.
2.	$P \vee R$	Given.
3.	$\sim P \vee R$	Given.
4.	$R \vee S$	Given.
5.	$R \vee \sim Q$	Given.
6.	$\sim S \vee \sim Q$	Given.
7.	$\sim R$	Negated conclusion.
8.	$Q \vee R$	Resolved from $P \vee Q$ and $\sim P \vee R$.
9.	$P \vee \sim S$	Resolved from $P \vee Q$ and $\sim S \vee \sim Q$.
10.	P	Resolved from $P \vee R$ and $\sim R$.
11.	$\sim P$	Resolved from $\sim P \vee R$ and $\sim R$.
12.	$R \vee \sim S$	Resolved from $\sim P \vee R$ and $P \vee \sim S$.
13.	R	Resolved from $\sim P \vee R$ and P .
14.	S	Resolved from $R \vee S$ and $\sim R$.
15.	$\sim Q$	Resolved from $R \vee \sim Q$ and $\sim R$.
16.	Q	Resolved from $\sim R$ and $Q \vee R$.
17.	$\sim S$	Resolved from $\sim R$ and $R \vee \sim S$.
18.		Resolved $\sim R$ and R to $\sim R \vee R$, which is in turn null.

A contradiction is found when $\sim R$ is assumed as true. Hence, R is true.