

MEDICAL COST PERSONAL DATASETS



CONTENTS



01

INTRODUCTION

Data Overview

02

DATA VISUALIZATION

03

MODEL

Multivariate linear regression
K-nearest neighbors (k-NN)


04

HYPOTHESIS TESTING







INTRODUCTION

The dataset named Medical Cost Personal Dataset from **Kaggle**. This dataset has a large number of clients from insurance companies of the USA and multiple personal information for each client such as age, sex, body mass index(BMI), number of children, smoker/non-smoker, residential area. We are going to see the relations between the charges (The charges are an important point of estimation for any insurance company.) and some interesting attributes as well as some criteria these companies have.




About this file

This dataset consists of 1338 rows.

# age	sex	# bmi	# children	✓ smoker	region	# charges
Edad del asegurado	Género	Índice de masa corporal	Número de hijos	Indicador si fuma	Región donde vive el asegurado	Prima del seguro
	male 51% female 49%			<div> <div>true 0 0%</div> <div>false 0 0%</div> </div>	southeast 27% southwest 24% Other (649) 49%	
18	female	27.9	0	yes	southwest	16884.924
19	male	33.77	1	no	southeast	1725.5523
28	male	33	3	no	southeast	4449.462
33	male	22.705	0	no	northwest	21984.47061
32	male	28.88	0	no	northwest	3866.8552
31	female	25.74	0	no	southeast	3756.6216
46	female	33.44	1	no	southeast	8240.5096
37	female	27.74	3	no	northwest	7281.5056
37	male	29.83	2	no	northeast	6406.4107
60	female	25.84	0	no	northwest	28923.13692

02. DATA VISUALIZATION





Data Visualization

Response (dependent) variable: charges

```
summary(insurance$charges)
```


Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
1122	4740	9382	13270	16640	63770

Because the mean value is greater than the median, this implies that the distribution of insurance expenses is right-skewed




Data Visualization

Correlation between Charges and smoker




Correlation between Charges and Children




Data Visualization

Correlation between Charges and Gender




Correlation between Charges and region



Data Visualization

Variables is related to the amount of charges



03. MODEL

Multivariate linear regression
K-nearest neighbors (k-NN)



MULTIVARIATE LINEAR REGRESSION

charges ~ age + sex + bmi + children + smoker + region

```
> # lm all column  
> mul_model <- lm(charges ~ age + sex + bmi + children + smoker + region, data = insurance)  
> # mul_model_ <- lm(charges ~ ., data = insurance_new)  
> summary(mul_model)
```

Call:

```
lm(formula = charges ~ age + sex + bmi + children + smoker +  
    region, data = insurance)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-11304.9	-2848.1	-982.1	1393.9	29992.8

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-11938.5	987.8	-12.086	< 2e-16 ***
age	256.9	11.9	21.587	< 2e-16 ***
sexmale	-131.3	332.9	-0.394	0.693348
bmi	339.2	28.6	11.860	< 2e-16 ***
children	475.5	137.8	3.451	0.000577 ***
smoker	23848.5	413.1	57.723	< 2e-16 ***
regionnorthwest	-353.0	476.3	-0.741	0.458769
regionsoutheast	-1035.0	478.7	-2.162	0.030782 *
regionsouthwest	-960.0	477.9	-2.009	0.044765 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 6062 on 1329 degrees of freedom
Multiple R-squared: 0.7509, Adjusted R-squared: 0.7494
F-statistic: 500.8 on 8 and 1329 DF, p-value: < 2.2e-16

MULTIVARIATE LINEAR REGRESSION

charges ~ age + bmi + children + smoker

```
> # column that have significance  
> mul_model <- lm(charges ~ age + bmi + children + smoker, data = insurance)  
> # mul_model_ <- lm(charges ~ ., data = insurance_new)  
> summary(mul_model)
```

Call:
lm(formula = charges ~ age + bmi + children + smoker, data = insurance)

Residuals:

	Min	1Q	Median	3Q	Max
	-11897.9	-2920.8	-986.6	1392.2	29509.6

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	-12102.77	941.98	-12.848	< 2e-16	***
age	257.85	11.90	21.675	< 2e-16	***
bmi	321.85	27.38	11.756	< 2e-16	***
children	473.50	137.79	3.436	0.000608	***
smoker	23811.40	411.22	57.904	< 2e-16	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 6068 on 1333 degrees of freedom
Multiple R-squared: 0.7497, Adjusted R-squared: 0.7489
F-statistic: 998.1 on 4 and 1333 DF, p-value: < 2.2e-16

```
> sqrt(0.7489)  
[1] 0.8653901
```

MULTIVARIATE LINEAR REGRESSION

The effect of changing one predictor variable while controlling the values of the other predictor variables.

Multiple Linear Regression

$$y = b_0 + b_1x_1 + b_2x_2 + \dots + b_nx_n$$

Dependent variable (DV) Independent variables (IVs)

Constant Coefficients

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	-12102.77	941.98	-12.848	< 2e-16	***
age	257.85	11.90	21.675	< 2e-16	***
bmi	321.85	27.38	11.756	< 2e-16	***
children	473.50	137.79	3.436	0.000608	***
smoker	23811.40	411.22	57.904	< 2e-16	***

$$Y = -12102.77 + 257.85(\text{age}) + 321.85(\text{bmi}) + 473.50(\text{Children}) + 23811.40(\text{smoker})$$

```
insurance$Multi <- -12102.77 + 257.85*insurance$age + 321.85*insurance$bmi +  
473.50*insurance$children + 23811.40*insurance$smoker
```

```
insurance$prediction <- predict(mul_model, newdata = insurance)
```

	age	bmi	children	smoker	charges	Multi	prediction
1	19	27.900	0	1	16884.924	25587.3950	25587.4252
2	18	33.770	1	0	1725.552	3880.9045	3880.9459
3	28	33.000	3	0	4449.462	7158.5800	7158.6201
4	33	22.705	0	0	21984.471	3713.8843	3713.9005
5	32	28.880	0	0	3866.855	5443.4580	5443.4834

MULTIVARIATE LINEAR REGRESSION



K-NEAREST NEIGHBORS (K-NN)

sampling approach 2 variables: Smoker (n=274) and Non-Smoker (n=1064)

smoker	count	min	median	max	IQR
<chr>	<int>	<dbl>	<dbl>	<dbl>	<dbl>
yes	274	12829.	34456.	63770.	20193.
no	1064	1122.	7345.	36911.	7376.

Population (n=1338): set test 20% (n sample = 270) of smoker and non-smoker.
set train: 80% (n sample = 1068).

```
set.seed(612)
test_no <- sample_n(smoker_no, 135, fac = "ID")$ID
test_yes <- sample_n(smoker_yes, 135, fac = "ID")$ID
test <- c(test_no, test_yes)

# keep just the test data points/rows
all_test <- df[test, -1]
all_train <- df[-(test), -1]
```

K-NEAREST NEIGHBORS (K-NN)

Check the data structure

```
> str(all_train)
'data.frame': 1198 obs. of 7 variables:
 $ age      : int  18 28 32 31 46 37 37 60 25 62 ...
 $ sex      : chr   "male" "male" "male" "female" ...
 $ bmi      : num   33.8 33 28.9 25.7 33.4 ...
 $ children: int    1 3 0 0 1 3 2 0 0 0 ...
 $ smoker   : chr   "no" "no" "no" "no" ...
 $ region   : chr   "southeast" "southeast" "northwest"
 $ charges  : num   1726 4449 3867 3757 8241 ...
```

change the "smoker" variable to be of a factor type

```
# Change the "smoker" variable to be of a factor type
all_train$smoker <- as.factor(all_train$smoker)
all_test$smoker <- as.factor(all_test$smoker)
```

```
> str(all_train)
'data.frame': 1198 obs. of 7 variables:
 $ age      : int  18 28 32 31 46 37 37 60 25 62 ...
 $ sex      : chr   "male" "male" "male" "female" ...
 $ bmi      : num   33.8 33 28.9 25.7 33.4 ...
 $ children: int    1 3 0 0 1 3 2 0 0 0 ...
 $ smoker   : Factor w/ 2 levels "no","yes": 1 1 1 1 1
 $ region   : chr   "southeast" "southeast" "northwest"
 $ charges  : num   1726 4449 3867 3757 8241 ...
```

```
> str(all_test)
'data.frame': 140 obs. of 7 variables:
 $ age      : int  19 43 23 31 49 18 20 41 34 53 ...
 $ sex      : chr   "male" "male" "female" "male" ...
 $ bmi      : num   20.6 26 28.1 31.1 34.8 ...
 $ children: int    2 0 0 3 1 0 1 1 1 1 ...
 $ smoker   : chr   "no" "no" "no" "no" ...
 $ region   : chr   "northwest" "northeast" "northwest"
 $ charges  : num   2804 6837 2690 5425 9584 ...
```

```
> str(all_test)
'data.frame': 140 obs. of 7 variables:
 $ age      : int  19 43 23 31 49 18 20 41 34 53 ...
 $ sex      : chr   "male" "male" "female" "male" ...
 $ bmi      : num   20.6 26 28.1 31.1 34.8 ...
 $ children: int    2 0 0 3 1 0 1 1 1 1 ...
 $ smoker   : Factor w/ 2 levels "no","yes": 1 1 1 1 1
 $ region   : chr   "northwest" "northeast" "northwest"
 $ charges  : num   2804 6837 2690 5425 9584 ...
```


K-NEAREST NEIGHBORS (K-NN)

```
> # see the model's details
> model_knn
```

Call:

```
train.kknn(formula = smoker ~ ., data = all_train, kmax = 9)
```

Type of response variable: nominal

Minimal misclassification: 0.04307116

Best kernel: optimal

Best k: 1

```
> # Do a prediction on the test data
```

```
> prediction <- predict(model_knn, all_test[, -5])
```

```
> prediction
```

```
[1] no no no no no no no no no no no no no no no no no
[20] no no no no no no no yes no no no no yes no no no no no
[39] no no no no yes no no no no no no no no yes no no no no
[58] no no no no no no no no no no no no no no yes no no no
[77] no no no no no no no no no no no no no no no yes no no
[96] no no no no no no no no no no no no no no no no no yes
[115] no no no no no yes no no no no no no no no no no no no
[134] no no yes yes no yes yes yes yes yes yes yes yes yes yes
[153] yes yes yes yes yes yes yes no yes yes yes no yes no no yes
[172] yes no no yes yes yes yes yes no no yes no yes yes yes yes
[191] yes no yes yes yes yes yes yes yes yes yes yes no yes no yes
[210] yes no yes yes yes yes yes yes yes yes yes yes yes no yes yes
[229] yes yes yes yes yes yes yes yes yes yes yes no no yes yes yes
[248] yes yes yes yes yes no no no yes yes yes yes yes yes yes yes
[267] yes yes yes no
```

Levels: no yes

K-NEAREST NEIGHBORS (K-NN)

```
> #Display results
```

```
> solution
```

Confusion Matrix and Statistics

	Reference	
Prediction	no	yes
no	127	28
yes	8	107

Accuracy : 0.8667

95% CI : (0.8202, 0.9048)

No Information Rate : 0.5

P-Value [Acc > NIR] : < 2.2e-16

Kappa : 0.7333

McNemar's Test P-Value : 0.001542

Sensitivity : 0.9407

Specificity : 0.7926

Pos Pred Value : 0.8194

Neg Pred Value : 0.9304

Prevalence : 0.5000

Detection Rate : 0.4704

Detection Prevalence : 0.5741

Balanced Accuracy : 0.8667

'Positive' Class : no

	NO	YES
NO	127	8
YES	28	107


Accuracy = 0.867

Precision = 0.819

Sensitivity = 0.941

Specificity = 0.793

04. HYPOTHESIS TESTING



HYPOTHESIS TESTING

Exploratory data analysis has indicated that smoking has an effect on charges.

```
> # Read in our dataset
> df <- read.csv("insurance.csv")
> df %>%
+   group_by(smoker) %>%
+   summarise(
+     count = n(),
+     median = median(charges),
+     mean = mean(charges),
+     SD = sd(charges),
+     Var = var(charges)
+   ) %>%
+   arrange(desc(median))
# A tibble: 2 × 6
  smoker count median    mean    SD      Var
  <chr>   <int>   <dbl>   <dbl>   <dbl>   <dbl>
1 yes      274 34456. 32050. 11542. 133207311.
2 no     1064  7345.  8434.  5994.  35925420.
```

HYPOTHESIS TESTING

Step 1: Define null and alternative hypothesis

$H_0: \mu_1 - \mu_2 = 0$ The average charges of smokers is equal to non-smokers

$H_a: \mu_1 - \mu_2 > 0$ The average charges of smokers is greater than non-smokers

Test at the 5% level of significance: $\alpha = 0.05$



Smoker: yes

n: 274

Mean: 32050

SD: 11541.55

Variance: 133207311

Smoker: no

n: 1064

Mean: 8434

SD: 5993.782

Variance: 35925420



HYPOTHESIS TESTING


Step 2: Confidence Intervals Two-sample hypothesis

```
z.test(df$charges[df$smoker== "yes"], df$charges[df$smoker== "no"],  
       mu = 0, sigma.x = sd(df$charges[df$smoker== "yes"]),  
       sigma.y = sd(df$charges[df$smoker== "no"]), conf.level = 0.95)
```

Two-sample z-Test

```
data: df$charges[df$smoker == "yes"] and df$charges[df$smoker == "no"]  
z = 32.752, p-value < 2.2e-16  
alternative hypothesis: true difference in means is not equal to 0  
95 percent confidence interval:  
 22202.72 25029.21  
sample estimates:  
mean of x mean of y  
32050.232 8434.268
```

We are 95% confident that the difference in the population means lies in the interval [22202.72 , 25029.21]



HYPOTHESIS TESTING

Step 3: Since the samples are independent and both are large the test statistic is

Where D_0 = hypothesized difference between the means

Test Statistic

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - D_0}{\sigma_{(\bar{x}_1 - \bar{x}_2)}}$$

where

$$\sigma_{(\bar{x}_1 - \bar{x}_2)} = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Rejection region: $z < -z_\alpha$

[or $z > z_\alpha$ when $H_a: (\mu_1 - \mu_2) > D_0$]

Rejection region: $|z| > z_{\alpha/2}$


Step 4: Inserting the data into the formula for the test statistic gives

Test Statistic:

$$Z^* = \frac{(32050 - 8434) - 0}{\sqrt{\frac{133207311}{274} + \frac{35925420}{1064}}}$$
$$= \frac{23616}{\sqrt{486158.1 + 33764.49}} = 32.752$$

HYPOTHESIS TESTING

Step 5: Since the symbol in H_a is ">" this is a right-tailed test, so there is a single critical value, $z_{\alpha}=0.05$, which from the last line in Figure we read off as 2.576. The rejection region is $[2.576, \infty)$



Step 6: As shown in Figure "Rejection Region and Test Statistic for " the test statistic falls in the rejection region. We reject the null hypothesis and can conclude that people who smoke have on an average larger medical claim compared to people who don't smoke.

Step 7: Find P-Value

Test Statistic:

$$Z = \frac{(32050 - 8434) - 0}{\sqrt{\frac{133207311}{274} + \frac{35925420}{1064}}}$$
$$= \frac{23616}{\sqrt{486158.1 + 33764.49}} = 32.752$$

Step 8: The observed significance or p -value of the test is the area of the right tail of the standard normal distribution that is cut off by the test statistic $Z = 32.752$. The number 5.684 is too large to appear in Z-table the area of the *right* tail, is therefore $1 - 1.0000 = 0.0000$ (The actual value is approximately 0.00000000000000022 or $2.2e^{-16}$)

Two-sample z-Test

```
data: smoker_yes$charges and smoker_no$charges
z = 32.752, p-value < 2.2e-16
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 22202.72 25029.21
sample estimates:
mean of x mean of y
32050.232 8434.268
```

Step 9: Since $2.2e^{-16} < 0.05(2.576)$, p -value $< \alpha$ so the decision is to reject the null hypothesis : The data provide sufficient evidence, at the 5% level of significance, to conclude that the mean charges satisfaction for smoker is higher than for non-smoker.

THANKS

Kanyawee Sadubjit ID: 64130701719

Chonkeait Kanongsilp ID: 64130701721

