3.3 Graphs	plot and draw graphs with equation: $y = Ax^3 + Bx^2 + Cx + D$ in which: (i) the constants are integers and some could be zero (ii) the letters x and y can be replaced with any other two letters or: $y = Ax^3 + Bx^2 + Cx + D + E/x + F/x^2$ in which: (i) the constants are numerical and at least three of them are zero (ii) the letters x and y can be replaced with any other two letters	$y = x^3$, $y = 3x^3 - 2x^2 + 5x - 4$, $y = 2x^3 - 6x + 2$, $V = 60w(60 - w)$ $y = \frac{1}{x}, x \neq 0$, $y = 2x^2 + 3x + 1/x$, $x \neq 0$, $y = \frac{1}{x}(3x^2 - 5)$, $x \neq 0$, $W = \frac{5}{d^2}, d \neq 0$
	find the gradients of non-linear graphs	By drawing a tangent

	find the intersection points of two graphs, one linear (y_1) and one non-linear (y_2), and recognise that the solutions correspond to the solutions of $y_2 - y_1 = 0$	The x-values of the intersection of the two graphs: $y = 2x + 1$ $y = x^2 + 3x - 2$ are the solutions of: $x^2 + x - 3 = 0$ Similarly, the x-values of the intersection of the two graphs: $y = 5$ $y = x^3 - 3x^2 + 7$ are the solutions of: $x^3 - 3x^2 + 2 = 0$
	calculate the gradient of a straight line given the coordinates of two points	
	recognise that equations of the form $y = mx + c$ are straight line graphs with gradient m and intercept on the y axis at the point $(0, c)$	Find the equation of the straight line through $(1, 7)$ and $(2, 9)$
	find the equation of a straight line parallel to a given line	