

	significance level, test statistic, 1- tail test, 2-tail test, critical value, critical region, and acceptance region and p -value.	
8.3	Know that a sample is being used to make an inference about the population and appreciate the need for a random sample and of the necessary conditions.	Students should be aware that a conclusion to a hypothesis test should not be stated as definite.
8.4	Choose the appropriate hypothesis test to carry out in particular circumstances.	
8.5	Conduct a statistical hypothesis test for the proportion in the binomial distribution and interpret the results in context using exact probabilities or, where appropriate, a normal approximation.	In a hypothesis test on a population proportion, candidates may use either π or p as the parameter in their hypotheses.
8.6	Conduct a statistical hypothesis test for the mean of a normal distribution with known or assumed variance, from a large sample, and interpret the results in context.	
8.7	Know the importance of appropriate sampling when using hypothesis tests and be able to critique the conclusions drawn from rejecting or failing to reject a null hypothesis by considering the test performed.	

9 Contingency tables

Subject content	Additional information
9.1 Construct contingency tables from real data, combining data where appropriate, and interpret results in context.	
9.2 Use a χ^2 test with the appropriate number of degrees of freedom to test for independence in a contingency table and interpret the results of such a test.	Students will be expected to know how to find the number of degrees of freedom and that questions set will not require the use of Yates' correction.
9.3 Know that expected frequencies must be greater than, or equal to, 5 for a χ^2 test to be carried out and understand the requirement for combining classes if that is not the case.	Students must combine classes when expected frequencies are smaller than 5, and 'pooling' is sensible.