

PRACTICE PAPER

JEE Advanced**JEE WORK CUTS**

www.mtg.in | September 2021 | Pages 76 | ₹ 40

Class
XI-XII**CBSE** warm up **TERM-I**

MATHEMATICS


today

India's #1
MATHEMATICS MONTHLY
for JEE (Main & Advanced)



**OLYMPIAD
CORNER**

**MONTHLY
TEST DRIVE**
CLASS XI-XII



SOLVED PAPER
JEE MAIN SESSION-3

CONCEPT MAP
CLASS XI & XII

mtG

Trust of more than
1 Crore Readers
Since 1982



**BEAT THE
TIME TRAP**

You Ask ?
We Answer ✓

Embark on the
Avenues of Excellence

Admissions Open for 2021-22

Programs Offered:

BA, MA, B.Com, M.Com, B.Sc, M.Sc, BCA,
MCA, B.Ed, M.Ed across various disciplines

HOW SRMIST ACCELERATES YOUR FUTURE?

- Futuristic infrastructure with a conducive ambience for teaching and learning
- Hands-on training with advanced research equipment
- Syllabus based on Learning Outcome-based Curriculum Framework (LOCF) for skill development with in-built specialized placement training
- State-of-the-art Computer Labs with over 500 high-end systems equipped with the latest software
- In-house Coding Club to nurture students who are passionate about coding and learning new technologies
- International exposure through Semester Abroad Program (SAP)

PLACEMENTS THAT MAKE YOU GO PLACES

- Amid the pandemic, over 200 companies hired SRMIST students through the virtual hiring process
- The top recruiters include: Cognizant, Wipro, TCS, Infosys, Zifo R&D, McKinsey, Nielsen IQ, Amdocs, NatWest, L&T construction, City Union Bank, Intellepat, Deloitte, Cerner Corporation

- Highest Salary Package of ₹ 7 Lakh Per Annum



SRM
INSTITUTE OF SCIENCE & TECHNOLOGY
(Deemed to be University u/s 3 of UGC Act, 1956)

KATTANKULATHUR CAMPUS

APPLY NOW



Website: www.srmist.edu.in | Email: admissions.india@srmist.edu.in

Helpline: +91 44 6737 0700, +91 44 2745 5510, +91 44 4743 7500

MATHEMATICS today

Vol. XXXIX No. 9 September 2021

Corporate Office:

Plot 99, Sector 44 Institutional Area,
Gurugram - 122 003 (H.R.), Tel : 0124-6601200
e-mail : info@mtg.in website : www.mtg.in

Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital,
Ring Road, New Delhi - 110029.

Managing Editor : Mahabir Singh
Editor : Anil Ahlawat

CONTENTS

Competition Edge

Class XI

Class XII

- 4 JEE Main *Solved Paper Session 3*
- 14 JEE Advanced *Practice Paper*
- 25 Challenging Problems
- 31 Beat the Time Trap
- 34 Olympiad Corner
- 37 You Ask We Answer
- 40 JEE Work Outs
- 38 Concept Map *Sequences and Series*
- 50 CBSE Warm Up! *Complex Numbers and Quadratic Equations*
- 58 Monthly Test Drive *Trigonometric Functions*
- 39 Concept Map *Definite Integrals and Application of Integrals*
- 61 CBSE Warm Up! *Inverse Trigonometric Functions*
- 71 Monthly Test Drive *Matrices and Determinants*

14 PRACTICE PAPER 40
JEE Advanced JEE WORK CUTS Class 10-12 CBSE WARM UP TERM-I
50 61

India's #1 MATHEMATICS MONTHLY for JEE (Main & Advanced)

14 PRACTICE PAPER 40
JEE Advanced JEE WORK CUTS Class 10-12 CBSE WARM UP TERM-I
50 61

MATHEMATICS today

34 OLYMPIAD CORNER

4 SOLVED PAPER SESSION 3 JEE MAIN

MONTHLY TEST DRIVE 71 58

CONCEPT MAP CLASS 11 & 12

MTG Trust of more than 100 years Since 1921

31 BEAT THE TIME TRAP

37 You Ask ? We Answer ✓

Subscribe online at www.mtg.in

Individual Subscription Rates

	9 months	15 months	27 months
Mathematics Today	300	500	850
Chemistry Today	300	500	850
Physics For You	300	500	850
Biology Today	300	500	850

Combined Subscription Rates

	9 months	15 months	27 months
PCM	900	1400	2500
PCB	900	1400	2500
PCMB	1200	1900	3400

Send DD/M.O in favour of MTG Learning Media (P) Ltd.

Payments should be made directly to : MTG Learning Media (P) Ltd.

Plot 99, Sector 44 Institutional Area, Gurugram - 122 003, Haryana.

We have not appointed any subscription agent.

Printed and Published by Mahabir Singh on behalf of MTG Learning Media Pvt. Ltd. Printed at HT Media Ltd., B-2, Sector-63, Noida, UP-201307 and published at 406, Taj Apartment, Ring Road, Near Safdarjung Hospital, New Delhi - 110029.

Editor :

Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/InFocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.

Copyright © MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.

JEE MAIN 2021

SECTION-A (MULTIPLE CHOICE QUESTIONS)

1. Let $A = [a_{ij}]$ be a 3×3 matrix, where

$$a_{ij} = \begin{cases} 1, & \text{if } i=j \\ -x, & \text{if } |i-j|=1 \\ 2x+1, & \text{otherwise} \end{cases}$$

Let a function $f: R \rightarrow R$ be defined as $f(x) = \det(A)$. Then the sum of maximum and minimum values of f on R is equal to

- (a) $\frac{88}{27}$ (b) $\frac{20}{27}$ (c) $-\frac{88}{27}$ (d) $-\frac{20}{27}$

2. If z and ω are two complex numbers such that $|z\omega|=1$ and $\arg(z) - \arg(\omega) = \frac{3\pi}{2}$, then $\arg\left(\frac{1-2z\omega}{1+2z\omega}\right)$ is

- (a) $-\frac{\pi}{4}$ (b) $-\frac{3\pi}{4}$ (c) $\frac{3\pi}{4}$ (d) $\frac{\pi}{4}$

3. The value of the integral $\int_{-1}^1 \log_e(\sqrt{1-x} + \sqrt{1+x}) dx$ is equal to

- (a) $\frac{1}{2} \log_e 2 + \frac{\pi}{4}$ (b) $2 \log_e 2 + \frac{\pi}{4} - 1$
 (c) $\log_e 2 + \frac{\pi}{2} - 1$ (d) $2 \log_e 2 + \frac{\pi}{2} - \frac{1}{2}$

4. The mean of 6 distinct observations is 6.5 and their variance is 10.25. If 4 out of 6 observations are 2, 4, 5 and 7, then the remaining two observations are
 (a) 8, 13 (b) 1, 20 (c) 10, 11 (d) 3, 18

5. Let $y = y(x)$ be the solution of the differential equation $e^x \sqrt{1-y^2} dx + \left(\frac{y}{x}\right) dy = 0$, $y(1) = -1$. Then the value of $(y(3))^2$ is equal to

- (a) $1 + 4e^6$ (b) $1 - 4e^3$
 (c) $1 - 4e^6$ (d) $1 + 4e^3$

6. Let $y = y(x)$ be the solution of the differential equation

$$x \tan\left(\frac{y}{x}\right) dy = \left(y \tan\left(\frac{y}{x}\right) - x\right) dx, -1 \leq x \leq 1, y\left(\frac{1}{2}\right) = \frac{\pi}{6}.$$

Then the area of the region bounded by the curves $x = 0$, $x = \frac{1}{\sqrt{2}}$ and $y = y(x)$ in the upper half plane is

- (a) $\frac{1}{8}(\pi-1)$ (b) $\frac{1}{6}(\pi-1)$
 (c) $\frac{1}{12}(\pi-3)$ (d) $\frac{1}{4}(\pi-2)$

7. The number of real roots of the equation

$$\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{4}$$

- (a) 2 (b) 0 (c) 4 (d) 1

8. Let the tangent to the parabola $S: y^2 = 2x$ at the point $P(2, 2)$ meet the x -axis at Q and normal to it meet the parabola S at the point R . Then the area (in sq.units) of the triangle PQR is equal to

- (a) 25 (b) $\frac{15}{2}$ (c) $\frac{25}{2}$ (d) $\frac{35}{2}$

9. Let a function $f: R \rightarrow R$ be defined as

$$f(x) = \begin{cases} \sin x - e^x & \text{if } x \leq 0 \\ a + [-x] & \text{if } 0 < x < 1 \\ 2x - b & \text{if } x \geq 1 \end{cases}$$

where $[x]$ is the greatest integer less than or equal to x . If f is continuous on R , then $(a+b)$ is equal to

- (a) 3 (b) 4 (c) 5 (d) 2

10. Let ' a ' be a real number such that the function $f(x) = ax^2 + 6x - 15$, $x \in R$ is increasing in $\left(-\infty, \frac{3}{4}\right)$

and decreasing in $\left(\frac{3}{4}, \infty\right)$. Then the function $g(x) = ax^2 - 6x + 15$, $x \in R$ has a

- (a) local maximum at $x = \frac{3}{4}$
 (b) local minimum at $x = -\frac{3}{4}$
 (c) local maximum at $x = -\frac{3}{4}$
 (d) local minimum at $x = \frac{3}{4}$
11. Let $A = \begin{bmatrix} 2 & 3 \\ a & 0 \end{bmatrix}$, $a \in R$ be written as $P + Q$, where P is a symmetric matrix and Q is a skew symmetric matrix. If $\det(Q) = 9$, then the modulus of the sum of all possible values of determinant of P is equal to
 (a) 36 (b) 24 (c) 18 (d) 45
12. If in a triangle ABC , $AB = 5$ units, $\angle B = \cos^{-1}\left(\frac{3}{5}\right)$ and radius of circumcircle of ΔABC is 5 units, then the area (in sq. units) of ΔABC is
 (a) $8+2\sqrt{2}$ (b) $10+6\sqrt{2}$
 (c) $4+2\sqrt{3}$ (d) $6+8\sqrt{3}$
13. Let a be a positive real number such that
 $\int_0^a e^{x-[x]} dx = 10e - 9$, where $[x]$ is the greatest integer less than or equal to x . Then a is equal to
 (a) $10 + \log_e 2$ (b) $10 + \log_e 3$
 (c) $10 + \log_e(1 + e)$ (d) $10 - \log_e(1 + e)$
14. Let $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$ and $\vec{b} = \hat{i} + \hat{j}$. If \vec{c} is a vector such that $\vec{a} \cdot \vec{c} = |\vec{c}|, |\vec{c} - \vec{a}| = 2\sqrt{2}$ and the angle between $(\vec{a} \times \vec{b})$ and \vec{c} is $\frac{\pi}{6}$, then the value of $|(\vec{a} \times \vec{b}) \times \vec{c}|$ is
 (a) 3 (b) 4 (c) $\frac{3}{2}$ (d) $\frac{2}{3}$
15. The probability of selecting integers $a \in [-5, 30]$ such that $x^2 + 2(a+4)x - 5a + 64 > 0$, for all $x \in R$, is
 (a) $\frac{1}{4}$ (b) $\frac{2}{9}$
 (c) $\frac{7}{36}$ (d) $\frac{1}{6}$
16. Words with or without meaning are to be formed using all the letters of the word EXAMINATION. The probability that the letter M appears at the fourth position in any such word is
 (a) $\frac{2}{11}$ (b) $\frac{1}{11}$ (c) $\frac{1}{9}$ (d) $\frac{1}{66}$

17. The Boolean expression $(p \wedge \sim q) \Rightarrow (q \vee \sim p)$ is equivalent to
 (a) $\sim q \Rightarrow p$ (b) $p \Rightarrow \sim q$
 (c) $p \Rightarrow q$ (d) $q \Rightarrow p$
18. The coefficient of x^{256} in the expansion of $(1-x)^{101}(x^2+x+1)^{100}$ is
 (a) ${}^{100}C_{16}$ (b) ${}^{100}C_{15}$ (c) $-{}^{100}C_{16}$ (d) $-{}^{100}C_{15}$
19. Let $[x]$ denote the greatest integer $\leq x$, where $x \in R$. If the domain of the real valued function $f(x) = \sqrt{\frac{[x]-2}{[x]-3}}$ is $(-\infty, a) \cup [b, c] \cup [4, \infty)$, $a < b < c$, then the value of $a+b+c$ is
 (a) 8 (b) -2 (c) -3 (d) 1
20. If α and β are the distinct roots of the equation $x^2 + (3)^{1/4}x + 3^{1/2} = 0$, then the value of $\alpha^{96}(\alpha^{12}-1) + \beta^{96}(\beta^{12}-1)$ is equal to
 (a) 52×3^{24} (b) 56×3^{25}
 (c) 28×3^{25} (d) 56×3^{24}

PUZZLE CORNER



MATHDOKU

Introducing MATHDOKU, a mixture of ken-ken, sudoku and Mathematics. In this puzzle 6×6 grid is given, your objective is to fill the digits 1-6 so that each appear exactly once in each row and each column. Notice that most boxes are part of a cluster. In the upper-left corner of each multibox cluster is a value that is combined using a specified operation on its numbers. For example, if that value is 3 for a two-box cluster and operation is multiply, you know that only 1 and 3 can go in there. But it is your job to determine which number goes where! A few cluster may have just one box and that is the number that fills that box.

72x	4-		60x	
16+				
16+			18x	
9+	3+			2-
		36x		
	2-			3-

Readers can send their responses at editor@mtg.in or post us with complete address. Winners' name with their valuable feedback will be published in next issue.

SECTION-B (NUMERICAL VALUE TYPE)

Attempt any 5 questions out of 10.

21. If the shortest distance between the lines $\vec{r}_1 = \alpha\hat{i} + 2\hat{j} + 2\hat{k} + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$, $\lambda \in R$, $\alpha > 0$ and $\vec{r}_2 = -4\hat{i} - \hat{k} + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$, $\mu \in R$ is 9, then, α is equal to _____.
 22. Let a, b, c, d be in arithmetic progression with common difference λ . If $\begin{vmatrix} x+a-c & x+b & x+a \\ x-1 & x+c & x+b \\ x-b+d & x+d & x+c \end{vmatrix} = 2$, then value of λ^2 is equal to _____.
 23. Let $y = mx + c$, $m > 0$ be the focal chord of $y^2 = -64x$, which is tangent to $(x+10)^2 + y^2 = 4$. Then the value of $4\sqrt{2}(m+c)$ is equal to _____.
 24. There are 15 players in a cricket team, out of which 6 are bowlers, 7 are batsmen and 2 are wicket keepers. The number of ways, a team of 11 players be selected from them so as to include at least 4 bowlers, 5 batsmen and 1 wicket keeper, is _____.
 25. If the value of $\lim_{x \rightarrow 0} (2 - \cos x \sqrt{\cos 2x})^{\left(\frac{x+2}{x^2}\right)}$ is equal to e^a , then a is equal to _____.
 26. Let $\vec{a}, \vec{b}, \vec{c}$ be three mutually perpendicular vectors of the same magnitude and equally inclined at an angle θ , with the vector $\vec{a} + \vec{b} + \vec{c}$. Then $36\cos^2 2\theta$ is equal to _____.
 27. Let P be a plane passing through points $(1, 0, 1)$, $(1, -2, 1)$ and $(0, 1, -2)$. Let a vector $\vec{a} = \alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$ be such that \vec{a} is parallel to the plane P , perpendicular to $(\hat{i} + 2\hat{j} + 3\hat{k})$ and $\vec{a} \cdot (\hat{i} + \hat{j} + 2\hat{k}) = 2$, then $(\alpha - \beta + \gamma)^2$ equals _____.
 28. Let $A = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}$ and $B = 7A^{20} - 20A^7 + 2I$, where I is an identity matrix of order 3×3 . If $B = [B_{ij}]$, then b_{13} is equal to _____.
 29. Let T be the tangent to the ellipse $E: x^2 + 4y^2 = 5$ at the point $P(1, 1)$. If the area of the region bounded by tangent T , ellipse E , lines $x = 1$ and $x = \sqrt{5}$ is $\alpha\sqrt{5} + \beta + \gamma\cos^{-1}\left(\frac{1}{\sqrt{5}}\right)$, then $|\alpha + \beta + \gamma|$ is equal to _____.

30. The number of rational terms in the binomial expansion of $\left(\frac{1}{4^4} + 5^6\right)^{120}$ is _____.

SOLUTIONS

1. (c) : We have, $A = \begin{bmatrix} 1 & -x & 2x+1 \\ -x & 1 & -x \\ 2x+1 & -x & 1 \end{bmatrix}$

$\therefore f(x) = \det(A) = 4x^3 - 4x^2 - 4x$
 $\Rightarrow f'(x) = 12x^2 - 8x - 4 = 4(x-1)(3x+1)$

Now, $f'(x) = 0 \Rightarrow x = 1, -\frac{1}{3}$

Also, $f''(x) = 24x - 8$

At $x = 1, f''(x) = 16 > 0$

At $x = -\frac{1}{3}, f''(x) = -16 < 0$

\therefore Sum of maximum and minimum values of f
 $= f\left(\frac{-1}{3}\right) + f(1) = \frac{20}{27} - 4 = \frac{-88}{27}$

2. (b) : Let $z = r_1 e^{i\theta}$ and $\omega = r_2 e^{i\phi}$

$\Rightarrow \bar{z} = r_1 e^{-i\theta}$

Given $|z\omega| = 1 \Rightarrow |r_1 r_2 e^{i(\theta + \phi)}| = 1$

$\Rightarrow r_1 r_2 = 1$

... (i)

Given $\arg(z) - \arg(\omega) = \frac{3\pi}{2}$

$\Rightarrow \theta - \phi = \frac{3\pi}{2}$

... (ii)

Now, $\bar{z}\omega = r_1 e^{-i\theta} \cdot r_2 e^{i\phi}$

$= r_1 r_2 e^{-i(\theta - \phi)}$

$= \cos \frac{3\pi}{2} - i \sin \frac{3\pi}{2}$

[From (i) and (ii)]

$\Rightarrow \bar{z}\omega = i$

$\therefore \arg\left(\frac{1-2\bar{z}\omega}{1+3\bar{z}\omega}\right) = \arg\left(\frac{1-2i}{1+3i}\right)$

$= \arg\left(\frac{1-2i}{1-3i} \times \frac{1-3i}{1-3i}\right) = \arg\left(\frac{-1}{2}(1+i)\right)$

$= -\pi + \tan^{-1}(1)$

$[\because \arg(z) = -\pi + \tan^{-1}\frac{y}{x}, \text{ if } x < 0 \text{ and } y < 0]$

$= -\pi + \frac{\pi}{4} = -\frac{3\pi}{4}$

3. (c) : Let $I = 2 \int_0^1 \log_e(\sqrt{1-x} + \sqrt{1+x}) \cdot 1 dx$


$(\because \int_a^b f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(x) = f(-x))$

$= 2 \left[\left(\log_e(\sqrt{1-x} + \sqrt{1+x}) \cdot x \right)_0^1 \right]$

$- \int_0^1 x \left(\frac{1}{\sqrt{1-x} + \sqrt{1+x}} \right) \cdot \left(\frac{1}{2\sqrt{1+x}} - \frac{1}{2\sqrt{1-x}} \right) dx$



What is JEE (MAIN) like?



Visit
www.mtg.in
for latest offers
and to buy
online!

Thinking about competing in the JEE? Unsure of what it's like? Get a first-hand feel with MTG's JEE Main Explorer. With solved papers from the past 8 years, glance through the changing pattern of examination. Or attempt the papers to check your chances of success. Either way, at ₹ 700, isn't this a steal for JEE aspirants like you? So what are you waiting for? Order MTG's JEE Main Explorer today.

Available at all leading book shops throughout the country. To buy online visit www.mtg.in.
For more information or for help in placing your order, call 0124-6601200 or email: info@mtg.in



Scan now with your
smartphone or tablet
Application to read
QR codes required

$$= 2[(\log_e \sqrt{2} - 0)] - \int_0^1 \frac{x(\sqrt{1-x} - \sqrt{1+x})}{(\sqrt{1-x} + \sqrt{1+x})\sqrt{1-x^2}} dx$$

$$= \log_e 2 + \int_0^1 \left(\frac{1-\sqrt{1-x^2}}{\sqrt{1-x^2}} \right) dx$$

$$= \log_e 2 + (\sin^{-1} x - x)_0^1 = \log_e 2 + \frac{\pi}{2} - 1$$

4. (c) : Let the remaining two observations be a and b .

$$\text{Then, mean } = \frac{2+4+5+7+a+b}{6} = 6.5 \text{ (Given)}$$

$$\Rightarrow a+b = 21$$

Also, variance = 10.25 (Given)

$$\Rightarrow \frac{1}{6}(2^2 + 4^2 + 5^2 + 7^2 + a^2 + b^2) - (6.5)^2 = 10.25$$

$$\Rightarrow a^2 + b^2 = 315 - 94 = 221$$

$$\text{Now, } (a+b)^2 + (a-b)^2 = 2(a^2 + b^2)$$

$$\Rightarrow (a-b)^2 = 2 \times 221 - (21)^2 \quad [\text{From (i) and (ii)}]$$

$$\Rightarrow (a-b)^2 = 1 \Rightarrow a-b = \pm 1$$

If $a-b = 1$, then $a = 11, b = 10$

If $a-b = -1$, then $a = 10, b = 11$

$$\text{5. (c) : We have, } e^x \sqrt{1-y^2} dx + \left(\frac{y}{x} \right) dy = 0$$

$$\Rightarrow x e^x dx = \frac{-y}{\sqrt{1-y^2}} dy$$

Integrating both sides, we get

$$\int x e^x dx = - \int \frac{2y}{2\sqrt{1-y^2}} dy$$

$$\Rightarrow x e^x - \int e^x dx = \sqrt{1-y^2} + c$$

$$\Rightarrow x e^x - e^x = \sqrt{1-y^2} + c$$

$$\text{At } y(1) = -1, c = 0$$

$$\therefore e^x(x-1) = \sqrt{1-y^2}$$

$$\Rightarrow y^2 = 1 - (e^x(x-1))^2$$

$$\therefore (y(3))^2 = 1 - 4e^6$$

6. (a) : We gave,

$$x \tan\left(\frac{y}{x}\right) dy = \left(y \tan\left(\frac{y}{x}\right) - x \right) dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{x \left(\frac{y}{x} \tan\left(\frac{y}{x}\right) - 1 \right)}{x \tan\left(\frac{y}{x}\right)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{x} - \cot\left(\frac{y}{x}\right) \quad \dots(i)$$

$$\text{Putting } \frac{y}{x} = v \Rightarrow y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$\text{From (i), } v + x \frac{dv}{dx} = v - \cot(v) \Rightarrow x \frac{dv}{dx} = -\frac{1}{\tan v}$$

$$\Rightarrow \int (\tan v) dv = - \int \frac{1}{x} dx$$

$$\Rightarrow \log \left| \sec\left(\frac{y}{x}\right) \right| = -\log|x| + c \quad \dots(iii)$$

$$\text{At } y\left(\frac{1}{2}\right) = \frac{\pi}{6}, c = 0$$

$$\therefore \sec\left(\frac{y}{x}\right) = \frac{1}{x} \Rightarrow \cos\left(\frac{y}{x}\right) = x \Rightarrow y = x \cos^{-1} x$$

7. (b) : Required area

$$= \int_0^{1/\sqrt{2}} y dx = \int_0^{1/\sqrt{2}} (x \cos^{-1} x) dx$$

$$= \left[\left(2x^2 - 1 \right) \frac{\cos^{-1} x}{4} - \frac{x}{4} \sqrt{1-x^2} \right]_0^{1/\sqrt{2}} = -\frac{1}{8} + \frac{\pi}{8}$$

$$= \frac{1}{8}(\pi - 1)$$

7. (b) : We have,

$$\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{4} \quad \dots(i)$$

The equation (i) is defined for $x^2 + x \geq 0$

$$\Rightarrow x(x+1) \geq 0$$

$$\Rightarrow x \geq 0 \& x \leq -1 \text{ and } 0 \leq x^2 + x + 1 \leq 1$$

$$\Rightarrow -1 \leq x \leq 0$$

$\Rightarrow x = 0, -1$ is the only solution but it does not satisfy (i).

8. (c) : Equation of tangent to the parabola $y^2 = 2x$ at the point $P(2, 2)$ is given by $yy_1 = 2a(x + x_1)$

$$\therefore y(2) = 2 \times \left(\frac{1}{2} \right) (x+2)$$

$$\Rightarrow 2y = x + 2$$

Equation (i) meet the x -axis at $Q \therefore Q = (-2, 0)$

Equation of normal to the parabola $y^2 = 2x$ at point $P(2, 2)$ is given by

$$y - 2 = -\frac{2}{2\left(\frac{1}{2}\right)}(x-2) \Rightarrow y - 2 = -2(x-2)$$

$$\Rightarrow y = 6 - 2x$$

... (ii)



Equation (ii) meet the $y^2 = 2x$ at $R \therefore R\left(\frac{9}{2}, -3\right)$

$$\therefore \text{Area} (\Delta PQR) = \frac{1}{2} \begin{vmatrix} 2 & 2 & 1 \\ -2 & 0 & 1 \\ \frac{9}{2} & -3 & 1 \end{vmatrix} = \frac{25}{2} \text{ sq. units}$$

9. (a): We have, $f(x)$ is Continuous at $x = 0$.

$$\therefore \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^-} f(x)$$

$$\Rightarrow a - 1 = 0 - e^0$$

$$\Rightarrow a - 1 = -1 \Rightarrow a = 0$$

Also, $f(x)$ is continuous at $x = 1$.

$$\therefore \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^-} f(x)$$

$$\Rightarrow 2(1) - b = a + (-1)$$

$$\Rightarrow b = 2 - a + 1$$

$$\Rightarrow a + b = 3$$

10. (c): Since, $f(x) = ax^2 + 6x - 15$ is increasing in

$$\left(-\infty, \frac{3}{4}\right) \text{ and decreasing in } \left(\frac{3}{4}, \infty\right).$$

$$\therefore f(x) \text{ has a local maxima at } x = \frac{3}{4}.$$

Now, $f'(x) = 2ax + 6$

$$\Rightarrow f'\left(\frac{3}{4}\right) = 0$$

$$\Rightarrow 2a\left(\frac{3}{4}\right) + 6 = 0 \Rightarrow a = -4$$

$$\therefore g(x) = -4x^2 - 6x + 15 \Rightarrow g'(x) = -8x - 6$$

Now, $g'(x) = 0$

$$\Rightarrow -8x - 6 = 0 \Rightarrow x = -\frac{3}{4}$$

$$g''(x) = -8 < 0, \text{ at } x = -\frac{3}{4}.$$

$$\therefore g(x) \text{ has local maximum at } x = -\frac{3}{4}.$$

$$\boxed{11. (a):} \text{ We have, } A = \begin{bmatrix} 2 & 3 \\ a & 0 \end{bmatrix} \therefore A^T = \begin{bmatrix} 2 & a \\ 3 & 0 \end{bmatrix}$$

We can write A as, $A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T) = P + Q$

$$\therefore P = \begin{bmatrix} 2 & \frac{3+a}{2} \\ \frac{3+a}{2} & 0 \end{bmatrix} \text{ and } Q = \begin{bmatrix} 0 & \frac{3-a}{2} \\ \frac{-(3-a)}{2} & 0 \end{bmatrix}$$

$$\text{Now, } \det(Q) = 9 \Rightarrow 0 + \left(\frac{3-a}{2}\right)^2 = 9$$

$$\Rightarrow (3-a) = \pm 6 \Rightarrow a = -3, 9$$

$$\text{At } a = -3, P = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}, \det(P) = 0$$

$$\text{At } a = 9, P = \begin{bmatrix} 2 & 6 \\ 6 & 0 \end{bmatrix}, \det(P) = -36$$


$$\therefore \text{Required sum} = |0 + (-36)| = |-36| = 36$$

$$\boxed{12. (d):} \text{ Given, } \cos B = \frac{3}{5}$$

$$\therefore \sin B = \frac{4}{5}$$

We have, $R = 5$

$$\therefore \frac{c}{\sin C} = 2R$$



SAMURAI SUDOKU

ANSWER • AUGUST 2021



2	6	1	3	4	9	8	7	5
4	3	7	8	1	5	2	6	9
8	5	9	2	7	6	3	1	4
6	4	2	7	8	3	9	5	1
3	1	5	6	9	2	7	4	8
7	9	8	4	5	1	6	3	2
5	2	3	1	6	8	4	9	7
1	8	4	9	3	7	5	2	6
9	7	6	5	2	4	1	3	8
8	3	4	6	9	2	1	5	7
6	7	9	1	3	5	8	2	4
2	5	1	8	4	7	3	9	6
4	5	6	7	8	3	9	1	2
7	8	9	4	2	1	3	6	5
3	2	1	6	9	5	7	4	8
6	4	5	2	1	7	8	9	3
1	9	3	5	6	8	2	7	4
8	7	2	9	3	4	1	5	6
9	6	7	8	5	2	4	3	1
5	1	8	3	4	9	6	2	7
2	3	4	1	7	6	5	8	9

$$\therefore \sin C = \frac{5}{10} \Rightarrow C = 30^\circ$$

$$\text{Now, } \frac{b}{\sin B} = 2R \Rightarrow b = 2 \times 5 \times \frac{4}{5} = 8$$

Now, by cosine formula, we have

$$\begin{aligned}\cos B &= \frac{a^2 + c^2 - b^2}{2ac} \Rightarrow \frac{3}{5} = \frac{a^2 + 25 - 64}{2 \times a \times 5} \\ \Rightarrow a^2 - 6a - 3 &= 0 \\ \Rightarrow a &= \frac{6 \pm \sqrt{192}}{2} \Rightarrow a = 3 + 4\sqrt{3} \quad (\because a \neq 3 - 4\sqrt{3})\end{aligned}$$

$$\text{Now, Area } (\Delta ABC) = \frac{abc}{4R} = \frac{(3+4\sqrt{3})(8)(5)}{(4)(5)} = 2(3+4\sqrt{3}) \\ = (6+8\sqrt{3}) \text{ sq. units}$$

13. (a): a is positive integer i.e. $a > 0 \therefore a = [a] + \{a\}$, where $[\cdot]$ and $\{ \cdot \}$ denote the G.I.F and fractional part

Let $n \leq a < n+1, n \in W$

Here, $[a] = n$

$$\text{Now } \int_0^a e^{x-[x]} dx = 10e - 9 \\ \Rightarrow \int_0^n e^{x-[x]} dx + \int_n^a e^{x-[x]} dx = 10e - 9$$

$$\therefore n \int_0^1 e^x dx + \int_n^a e^{x-n} dx = 10e - 9$$

$$\Rightarrow n(e-1) + (e^{a-n} - 1) = 10e - 9$$

$$\therefore n = 10 \text{ and } \{a\} = \log_2 2$$

$$\text{So, } a = [a] + \{a\} = 10 + \log_2 2$$

14. (c): We have, $|\vec{a}| = 3$ and $\vec{a} \cdot \vec{c} = |\vec{c}|$

$$\text{Now, } |\vec{c} - \vec{a}| = 2\sqrt{2}$$

$$\Rightarrow |\vec{c}|^2 + |\vec{a}|^2 - 2\vec{c} \cdot \vec{a} = 8$$

$$\Rightarrow |\vec{c}|^2 + 9 - 2|\vec{c}| = 8$$

$$\Rightarrow |\vec{c}|^2 - 2|\vec{c}| + 1 = 0 \Rightarrow |\vec{c} - 1|^2 = 0 \Rightarrow |\vec{c}| = 1$$

$$\begin{array}{ccc} \hat{i} & \hat{j} & \hat{k} \\ \vec{a} \times \vec{b} & = & 2 & 1 & -2 \\ & & 1 & 1 & 0 \end{array} = 2\hat{i} - 2\hat{j} + \hat{k}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = 3$$

$$\therefore |(\vec{a} \times \vec{b}) \times \vec{c}| = |\vec{a} \times \vec{b}| |\vec{c}| \sin \frac{\pi}{6} = 3 \times 1 \times \frac{1}{2} = \frac{3}{2}$$

15. (b): We have, $x^2 + 2(a+4)x - 5a + 64 > 0 \forall x \in R$

$$\therefore D < 0$$

$$\Rightarrow 4(a+4)^2 - 4(-5a+64) < 0$$

$$\Rightarrow a^2 + 16 + 8a + 5a - 64 < 0$$

$$\Rightarrow a^2 + 13a - 48 < 0 \Rightarrow (a+16)(a-3) < 0$$

$$\therefore a \in (-16, 3)$$

16. (b): Possibilities for a are $\{-5, -4, \dots, 2\}$ i.e., 8 in number.
In set $[-5, 30]$, total integers = 36

$$\therefore \text{Required probability} = \frac{8}{36} = \frac{2}{9}$$

16. (b): In the word EXAMINATION, "A", N and I appears two times.

$$\therefore \text{Total number of words with M at fourth place} \\ = \frac{10!}{2! \cdot 2! \cdot 2!}$$

Also, total number of words formed using all letters of word EXAMINATION = $\frac{11!}{2! \cdot 2! \cdot 2!}$

$$\therefore \text{Required probability} = \frac{10!}{11!} = \frac{1}{11}$$

17. (c):

p	q	$\sim p$	$\sim q$	$p \wedge \sim q$	$q \vee \sim p$	$(p \wedge \sim q) \Rightarrow (q \vee \sim p)$	$p \Rightarrow q$
T	F	F	T	T	F	F	F
T	T	F	F	F	T	T	T
F	F	T	T	F	T	T	T
F	T	T	F	F	T	T	T

18. (b): We have, $(1-x)^{101} \cdot (x^2+x+1)^{100}$

$$= (1-x)^{100} \cdot (1-x) \cdot (x^2+x+1)^{100}$$

$$= ((1-x)(x^2+x+1))^{100} \cdot (1-x)$$

$$= (1-x^3)^{100} \cdot (1-x)$$

$$= (1-x^3)^{100} - x(1-x^3)^{100}$$

$$\therefore \text{Required coefficient} = (-1) \times (-{}^{100}C_{85}) \\ = {}^{100}C_{85} = {}^{100}C_{15}$$

19. (b): For domain, $\frac{|[x]|-2}{|[x]|-3} \geq 0$

Case I: When $|[x]| - 2 \geq 0$ and $|[x]| - 3 > 0$

$$\therefore x \in (-\infty, -3) \cup [4, \infty) \quad \dots(i)$$

Case II: When $|[x]| - 2 \leq 0$ and $|[x]| - 3 < 0$

$$\therefore x \in [-2, 3) \quad \dots(ii)$$

From (i) and (ii), we get

Domain of function = $(-\infty, -3) \cup [-2, 3) \cup [4, \infty)$

$$\therefore (a+b+c) = -3 + (-2) + 3 = -2$$

20. (a): As, $(\alpha^2 + \sqrt{3}) = -(3)^{1/4} \alpha$

$$\Rightarrow (\alpha^4 + 2\sqrt{3}\alpha^2 + 3) = \sqrt{3}\alpha^2$$

(On squaring both sides)

$$\therefore (\alpha^4 + 3) = -\sqrt{3}\alpha^2$$

$$\Rightarrow \alpha^8 + 6\alpha^4 + 9 = 3\alpha^4 \quad (\text{Again squaring both sides})$$

$$\Rightarrow \alpha^8 = -9 - 3\alpha^4$$


Multiplying by α^4 , we get

$$\alpha^{12} = -9\alpha^4 - 3\alpha^8$$


Study right. Dig deep.




Build a solid foundation for success
in JEE Main



₹ 875



₹ 875



₹ 875

Are you a do-it-yourself type of a student? Then for success in JEE Main, choose MTG's JEE Main combo, comprising coursebooks for Physics, Chemistry & Mathematics. This combo is all class 11 and 12 students need for a solid and deep understanding of concepts in these three key subjects.

FEATURES

- Based on latest pattern of JEE Main
- Covers the entire syllabus
- Full of graphic illustrations for deep understanding of concepts
- Levelwise MCQs with detailed solutions
- NCERT Drill MCQs framed from NCERT Books
- Previous 10 Years' MCQs (2018-2009) of JEE Main / AIEEE
- Numerical Value Type chapterwise questions included
- JEE Main January & September 2020 - January & April 2019 Solved Papers Included

Note: Coursebooks are also available separately.

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or e-mail: info@mtg.in



Scan now with your
smartphone or tablet
Application to read
QR codes required

$$\begin{aligned}\Rightarrow \alpha^{12} &= -9\alpha^4 - 3(-9 - 3\alpha^4) \\ \Rightarrow \alpha^{12} &= -9\alpha^4 + 27 + 9\alpha^4 \Rightarrow \alpha^{12} = 27 \\ \Rightarrow (\alpha^{12})^8 &= (27)^8 \\ \Rightarrow \alpha^{96} &= (3)^{24} \\ \text{Similarly, } \beta^{96} &= (3)^{24} \\ \therefore \alpha^{96}(\alpha^{12}-1) + \beta^{96}(\beta^{12}-1) &= 2 \times 26 \times 3^{24} \\ &= (3)^{24} \times 52\end{aligned}$$

21. (6): We know that, if $\vec{r}_1 = \vec{a} + \lambda \vec{b}$ and $\vec{r}_2 = \vec{c} + \lambda \vec{d}$, their shortest distance between two lines is given by

$$d = \frac{|\vec{b} \times \vec{d}| \cdot |\vec{a} - \vec{c}|}{|\vec{b} \times \vec{d}|}$$

$$\text{Now, } \vec{a} - \vec{c} = (\alpha+4)\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\text{Also, } \vec{b} \times \vec{d} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 2 \\ 3 & -2 & -2 \end{vmatrix} = 8\hat{i} + 8\hat{j} + 4\hat{k}$$

$$\therefore |\vec{b} \times \vec{d}| = 12$$

$$\text{Now, } \frac{\vec{b} \times \vec{d}}{|\vec{b} \times \vec{d}|} = \frac{2\hat{i} + 2\hat{j} + \hat{k}}{3}$$

Required distance

$$= \frac{((\alpha+4)\hat{i} + 2\hat{j} + 3\hat{k})(2\hat{i} + 2\hat{j} + \hat{k})}{3} = 9$$

$$\Rightarrow 2\alpha + 8 + 4 + 3 = 27 \Rightarrow 2\alpha = 27 - 15$$

$$\Rightarrow 2\alpha = 12 \Rightarrow \alpha = 6$$

$$\begin{array}{|ccc|} \hline & x+a-c & x+b & x+a \\ \hline & x-1 & x+c & x+b \\ & x-b+d & x+d & x+c \\ \hline \end{array}$$

Applying $c_2 \rightarrow c_2 - c_3$, we get

$$\begin{array}{|ccc|} \hline & x-2\lambda & \lambda & x+a \\ & x-1 & \lambda & x+b \\ & x+2\lambda & \lambda & x+c \\ \hline \end{array}$$

Applying $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$, we get

$$\begin{array}{|ccc|} \hline & x-2\lambda & \lambda & x+a \\ & 2\lambda-1 & 0 & \lambda \\ & 4\lambda & 0 & 2\lambda \\ \hline \end{array}$$

$$= -\lambda(4\lambda^2 - 2\lambda - 4\lambda^2) = 2\lambda^2 = 2 \text{ (Given)}$$

$$\therefore \lambda^2 = 1$$

23. (34): We have, $y^2 = -64x$, focus $(-16, 0)$.

Now, $y = mx + c$ is focal chord.

$$\Rightarrow c = 16m$$

Also, $y = mx + c$ is tangent to $(x+10)^2 + y^2 = 4$

$$\therefore y = m(x+10) \pm 2\sqrt{1+m^2}$$

$$\Rightarrow c = 10m \pm 2\sqrt{1+m^2}$$

From (i) and (ii), we get

$$16m = 10m \pm 2\sqrt{1+m^2}$$

$$\Rightarrow 6m = 2\sqrt{1+m^2}$$

$$\Rightarrow 9m^2 = 1 + m^2$$

$$\Rightarrow m = \frac{1}{2\sqrt{2}} \text{ and } c = \frac{8}{\sqrt{2}}$$

$$\therefore 4\sqrt{2}(m+c) = 4\sqrt{2} \left(\frac{17}{2\sqrt{2}} \right) = 34$$

24. (777): Total number of players = 15

Number of bowler = 6

Number of batsman = 7

Number of wicket keepers = 2

\therefore Total number of ways for with at least 4 bowlers, 5 batsman and 1 wicket keeper is to be selected

$$= {}^6C_4 {}^7C_5 {}^2C_2 + {}^6C_4 {}^6C_6 {}^2C_1 + {}^6C_5 {}^7C_5 {}^2C_1 \\ = 315 + 210 + 252 = 777$$

25. (3): We have, $\lim_{x \rightarrow 0} (2 - \cos x \sqrt{\cos 2x})^{\frac{x+2}{x^2}}$

$$= e^{\lim_{x \rightarrow 0} (1 - \cos x \sqrt{\cos 2x}) \frac{x+2}{x^2}} \quad (\text{Indeterminate form})$$

$$= e^{\lim_{x \rightarrow 0} \frac{(1 - \cos x \sqrt{\cos 2x})(x+2)}{x^2}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{(1 - \cos x \sqrt{\cos 2x})(1 + \cos x \sqrt{\cos 2x})}{(1 + \cos x \sqrt{\cos 2x})} \times \frac{x+2}{x^2}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{1 - \cos^2 x (\cos 2x)}{1 + \cos x \sqrt{\cos 2x}} \times \frac{x+2}{x^2}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{1 - \cos^2 x (1 - 2\sin^2 x)}{1 + \cos x \sqrt{\cos 2x}} \times \frac{x+2}{x^2}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{1 - (1 - \sin^2 x)(1 - 2\sin^2 x)}{x^2} \times \frac{x+2}{1 + \cos x \sqrt{\cos 2x}}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{3\sin^2 x - 2\sin^4 x}{x^2} \times \frac{x+2}{1 + \cos x \sqrt{\cos 2x}}}$$

$$= e^{\lim_{x \rightarrow 0} \frac{3\sin^2 x}{x^2} \left(1 - \frac{2}{3}\sin^2 x\right) \times \frac{x+2}{1 + \cos x \sqrt{\cos 2x}}}$$

$$= e^3 = e^a \text{ (Given)} \Rightarrow a = 3$$

26. (4): $|\vec{a} + \vec{b} + \vec{c}|^2$

$$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2|\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c}| = 3$$

$$\dots \text{(i)} \Rightarrow |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}$$

Now, $\vec{a}(\vec{a} + \vec{b} + \vec{c}) = |\vec{a}||\vec{a} + \vec{b} + \vec{c}| \cos \theta = 1 \cdot \sqrt{3} \cos \theta$

$$\Rightarrow 1 = \sqrt{3} \cos \theta$$

(ii)

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{3}} \therefore \cos^2 \theta = \frac{1}{3}$$

$$\therefore 36 \cos^2 2\theta = 36(2\cos^2 \theta - 1)^2 = 36\left(\frac{2}{3} - 1\right)^2 = 4$$

27. (81) : Equation of plane passing through $(1, 0, 1)$, $(1, -2, 1)$ and $(0, 1, -2)$ is given by

$$\begin{vmatrix} x-1 & y-0 & z-1 \\ 1-1 & -2-0 & 1-1 \\ 0-1 & 1-0 & -2-1 \end{vmatrix} = 0 \Rightarrow 3x-z-2=0$$

$\vec{a} = \hat{\alpha}\hat{i} + \hat{\beta}\hat{j} + \hat{\gamma}\hat{k}$ is parallel to $3x - z - 2 = 0$

$$\Rightarrow 3\alpha - \gamma = 0 \quad \dots(i)$$

\vec{a} is perpendicular to $\hat{i} + 2\hat{j} + 3\hat{k}$

$$\Rightarrow \alpha + 2\beta + 3\gamma = 0 \quad \dots(ii)$$

Also, $\vec{a} \cdot (\hat{i} + \hat{j} + 2\hat{k}) = 2$

$$\Rightarrow \alpha + \beta + 2\gamma = 2 \quad \dots(iii)$$

Solving (i), (ii) and (iii), we get

$$\alpha = 1, \beta = -5, \gamma = 3$$

$$\therefore (\alpha - \beta + \gamma)^2 = 9^2 = 81$$

28. (910) : Let $A = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} = I + P$,


where $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, P = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}$

$$P^2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, P^3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\text{Now, } B = 7A^{20} - 20A^7 + 2I \\ = 7(I + P)^{20} - 20(I + P)^7 + 2I \\ = 7(I + 20P + {}^{20}C_2P^2) - 20(I + 7P + {}^7C_2P^2) + 2I \\ \therefore b_{13} = 7 \times {}^{20}C_2 - 20 \times {}^7C_2 = 910$$

29. (1.25) : We have, $E : x^2 + 4y^2 = 5$

Equation of tangent to E at $P(1,1)$ is : $x + 4y = 5$



Required area

$$\begin{aligned} &= \int_1^{\sqrt{5}} \left(\frac{5-x}{4} - \frac{\sqrt{5-x^2}}{2} \right) dx \\ &= \left[\frac{5x}{4} - \frac{x^2}{8} - \frac{x}{4} \sqrt{5-x^2} - \frac{5}{4} \sin^{-1} \frac{x}{\sqrt{5}} \right]_1^{\sqrt{5}} \\ &= \frac{5}{4} \sqrt{5} - \frac{5}{4} - \frac{5}{4} \cos^{-1} \left(\frac{1}{\sqrt{5}} \right) \end{aligned}$$

On comparing with given area, we get

$$\alpha = \frac{5}{4}, \beta = -\frac{5}{4} \text{ and } \gamma = -\frac{5}{4}$$

$$\therefore |\alpha + \beta + \gamma| = \left| \frac{5}{4} - \frac{5}{4} - \frac{5}{4} \right| = 1.25$$

30. (21) : General term in the binomial expansion of

$$\left(\frac{1}{4^4} + \frac{1}{5^6} \right)^{120} = T_{r+1} = {}^{120}C_r (2^{1/2})^{120-r} (5)^{r/6}$$

For rational terms, $r = 6k$, $r = 120 - 2k$ and $0 \leq r \leq 120$

$$\Rightarrow r = 6k \text{ i.e., } r = 0, 6, 12, \dots, 120$$

So, total number of terms are 21.





ONLINE TEST SERIES

Practice Part Syllabus/ Full Syllabus
24 Mock Tests for

JEE Main



Now on your android smart phones with the same login of web portal.

Log on to test.pcmbtoday.com

JEE ADVANCED PRACTICE PAPER

Exam on 3rd October



PAPER-1

SINGLE OPTION CORRECT TYPE

1. Let S be the set of all complex numbers z satisfying $|z - 2 + i| \geq \sqrt{5}$. If the complex number z_0 is such that

$\frac{1}{|z_0 - 1|}$ is the maximum of the set $\left\{ \frac{1}{|z - 1|} : z \in S \right\}$,

then the principal argument of $\frac{4 - z_0 - \bar{z}_0}{z_0 - \bar{z}_0 + 2i}$ is

- (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{4}$ (c) $-\frac{\pi}{2}$ (d) $\frac{3\pi}{4}$

2. If $(1+k)\tan^2 x - 4\tan x - 1 + k = 0$ has real roots, then which one of the following is not true?

- (a) $k^2 \leq 5$ (b) $\tan(x_1 + x_2) = 2$
 (c) for $k=2$, $x_1 = \frac{\pi}{4}$ (d) none of these

3. If for non-zero x , $af(x) + bf\left(\frac{1}{x}\right) = \frac{1}{x} - 5$, where $a \neq b$, then $\int_1^2 f(x) dx =$

- (a) $\frac{1}{(a^2 + b^2)} \left[a \log 2 - 5a + \frac{7}{2}b \right]$
 (b) $\frac{1}{(a^2 - b^2)} \left[a \log 2 - 5a + \frac{7}{2}b \right]$
 (c) $\frac{1}{(a^2 - b^2)} \left[a \log 2 - 5a - \frac{7}{2}b \right]$
 (d) $\frac{1}{(a^2 + b^2)} \left[a \log 2 - 5a - \frac{7}{2}b \right]$

4. If $A(z_1)$, $B(z_2)$ and $C(z_3)$ are three points in the argand plane, where $|z_1 + z_2| = ||z_1| - |z_2||$ and

$|(1-i)z_1 + iz_3| = |z_1| + |z_3 - z_1|$, where $i = \sqrt{-1}$, then

- (a) A , B and C lie on a fixed circle with centre $\frac{(z_2 + z_3)}{2}$.
 (b) A , B and C are collinear points.

- (c) ABC form an equilateral triangle.
 (d) ABC form an obtuse angle triangle.

5. Given that $x_1 + x_2 + x_3 = 0$, $y_1 + y_2 + y_3 = 0$ and $x_1y_1 + x_2y_2 + x_3y_3 = 0$. Then

$$\frac{x_1^2}{x_1^2 + x_2^2 + x_3^2} + \frac{y_1^2}{y_1^2 + y_2^2 + y_3^2} =$$

- (a) $\frac{1}{3}$ (b) $\frac{2}{3}$
 (c) 1 (d) $\frac{4}{3}$

6. In ΔABC , $\sqrt{3} \sin C = 2 \sec A - \tan A$ and one of the sides is of length 2, then maximum area of the triangle ABC is

- (a) $2\sqrt{3}$ sq. units (b) $4\sqrt{3}$ sq. units
 (c) $\frac{\sqrt{3}}{2}$ sq. units (d) $\frac{\sqrt{3}}{4}$ sq. units

ONE OR MORE THAN ONE OPTION(S) CORRECT TYPE

7. Let $f: R \rightarrow R$ be given by

$$f(x) = \begin{cases} x^5 + 5x^4 + 10x^3 + 10x^2 + 3x + 1, & x < 0; \\ x^2 - x + 1, & 0 \leq x < 1; \\ (2/3)x^3 - 4x^2 + 7x - (8/3), & 1 \leq x < 3; \\ (x-2)\log_e(x-2) - x + (10/3), & x \geq 3. \end{cases}$$

Then which of the following options is/are correct?

- (a) f' is not differentiable at $x = 1$
 (b) f is increasing on $(-\infty, 0)$
 (c) f is onto
 (d) f' has a local maximum at $x = 1$

8. The roots of the equation, $(x^2 + 1)^2 = x(4x^2 + 5x + 4)$ are both real and imaginary of the form $A \pm \sqrt{B}$ and $C \pm iD$ respectively, then

(a) $D = \frac{\sqrt{3}}{2}$

(b) $A = \frac{5}{2}$

(c) $B = \frac{21}{4}$

(d) $C = -\frac{1}{2}$

9. Bag A contains 2 white and 3 red balls and bag B contains 4 white and 7 red balls, one bag is selected randomly and a ball is drawn from the bag then

(a) If drawn ball is found to be red, the probability that it was drawn from bag B is $\frac{35}{68}$

(b) If drawn ball is found to be white, the probability that it was drawn from bag A is $\frac{11}{21}$

(c) If drawn ball is found to be red, the probability that it was drawn from bag A is $\frac{33}{68}$

(d) If drawn ball is found to be white, the probability that it was drawn from bag B is $\frac{10}{21}$

10. The possible value of the expression

$$\frac{\sec^4 \alpha + \sec^4 \beta}{\tan^2 \beta - \tan^2 \alpha} \quad \left(\alpha, \beta \neq \frac{n\pi}{2}, n \in I \right)$$
 is/are

- (a) 6 (b) 8 (c) 10 (d) 12

11. Let f be a non negative function defined on the interval $[0, 1]$. If $\int_0^x \sqrt{1-(f'(t))^2} dt = \int_0^x f(t) dt$, $0 \leq x \leq 1$ and $f(0) = 0$, then

(a) $f\left(\frac{1}{2}\right) < \frac{1}{2}$ and $f\left(\frac{1}{3}\right) > \frac{1}{3}$

(b) $f\left(\frac{1}{2}\right) > \frac{1}{2}$ and $f\left(\frac{1}{3}\right) > \frac{1}{3}$

(c) $f\left(\frac{1}{2}\right) < \frac{1}{2}$ and $f\left(\frac{1}{3}\right) < \frac{1}{3}$

(d) $f\left(\frac{1}{2}\right) > 0$ and $f\left(\frac{1}{3}\right) < \frac{1}{3}$

12. The point P is the intersection of the straight line joining the points $Q(2, 3, 5)$ and $R(1, -1, 4)$ with the plane $5x - 4y - z = 1$. If S is the foot of the perpendicular drawn from the point $T(2, 1, 4)$ to QR , then the possible length of the line segment PS is/are

(a) $\frac{1}{\sqrt{2}}$

(b) $\sqrt{2}$

(c) 2

(d) $2\sqrt{2}$

NUMERICAL VALUE TYPE

13. If the foci of the ellipse $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$ and the hyperbola $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$ coincide, then the value of b^2 is _____.

14. If the vectors $\overline{AB} = 3\hat{i} + 4\hat{k}$ and $\overline{AC} = 5\hat{i} - 2\hat{j} + 4\hat{k}$ are the sides of a triangle ABC , then the length of the median through A is _____.

15. $\lim_{x \rightarrow 0} \frac{(1 - \cos 2x)(3 + \cos x)}{x \tan 4x}$ is equal to _____.

16. If for some natural number n , $(1 + 2 + 3 + \dots + n) + k = 2013$, where k is one of the numbers 1, 2, 3, ..., n then $n + k$ is _____.

17. In a certain test there are n questions. In this test 2^{n-1} students gave wrong answers to atleast i questions, where $i = 1, 2, 3, \dots, n$. If the total number of wrong answers given is 2047, then n is _____.

18. Tangents are drawn to the circle $x^2 + y^2 = 12$ at the points where it is met by the circle $x^2 + y^2 - 5x + 3y - 2 = 0$, then the x -coordinate of the point of intersection of these tangents is _____.

PAPER-2

SINGLE DIGIT INTEGER ANSWER TYPE

1. If the papers of 5 students can be checked by any one of the 5 teachers. If the probability that all the 5 papers are checked by exactly 2 teachers is m then the value of $\frac{125m}{2}$ equal to _____.

2. Let $P = [a_{ij}]$ be a 3×3 matrix and let $Q = [b_{ij}]$, where $b_{ij} = 2^{i+j} a_{ij}$ for $1 \leq i, j \leq 3$. If the determinant of P is 2,

then the determinant of the matrix Q is 2^{k+4} , where k is equal to _____.

3. If $(3 + x^{2008} + x^{2009})^{2010} = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$,

then the value of $a_0 - \frac{1}{2}a_1 - \frac{1}{2}a_2 + a_3 - \frac{1}{2}a_4 - \frac{1}{2}a_5 + a_6 - \dots - K^{1005}$, where K is equal to _____.

4. Let $u(x)$ and $v(x)$ be differentiable functions such that $\frac{u(x)}{v(x)} = 7$. If $\frac{u'(x)}{v'(x)} = p$ and $\left(\frac{u(x)}{v(x)}\right)' = q$, then $\frac{p+q}{p-q}$ has the value equal to _____. (u' is equal to derivative of ' u ')

5. Let \vec{a} and \vec{b} be two unit vectors such that $\vec{a} \cdot \vec{b} = 0$. For some $x, y \in R$, let $\vec{c} = x\vec{a} + y\vec{b} + (\vec{a} \times \vec{b})$. If $|\vec{c}| = 2$ and the vector \vec{c} is inclined at some angle α to both \vec{a} and \vec{b} , then the value of $8 \cos^2 \alpha$ is _____.

6. If $f(x+y) = f(x)f(y)$ and $f(x) = 1 + xg(x)H(x)$ where $\lim_{x \rightarrow 0} g(x) = 2$, $\lim_{x \rightarrow 0} H(x) = 3$, then $f'(x) = Kf(x)$ where $K = _____$.

ONE OR MORE THAN ONE OPTION(S) CORRECT TYPE

7. Let $f(x) = \frac{x}{1+x^2}$ and $g(x) = \frac{e^{-x}}{1+[x]}$, where $[\cdot]$ is the greatest integer less than or equal to x , then
 (a) Domain $(f+g) = R - [-2, 0]$
 (b) Domain $(f-g) = R - [-1, 0)$
 (c) Range $f \cap$ Range $g = \left[-2, \frac{1}{2}\right]$
 (d) Range $g = R - \{0\}$

8. Consider the function $f(x) = \sin^5 x + \cos^5 x - 1$, $x \in \left[0, \frac{\pi}{2}\right]$. Which of the following is/are correct?

- (a) $f(x)$ is monotonic increasing in $\left(0, \frac{\pi}{4}\right)$
 (b) $f(x)$ is monotonic decreasing in $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$.
 (c) \exists some $c \in \left(0, \frac{\pi}{2}\right)$ for which $f'(c) = 0$.
 (d) The equation $f(x) = 0$ has two roots in $\left[0, \frac{\pi}{2}\right]$.

9. Let $I_n = \int_0^{\sqrt{3}} \frac{dx}{1+x^n}$ ($n = 1, 2, 3, \dots$) and $\lim_{n \rightarrow \infty} I_n = I_0$ (say), then which of the following statement(s) is/are correct?

- (a) $I_1 > I_0$ (b) $I_2 < I_0$
 (c) $I_0 + I_1 + I_2 > 3$ (d) $I_0 + I_1 > 2$

10. A line L passing through the point $P(1, 4, 3)$, is perpendicular to both the lines $\frac{x-1}{2} = \frac{y+3}{1} = \frac{z-2}{4}$ and $\frac{x+2}{3} = \frac{y-4}{2} = \frac{z+1}{-2}$.

If the position vector of point Q on L is (a_1, a_2, a_3) such that $(PQ)^2 = 357$, then $(a_1 + a_2 + a_3)$ can be

- (a) 16 (b) 15 (c) 2 (d) 1

11. In a triangle PQR , let $\angle PQR = 30^\circ$ and the sides PQ and QR have lengths $10\sqrt{3}$ and 10, respectively. Then, which of the following statement(s) is (are) TRUE?

- (a) $\angle QPR = 45^\circ$
 (b) The area of the triangle PQR is $25\sqrt{3}$ and $\angle QRP = 120^\circ$
 (c) The radius of the incircle of the triangle PQR is $10\sqrt{3} - 15$
 (d) The area of the circumcircle of the triangle PQR is 100π

12. Triangles $A_1A_2A_3$ and $B_1B_2B_3$ have side lengths (a_1, a_2, a_3) and (b_1, b_2, b_3) respectively satisfying the relation $\sqrt{a_1 + a_2 + a_3} \sqrt{b_1 + b_2 + b_3} = \sqrt{a_1 b_1} + \sqrt{a_2 b_2} + \sqrt{a_3 b_3}$, then which one of the following statements is/are true ?

- (a) $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$ (b) $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$
 (c) $\Delta A_1A_2A_3$ and $\Delta B_1B_2B_3$ are similar
 (d) $\Delta A_1A_2A_3$ and $\Delta B_1B_2B_3$ are congruent

NUMERICAL VALUE TYPE

13. If α, β are roots of the equation $p(x^2 - x) + x + 5 = 0$ and p_1, p_2 are two values of p for which the roots α, β are connected by the relation $\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{4}{5}$, then the value of $\frac{p_1}{p_2} + \frac{p_2}{p_1} = _____$.

14. The number of real solutions of the equation

$$\sin^{-1} \left(\sum_{i=1}^{\infty} x^{i+1} - x \sum_{i=1}^{\infty} \left(\frac{x}{2} \right)^i \right) = \frac{\pi}{2} - \cos^{-1} \left(\sum_{i=1}^{\infty} \left(-\frac{x}{2} \right)^i - \sum_{i=1}^{\infty} (-x)^i \right)$$

lying in the interval $\left(-\frac{1}{2}, \frac{1}{2}\right)$ is _____.

(Here, the inverse trigonometric functions $\sin^{-1} x$ and $\cos^{-1} x$ assume values in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and $[0, \pi]$, respectively.)

15. Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11, ..., and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23, Then, the number of elements in the set $X \cup Y$ is _____.



16. For each positive integer n ,

$$\text{Let } y_n = \frac{1}{n}((n+1)(n+2) \dots (n+n))^n$$

For $x \in R$, let $[x]$ be the greatest integer less than or equal to x . If $\lim_{n \rightarrow \infty} y_n = L$, then the value of $[L]$ is ____.

17. $\lim_{x \rightarrow 0} \frac{\tan(e^x)x^4 - \tan(-e^x)x^4}{\sin^4 x} = \text{_____}$, [where $[\cdot]$ is G.I.E.]

18. Let $A = \begin{bmatrix} 1 & \frac{3}{2} \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & -3 \\ -2 & 2 \end{bmatrix}$

and $C_r = \begin{bmatrix} r \cdot 3^r & 2^r \\ 0 & (r-1)3^r \end{bmatrix}$


be given matrices. If $\sum_{r=1}^{50} \text{tr}((AB)^r C_r) = 3 + a \cdot 3^b$ where $\text{tr}(A)$ denotes trace of matrix A , then find the value of $\frac{1}{20}(a+b)$. [Where a and b are relatively prime]

SOLUTIONS

PAPER-1

1. (c) : $\frac{1}{|z-1|}$, $z \in S$, attains

its maximum when $|z-1|$ attains its minimum.



Also, the curve of $|z-2+i| \geq \sqrt{5}$ looks like the given figure.

$$\text{Consider } \frac{4-z_0-\bar{z}_0}{z_0-\bar{z}_0+2i} = \frac{4-2\operatorname{Re} z_0}{2i\operatorname{Im} z_0+2i} = \frac{2(2-\operatorname{Re} z_0)}{2i(\operatorname{Im} z_0+1)}$$

$$= -i\lambda, \text{ where } \lambda > 0$$

Since $2 - \operatorname{Re} z_0 > 0$ and $\operatorname{Im} z_0 > 0$

$$\therefore \text{The principal argument of } \frac{4-z_0-\bar{z}_0}{z_0-\bar{z}_0+2i} \text{ is } -\pi/2.$$

2. (d) : Let $\tan x_1$ and $\tan x_2$ are the roots of the equation $(1+k)\tan^2 x - 4\tan x + (k-1) = 0$.

$$\therefore \tan x_1 + \tan x_2 = \frac{4}{1+k} \text{ and } \tan x_1 \cdot \tan x_2 = \frac{k-1}{1+k}$$

$$\Rightarrow \tan(x_1 + x_2) = \frac{\tan x_1 + \tan x_2}{1 - \tan x_1 \tan x_2} = \frac{\frac{4}{1+k}}{1 - \left(\frac{k-1}{1+k}\right)} = 2$$

Since, the equation has real roots.

$$\therefore D \geq 0 \Rightarrow 16 - 4(k^2 - 1) \geq 0 \Rightarrow k^2 \leq 5$$

3. (b) : $af(x) + bf\left(\frac{1}{x}\right) = \frac{1}{x} - 5 \text{ (for each } x \neq 0\text{)} \quad \dots \text{(i)}$

Replacing x by $\frac{1}{x}$ in (i), we get

$$af\left(\frac{1}{x}\right) + bf(x) = x - 5 \quad \dots \text{(ii)}$$

Eliminating $f\left(\frac{1}{x}\right)$ from (i) and (ii), we get

$$(a^2 - b^2)f(x) = \frac{a}{x} - bx - 5a + 5b$$

$$\Rightarrow (a^2 - b^2) \int_1^x f(t) dt = \left[\left(a \log|x| - \frac{b}{2}x^2 - 5(a-b)x \right) \right]_1^x$$

SAMURAI SUDOKU



Samurai Sudoku puzzle consists of five overlapping sudoku grids. The standard sudoku rules apply to each 9×9 grid. Place digits from 1 to 9 in each empty cell. Every row, every column and every 3×3 box should contain one of each digit.

The puzzle has a unique answer.

2	6	7	9	8				
5					3			4
1	3					5		
					4	7		
					6	3		
2	6	5	1	8	2			
2	4	8	6		7			6
5					3	4	7	
					6			

7		5		2				
9	1							
6	3	4	9	2				
2			3	9				
			7					
					2			
2	6	5	1	8	2			
5					7			
					3	4	7	

7		4	6		8	9	6	5	2
3		9		8	3	9	7	4	8
	9	5		1		2			7
2			8	4		8			7 4
4		6					6	8	
		3		5			1		
3			1	7			4		7
		7	4	1			3		5 2
5	1			8		5	7	1	

Readers can send their responses at editor@mtg.in or post us with complete address. Winners' name with their valuable feedback will be published in next issue.

$$= a \log 2 - 2b - 10(a-b) - a \log 1 + \frac{b}{2} + 5(a-b)$$

$$= a \log 2 - 5a + \frac{7}{2}b$$

$$\Rightarrow \int_1^2 f(x)dx = \frac{1}{a^2 - b^2} \left[a \log 2 - 5a + \frac{7}{2}b \right]$$

4. (a) : $\because |z_1 + z_2| = ||z_1| - |z_2||$

$$\Rightarrow \arg\left(\frac{z_1}{z_2}\right) = \pm \pi$$

$$\text{and } |z_1 + i(z_3 - z_1)|$$

$$= |z_1| + |z_3 - z_1|$$

$$\Rightarrow \arg\left(\frac{z_1}{z_3 - z_1}\right) = \frac{\pi}{2}$$

$$\therefore \text{Centre of circle is } \left(\frac{z_2 + z_3}{2}\right)$$

and ΔABC is right angled triangle with $\angle A = 90^\circ$

5. (b) : Consider three vectors

$$\vec{n}_1 = (x_1, x_2, x_3), \vec{n}_2 = (y_1, y_2, y_3) \text{ and } \vec{n}_3 = (1, 1, 1).$$

From the given data,

$$\vec{n}_1 \cdot \vec{n}_2 = 0, \vec{n}_2 \cdot \vec{n}_3 = 0 \text{ and } \vec{n}_1 \cdot \vec{n}_3 = 0$$

i.e., \vec{n}_1, \vec{n}_2 and \vec{n}_3 are mutually \perp vectors.

Now, $\frac{x_1^2}{x_1^2 + x_2^2 + x_3^2}, \frac{y_1^2}{y_1^2 + y_2^2 + y_3^2}$ and $\frac{1}{3}$ are the squares of the projections of the vector $(1, 0, 0)$ on to the vectors of $\vec{n}_1, \vec{n}_2, \vec{n}_3$ respectively and hence their sum = 1

$$\text{i.e., } \frac{x_1^2}{x_1^2 + x_2^2 + x_3^2} + \frac{y_1^2}{y_1^2 + y_2^2 + y_3^2} + \frac{1}{3} = 1$$

6. (a) : On simplifying, we have

$$\sqrt{3} \cos A \sin C = 2 - \sin A$$

$$\Rightarrow \sqrt{3} \cos A \sin C + \sin A = 2$$


As $\sin C \leq 1$ and maximum value of $\sqrt{3} \cos A + \sin A = 2$

So, $\sin C = 1$, i.e., $C = 90^\circ, A = 30^\circ, B = 60^\circ$

So, maximum area = $\frac{1}{2} \cdot 2 \cdot 2\sqrt{3} = 2\sqrt{3}$ sq. units

7. (a, c, d) : We have,

$$f(x) = \begin{cases} (x+1)^5 - 2x, & x < 0; \\ x^2 - x + 1, & 0 \leq x < 1; \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3; \\ (x-2) \log_e(x-2) - x + \frac{10}{3}, & x \geq 3. \end{cases}$$



$$\text{Now, } f'(x) = \begin{cases} 5(x+1)^4 - 2, & x < 0 \\ 2x-1, & 0 < x < 1 \\ 2x^2 - 8x + 7, & 1 < x < 3 \\ \log_e(x-2), & x > 3 \end{cases}$$

$$\text{and } f''(x) = \begin{cases} 2, & 0 < x < 1 \\ 4x-8, & 1 < x < 3 \end{cases}$$

$$\text{Clearly } \lim_{x \rightarrow 1^-} f''(x) \neq \lim_{x \rightarrow 1^+} f''(x)$$

and $f''(x)$ changes sign from positive to negative.

$\therefore f'(x)$ has a local maxima at $x = 1$ and $f'(x)$ is not differentiable at $x = 1$.

$\therefore f'(-1) < 0 \therefore f$ is not increasing on $(-\infty, 0)$

$$\text{For } x < 0, \lim_{x \rightarrow -\infty} f(x) = -\infty, \lim_{x \rightarrow 0^-} f(x) = 1$$

\therefore The range of f in $(-\infty, 0)$ is $(-\infty, 1)$.

For $3 < x < \infty, f'(x) > 0 \therefore f$ is increasing and $f(3) = \frac{1}{3}$.

So, the range of f contains $\left(\frac{1}{3}, \infty\right)$.

Hence, the range of f is whole of R .

8. (a, b, c, d) : Given equation is

$$(x^2 + 1)^2 = x(4x^2 + 5x + 4) \quad \dots(i)$$

$$\Rightarrow x^4 - 4x^3 - 3x^2 - 4x + 1 = 0$$

$$\Rightarrow x^2 \left(x^2 - 4x - 3 - \frac{4}{x} + \frac{1}{x^2} \right) = 0$$

$$\Rightarrow \left(x^2 + \frac{1}{x^2} \right) - 4 \left(x + \frac{1}{x} \right) - 3 = 0$$

(as $x = 0$ does not satisfy (i))

$$\Rightarrow t^2 - 4t - 5 = 0, \text{ where } t = x + \frac{1}{x}$$

$$\Rightarrow (t+1)(t-5) = 0$$

$$\Rightarrow \left(x + \frac{1}{x} + 1 \right) \left(x + \frac{1}{x} - 5 \right) = 0$$

$$\Rightarrow (x^2 + x + 1)(x^2 - 5x + 1) = 0$$

$$\Rightarrow x^2 + x + 1 = 0, \quad x^2 - 5x + 1 = 0$$


$$\Rightarrow x = \frac{-1 \pm i\sqrt{3}}{2}, x = \frac{5 \pm \sqrt{21}}{2}$$

$$x = -\frac{1}{2} \pm i\frac{\sqrt{3}}{2} = C \pm iD, x = \frac{5}{2} \pm \sqrt{\frac{21}{4}} = A \pm \sqrt{B}$$

$$\Rightarrow C = \frac{-1}{2}, D = \frac{\sqrt{3}}{2}, \quad A = \frac{5}{2}, B = \frac{21}{4}$$

Precision Revision for JEE is Here!

mtG



₹625

- FEATURES:**
- 20 years solved papers with detailed solutions
 - 10 Model Test Papers
 - Chapter-wise indexing of questions

Now, create your own pre-JEE. Just like pre-boards. With previous years' papers and model test papers for JEE Advanced, complete with detailed solutions, identify your areas of weakness and work on addressing them in time. Multiple test papers ensure you do your dry runs again and again, till such time you feel confident of taking on the best. For it will indeed be the best you compete with in JEE Advanced. So what are you waiting for? **Order MTG's JEE Advanced Explorer today.**

Available at all leading book shops throughout India. To buy online visit www.mtg.in.

For more information or for help in placing your order, call 0124-6601200 or email:info@mtg.in



Scan now with your
smartphone or tablet
Application to read
QR codes required

9. (a, b, c, d) : Let E_1 and E_2 be the events of selecting bag A and B, respectively. Then,

$$P(E_1) = P(E_2) = \frac{1}{2},$$

$$P\left(\frac{\text{red}}{E_1}\right) = \frac{3}{5}, P\left(\frac{\text{red}}{E_2}\right) = \frac{7}{11}, P\left(\frac{\text{white}}{E_1}\right) = \frac{2}{5}$$

$$\text{and } P\left(\frac{\text{white}}{E_2}\right) = \frac{4}{11}$$

$$\therefore P\left(\frac{E_2}{\text{red}}\right) = \frac{P(E_2)P\left(\frac{\text{red}}{E_2}\right)}{P(E_1)P\left(\frac{\text{red}}{E_1}\right) + P(E_2)P\left(\frac{\text{red}}{E_2}\right)}$$

$$= \frac{\frac{1}{2} \cdot \frac{7}{11}}{\frac{1}{2} \cdot \frac{3}{5} + \frac{1}{2} \cdot \frac{7}{11}} = \frac{35}{68}$$

$$\therefore P\left(\frac{E_1}{\text{red}}\right) = 1 - P\left(\frac{E_2}{\text{red}}\right) = 1 - \frac{35}{68} = \frac{33}{68}$$

$$\text{Now, } P\left(\frac{E_1}{\text{white}}\right) = \frac{P(E_1)P\left(\frac{\text{white}}{E_1}\right)}{P(E_1)P\left(\frac{\text{white}}{E_1}\right) + P(E_2)P\left(\frac{\text{white}}{E_2}\right)}$$

$$= \frac{\frac{2}{5}}{\frac{2}{5} + \frac{4}{11}} = \frac{11}{21}$$

$$\therefore P\left(\frac{E_2}{\text{white}}\right) = 1 - P\left(\frac{E_1}{\text{white}}\right) = 1 - \frac{11}{21} = \frac{10}{21}$$

10. (b, c, d) : Let $\tan^2 \alpha = a$ and $\tan^2 \beta = b$, then the given expression reduces to

$$\frac{(a+1)^2}{b} + \frac{(b+1)^2}{a} = 2\left(\frac{a+b}{b+a}\right) + \left(\frac{a^2}{b} + \frac{b^2}{a} + \frac{1}{a} + \frac{1}{b}\right)$$

$$\geq 2(2) + 4(1) = 8$$

$$\text{11. (c) : } \int_0^x \sqrt{1-(f'(t))^2} dt = \int_0^x f(t) dt$$

Differentiating, we get $\sqrt{1-(f'(x))^2} = f(x)$

$$\Rightarrow 1 - (f'(x))^2 = (f(x))^2 \Rightarrow f'(x) = \sqrt{1-(f(x))^2}$$


$$\Rightarrow \int \frac{dt}{\sqrt{1-t^2}} = \int dx \Rightarrow \sin^{-1}(f(x)) = x + c$$

$$\Rightarrow f(x) = \sin(x+c), f(0) = 0 \Rightarrow c = 0 \therefore f(x) = \sin x$$

$$\text{Now, } f\left(\frac{1}{2}\right) = \sin \frac{1}{2} < \frac{1}{2}, f\left(\frac{1}{3}\right) = \sin \frac{1}{3} < \frac{1}{3}.$$

12. (a) : Clearly, the equation of line QR be

$$\frac{x-2}{1} = \frac{y-3}{4} = \frac{z-5}{1}$$



Let the point P be given as

$$P(2+t, 3+4t, 5+t)$$

As P lies in the plane $5x - 4y - z = 1$

$$\therefore 5(2+t) - 4(3+4t) - (5+t) = 1$$

$$\Rightarrow 10 + 5t - 12 - 16t - 5 - t = 1 \Rightarrow -12t - 7 = 1$$

$$\Rightarrow 12t = -8 \therefore t = -\frac{8}{12} = -\frac{2}{3}$$

The co-ordinates of P are $\left(\frac{4}{3}, \frac{1}{3}, \frac{13}{3}\right)$

Again let $S \equiv (2+m, 3+4m, 5+m)$, m being a constant. As S is the foot of perpendicular, we have

$$1(2+m-2) + 4(3+4m-1) + 1(5+m-4) = 0$$

$$\Rightarrow m + 4(4m+2) + m + 1 = 0$$

$$\Rightarrow 18m + 9 = 0 \therefore m = -\frac{1}{2}$$

$$\text{Thus } S \equiv \left(\frac{3}{2}, 1, \frac{9}{2}\right)$$

Now, by distance formula

$$PS = \sqrt{\left(\frac{3}{2} - \frac{4}{3}\right)^2 + \left(1 - \frac{1}{3}\right)^2 + \left(\frac{9}{2} - \frac{13}{3}\right)^2}$$

$$= \sqrt{\left(\frac{1}{6}\right)^2 + \left(\frac{2}{3}\right)^2 + \left(\frac{1}{6}\right)^2} = \sqrt{\frac{1}{36} + \frac{4}{9} + \frac{1}{36}} = \sqrt{\frac{9}{18}} = \frac{1}{\sqrt{2}}$$

$$\text{13. (7) : Hyperbola is } \frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$$

$$\therefore a = \sqrt{\frac{144}{25}}, b = \sqrt{\frac{81}{25}} \text{ and } e_1 = \sqrt{1 + \frac{81}{144}} = \sqrt{\frac{225}{144}} = \frac{15}{12} = \frac{5}{4}$$

$$\text{Now, foci} = (\pm ae_1, 0) = \left(\pm \frac{12}{5} \cdot \frac{5}{4}, 0\right) = (\pm 3, 0)$$

Also, focus of ellipse = (4e, 0).

$$\Rightarrow e = \frac{3}{4}. \text{ Hence, } b^2 = 16 \left(1 - \frac{9}{16}\right) = 7$$

14. (33) : $\overline{AM} = \frac{1}{2}(\overline{AB} + \overline{AC})$

$$= \frac{1}{2}[(3, 0, 4) + (5, -2, 4)]$$

$$= \frac{1}{2}(8, -2, 8) = (4, -1, 4)$$

$$\therefore |\overline{AM}| = \sqrt{4^2 + 1^2 + 4^2} = \sqrt{33}$$

15. (2) : $\lim_{x \rightarrow 0} \frac{(1-\cos 2x)(3+\cos x)}{x(\tan 4x)}$

$$= \lim_{x \rightarrow 0} \frac{1-\cos 2x}{x^2} \cdot \left(\frac{\tan 4x}{x} \right) (3+\cos x)$$

$$= \lim_{x \rightarrow 0} \left(\frac{2\sin^2 x}{x^2} \right) \cdot \left(\frac{x}{\tan 4x} \right) (3+\cos x) = 2 \times \frac{1}{4} \times 4 = 2$$

16. (122) : $\frac{n(n+1)}{2} + k = 2013$ and $n \geq k$

$$\text{i.e. } n \geq 2013 - \frac{n(n+1)}{2}$$

Solving, $n \in (61.9, 62.9)$

So, $n = 62$ (integral) and $k = 60$

17. (11) : Number of students who gave wrong answers to atleast i questions = $2^n - i$

Number of students who gave wrong answers to atleast n questions = $2^n - 1$

Number of students gave wrong answers to exactly i questions = $2^{n-i} - 2^n - (i+1)$

Number of wrong answers

$$= \sum_{i=1}^{n-1} i \left(2^{n-i} - 2^{n-(i+1)} \right) + (n) = 2047$$

$$\Rightarrow 1 + 2^1 + 2^2 + \dots + 2^{n-1} = 2047$$

$$\Rightarrow 2^n = 2048 \Rightarrow n = 11$$


18. (6) : The given circles are $x^2 + y^2 = 12$ and $x^2 + y^2 - 5x + 3y - 2 = 0$

Common chord say, AB is $5x - 3y - 10 = 0$. Let the coordinates of point of intersection of tangents be $P(\alpha, \beta)$.

Equation of chord of contact of two tangents, drawn $P(\alpha, \beta)$, with respect to $x^2 + y^2 = 12$ is $x\alpha + y\beta - 12 = 0$. Comparing the coefficients of common chord AB and

$$\text{chord of contact, we get } \frac{\alpha}{5} = \frac{\beta}{-3} = \frac{-12}{-10} \Rightarrow \alpha = 6$$

$\therefore x$ -coordinate is 6.



PAPER-2

1. (6) : Here, $n(S) =$ The number of ways in which papers of 5 students can be checked by the 5 teachers = 5^5

and $n(A) =$ choosing two teachers out of 5 \times the number of ways in which 5 papers can be checked by exactly 2 teachers = ${}^5C_2 \times (2^5 - 2) = 300$

\therefore Required probability

$$= \frac{n(A)}{n(S)} = \frac{300}{5^5} = \frac{12}{125} = m \quad (\text{given})$$

$$\text{Hence, } \frac{125m}{2} = 6$$

$$2. (9) : \text{We have, } Q = \begin{vmatrix} 2^2 a_{11} & 2^3 a_{12} & 2^4 a_{13} \\ 2^3 a_{21} & 2^4 a_{22} & 2^5 a_{23} \\ 2^4 a_{31} & 2^5 a_{32} & 2^6 a_{33} \end{vmatrix}$$

Your favourite MTG Books/Magazines available in MAHARASHTRA at

Shridhar Granth Bhardar - Ahmed Nagar Ph: 0241-2347307, 9922664979

Vithal Book Seller - Akola Ph: 0724-2439403, 2431003, 9370336111

Rathi Paper Traders - Akola Ph: 00724-2452797, 9767655760, 9422160329

Jalaram Agency - Amravati Ph: 957267, 9372452996, 7385903365

Aarti Book Centre - Aurangabad Ph: 0240-2329250, 2333366, 4125679, 942203836

Maya Book Centre - Aurangabad Ph: 0240-2360150, 9421983933, 9421688529

Saraswati Book Centre - Aurangabad Ph: 9823162604, 7776009085

Kashik Book Stall - Baramati Ph: 224640, 9764510999

Naveen Book Stall - Dhule Ph: 2562-232574, 234574, 9422296434

Sharda Books Incorporation - Jaigad Ph: 0257-223765, 2228567, 9850950666

Destry World Stationers - Kolhapur Ph: 0231-202129, 9415588033

Adash Book Centre - Latur Ph: 9049547705, 7030184786

Krishna Pustakalaya - Latur Ph: 02385-259783, 9423297710, 9423291399

Anjali Trading Company - Mumbai Ph: 022-26714024, 25, 9819737977

Modern Book Plaza - Mumbai Ph: 3254940, 28801233, 022-28801233, 8018391969

Vidyarthi Sales Agencies - Mumbai Ph: 23829330, 23867279, 23571416, 9819776110

Student Agencies Pvt. Ltd. - Mumbai Ph: 022-40496161, 40496131, 982124775

JMD Book Distributors - Nagpur Ph: 0712-2555213, 2557838, 9823073969

Newly Book Depot - Nagpur Ph: 0712-2555213, 2557838, 9823073969

Renuka Book Distributor - Nagpur Ph: 0712-2726122, 9785406133

India Book Agencies - Nanded Ph: 9890489460

Morya Books & Genera Stores - Nanded Ph: 9730868713, 7385571813

Mahalaxmi Pustakalaya - Nashik Ph: 0253-2578538, 9422253538, 9869013220

Deepak Book Store - New Farve Ph: 7765426858, 9869013220

Sri Nivas Books & Gen Store - Prabhan Ph: 9420529899, 9970600851

Goyal Book Distributors - Pune Ph: 020-24459470, 9970600943

Pragat Book Centre - Pune Ph: 020-66022707, 9850039311

Vikas Book House - Pune Ph: 020-24468737, 24476376, 9860286472, 9921331187

Career Books - Sangli - 0233-2324205, 9370142979, 9881007001

Harshal Book Agency - Solapur Ph: 9923875597, 9764249414

Jyoti Book Centre - Thane Ph: 022-25413409, 25413504, 9833944432

Unique Traders - Wardha Ph: 9960578652, 7152243617

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

$$= 2^9 \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ 2a_{21} & 2a_{22} & 2a_{23} \\ 2^2 a_{31} & 2^2 a_{32} & 2^2 a_{33} \end{vmatrix} = 2^9 \cdot 2 \cdot 2^2 \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= 2^{12} \det P = 2^{12} \cdot 2 = 2^{13} = 2^{k+4} \Rightarrow k = 9$$

3. (4) : Putting $x = \omega, \omega^2$, we get

$$(3 + \omega + \omega^2)^{2010} = a_0 + a_1\omega + a_2\omega^2 + \dots$$

$$\Rightarrow 2^{2010} = a_0 + a_1\omega + a_2\omega^2 + a_3 + a_4\omega + \dots \quad \dots(i)$$

$$\text{and } 2^{2010} = a_0 + a_1\omega^2 + a_2\omega + a_3 + a_4\omega^2 + \dots \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2^{2011} = 2a_0 - a_1 - a_2 + 2a_3 - a_4 - a_5 + 2a_6 - \dots$$

$$\therefore a_0 - \frac{1}{2}a_1 - \frac{1}{2}a_2 + a_3 - \frac{1}{2}a_4 - \frac{1}{2}a_5 + a_6 - \dots = 2^{2010}$$

$$= (2^2)^{1005} = k^{1005} \text{ (Given)} \Rightarrow k = 4$$

4. (1) : $u(x) = 7v(x) \Rightarrow u'(x) = 7v'(x) \Rightarrow p = 7$ (given)

$$\text{Again } \frac{u(x)}{v(x)} = 7 \Rightarrow \left(\frac{u(x)}{v(x)} \right)' = 0 \Rightarrow q = 0.$$

$$\text{Now, } \frac{p+q}{p-q} = \frac{7+0}{7-0} = 1$$

5. (3) : Here $\vec{c} = x\vec{a} + y\vec{b} + \vec{a} \times \vec{b}$

$$\text{Also, } \vec{a} \cdot \vec{c} = \vec{b} \cdot \vec{c} = 2 \cos \alpha$$

$$\text{We have, } \vec{c} \cdot \vec{a} = x\vec{a} \cdot \vec{a} + 0 + 0$$

$$\therefore x = \vec{c} \cdot \vec{a} = 2 \cos \alpha. \text{ Similarly, } y = 2 \cos \alpha$$

$$\text{Now, as } c^2 = x^2 + y^2 + 1$$

$$\therefore \text{ We have } 4 = 2(4 \cos^2 \alpha) + 1 \Rightarrow 3 = 8 \cos^2 \alpha$$

6. (6) : We know, $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$= \lim_{h \rightarrow 0} \frac{f(x)f(h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x)[1 + h g(h)H(h) - 1]}{h}$$

$$= f(x) \lim_{h \rightarrow 0} g(h) \cdot H(h) = 6f(x)$$

7. (b, d) : Domain of $f = R$, Domain $g = R - [-1, 0]$;

because $-1 \leq x < 0 \Rightarrow 1 + [x] = 0$

\therefore Domain of $f - g = R - [-1, 0)$

Since $e^{-x} > 0 \Rightarrow (1 + [x])y > 0$

It can be observed that $y > 0 \Rightarrow 1 + [x] > 0$,

or $y < 0 \Rightarrow 1 + [x] < 0$

\therefore Range $g = R - \{0\}$

8. (c, d) : We have, $f'(x) = 5 \sin^4 x \cos x - 5 \cos^4 x \sin x = 5 \sin x \cos x (\sin x - \cos x)(1 + \sin x \cos x)$

$$\therefore f'(x) = 0 \text{ at } x = \frac{\pi}{4}. \text{ Also } f(0) = f\left(\frac{\pi}{2}\right) = 0$$

Hence \exists some $c \in \left(0, \frac{\pi}{2}\right)$ for which $f'(c) = 0$

(By Rolle's Theorem)

Also in $\left(0, \frac{\pi}{4}\right)$, $f(x)$ is decreasing and in $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$, $f(x)$ is increasing.

$$\Rightarrow \text{ Minimum at } x = \frac{\pi}{4}$$

$$\text{As } f(0) = f\left(\frac{\pi}{2}\right) = 0 \Rightarrow 2 \text{ roots exists.}$$

9. (a, c, d) : $I_1 = \ln(1 + \sqrt{3})$

$$I_2 = \frac{\pi}{3}$$

$$I_0 = \lim_{n \rightarrow \infty} I_n = \lim_{n \rightarrow \infty} \left[\int_0^1 \frac{dx}{1+x^n} + \underbrace{\int_{\frac{1}{1+x^n}}^{\frac{\sqrt{3}}{1+x^n}} \frac{dx}{1+x^n}}_{\text{zero}} \right] = \int_0^1 dx = 1.$$

Hence $I_0 = 1$.

10. (b, d) : Equation of the line passing through $P(1, 4, 3)$ and direction ratio's $a, b & c$ is

$$\frac{x-1}{a} = \frac{y-4}{b} = \frac{z-3}{c} \quad \dots(i)$$

$$\text{Since (i) is perpendicular to } \frac{x-1}{2} = \frac{y+3}{1} = \frac{z-2}{4}$$

$$\text{and } \frac{x+2}{3} = \frac{y-4}{2} = \frac{z+1}{-2}$$

Hence $2a + b + 4c = 0$ and $3a + 2b - 2c = 0$

$$\therefore \frac{a}{-2-8} = \frac{b}{12+4} = \frac{c}{4-3} \Rightarrow \frac{a}{-10} = \frac{b}{16} = \frac{c}{1}$$

Hence the equation of the line is

$$\frac{x-1}{-10} = \frac{y-4}{16} = \frac{z-3}{1} \quad \dots(ii)$$

Now any point Q on (ii) can be taken as $(1 - 10\lambda, 16\lambda + 4, \lambda + 3)$

$$\therefore \text{ Distance of } Q \text{ from } P(1, 4, 3) = (10\lambda)^2 + (16\lambda)^2 + \lambda^2 = 357$$

$$\Rightarrow (100 + 256 + 1)\lambda^2 = 357$$

$$\Rightarrow \lambda = 1 \text{ or } -1$$

$$\therefore Q \text{ is } (-9, 20, 4) \text{ or } (11, -12, 2)$$

Hence, $a_1 + a_2 + a_3 = 15$ or 1

11. (b, c, d) : Using cosine rule,

$$\cos 30^\circ = \frac{\sqrt{3}}{2} = \frac{(10\sqrt{3})^2 + 10^2 - PR^2}{2 \cdot 10 \cdot 10\sqrt{3}}$$

$$\Rightarrow \frac{\sqrt{3}}{2} \cdot 2 \times 10\sqrt{3} \times 10 = 400 - PR^2$$

$$\Rightarrow 300 = 400 - PR^2 \Rightarrow PR^2 = 100 \quad \therefore PR = 10$$

So, $QR = PR \quad \therefore \angle QPR = 30^\circ$

$$ar(\Delta PQR) = \frac{1}{2} \cdot 10\sqrt{3} \cdot 10 \sin 30^\circ = \frac{1}{2} \cdot 10 \cdot 10\sqrt{3} \cdot \frac{1}{2} = 25\sqrt{3}$$

Also, $\angle QRP = 180^\circ - 2 \times 30^\circ = 120^\circ$

$$\text{Now, } r = \frac{\Delta}{s} = \frac{25\sqrt{3}}{\frac{10+10+10\sqrt{3}}{2}} = \frac{25\sqrt{3}}{10+5\sqrt{3}} = \frac{25\sqrt{3}}{5(2+\sqrt{3})}$$

$$= 5\sqrt{3}(2-\sqrt{3}) = 10\sqrt{3}-15$$

$$\text{Also, } R = \frac{10}{2 \sin 30^\circ} = 10$$

Area of circumcircle = $\pi R^2 = 100\pi$

12. (a, c) : We have, $\sqrt{(a_1+a_2+a_3)(b_1+b_2+b_3)}$
 $= \sqrt{a_1 b_1} + \sqrt{a_2 b_2} + \sqrt{a_3 b_3}$

$$\text{Let } a_1 = P_1^2, a_2 = P_2^2, a_3 = P_3^2$$

$$\text{and } b_1 = Q_1^2, b_2 = Q_2^2, b_3 = Q_3^2$$

$$\Rightarrow (P_1^2 + P_2^2 + P_3^2)(Q_1^2 + Q_2^2 + Q_3^2) = (P_1 Q_1 + P_2 Q_2 + P_3 Q_3)^2$$

$$\Rightarrow (P_1 Q_2 - P_2 Q_1)^2 + (P_2 Q_3 - P_3 Q_2)^2 + (P_3 Q_1 - P_1 Q_3)^2 = 0$$

$$\Rightarrow \frac{P_1}{Q_1} = \frac{P_2}{Q_2} = \frac{P_3}{Q_3} = \lambda$$

$$\text{or } \frac{\sqrt{a_1}}{\sqrt{b_1}} = \frac{\sqrt{a_2}}{\sqrt{b_2}} = \frac{\sqrt{a_3}}{\sqrt{b_3}} \quad \therefore \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

13. (254) : Here the given equation is

$$p(x^2 - x) + x + 5 = 0 \Rightarrow px^2 - (p-1)x + 5 = 0$$

$$\therefore \alpha + \beta = \frac{p-1}{p} \text{ and } \alpha\beta = \frac{5}{p}$$

$$\text{Now } \frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{4}{5} \Rightarrow \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{4}{5}$$

$$\Rightarrow \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} = \frac{4}{5} \Rightarrow \frac{(p-1)^2 - 10p}{5p} = \frac{4}{5}$$

$$\Rightarrow p^2 - 16p + 1 = 0 \Rightarrow p_1 + p_2 = 16 \text{ and } p_1 p_2 = 1$$

$$\text{Now, } \frac{p_1}{p_2} + \frac{p_2}{p_1} = \frac{(p_1 + p_2)^2 - 2p_1 p_2}{p_1 p_2} = \frac{256 - 2}{1} = 254$$

14. (2) : Let $f(x) = \sum_{i=1}^{\infty} x^{i+1} - x \sum_{i=1}^{\infty} \left(\frac{x}{2}\right)^i$

$$= (x^2 + x^3 + \dots \text{ to } \infty) - x \left(\frac{x}{2} + \left(\frac{x}{2}\right)^2 + \dots \text{ to } \infty \right)$$

$$= \frac{x^2}{1-x} - \frac{x \cdot \frac{x}{2}}{1-\frac{x}{2}} = \frac{x^2}{1-x} - \frac{x^2}{2-x}$$

Again, let $g(x) = \sum_{i=1}^{\infty} \left(-\frac{x}{2}\right)^i - \sum_{i=1}^{\infty} (-x)^i$



$$= \left[\left(-\frac{x}{2}\right) + \left(\frac{-x}{2}\right)^2 + \dots \text{ to } \infty \right] - \left[(-x) + (-x)^2 + \dots \text{ to } \infty \right]$$



COMIC CAPSULE

BUT I DON'T KNOW
HOW TO DANCE.

I CAN LEAD.



$$= \frac{-x}{\frac{x}{1+x} - \frac{x}{2+x}} = \frac{x}{1+x} - \frac{x}{2+x}$$

The given equation becomes

$$\sin^{-1}f(x) + \cos^{-1}g(x) = \pi/2$$

So we must have $f(x) = g(x)$

$$\Rightarrow \frac{x^2}{1-x} - \frac{x^2}{2-x} = \frac{x}{1+x} - \frac{x}{2+x}$$

$$\Rightarrow \frac{x^2}{(1-x)(2-x)} = \frac{x}{(2+x)(1+x)}$$

$$\Rightarrow x = 0 \text{ and } x(2+x)(1+x) = (1-x)(2-x)$$

$$\text{Let } h(x) = x(2+x)(1+x) - (1-x)(2-x)$$

$$= x^3 + 2x^2 + 5x - 2$$

$$\text{Now, } h\left(-\frac{1}{2}\right) = -\frac{1}{2}\left(2-\frac{1}{2}\right)\left(1-\frac{1}{2}\right) - \left(1+\frac{1}{2}\right)\left(2+\frac{1}{2}\right) \\ = -\frac{1}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} - \frac{3}{2} \cdot \frac{5}{2} = \frac{3}{2} \left(-\frac{1}{4} - \frac{5}{2}\right) < 0$$

$$h\left(\frac{1}{2}\right) = \frac{1}{2}\left(2+\frac{1}{2}\right)\left(1+\frac{1}{2}\right) - \left(1-\frac{1}{2}\right)\left(2-\frac{1}{2}\right) \\ = \frac{1}{2} \cdot \frac{5}{2} \cdot \frac{3}{2} - \frac{1}{2} \cdot \frac{3}{2} > 0$$

$\therefore \exists$ a root between $-\frac{1}{2}$ and $\frac{1}{2}$.

$$\text{Also, } h'(x) = 3x^2 + 4x + 5 > 0$$

$\Rightarrow h(x)$ has exactly one real root in $\left(-\frac{1}{2}, \frac{1}{2}\right)$

15. (3748) : Here $X = \{1, 6, 11, \dots, 10086\}$

and $Y = \{9, 16, 23, \dots, 14128\}$

The intersection of X and Y is an A.P. with 16 as first term and 35 as common difference.

The sequence becomes 16, 51, 86,

Now, k^{th} term $= 16 + (k-1) 35 \leq 10086$

$$\text{i.e. } k \leq \frac{10105}{35} \quad \therefore k \leq 288 \text{ (As } k \text{ is to be an integer)}$$

$$\text{Hence, } n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$$

$$= 2018 + 2018 - 288 = 3748$$

16. (1) : We have

$$\log y_n = \frac{1}{n} \left\{ \log \left(1 + \frac{1}{n}\right) + \log \left(1 + \frac{2}{n}\right) + \dots + \log \left(1 + \frac{x}{n}\right) \right\}$$

$$\therefore \lim_{n \rightarrow \infty} \log y_n = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \log \left(1 + \frac{r}{n}\right)$$

$$= \int_0^1 \log(1+x) dx = \int_1^2 \log x dx$$

$$= [x \log x - x]_1^2 = 2 \log 2 - 1 = \log \frac{4}{e}$$

$$\therefore L = 4/e \Rightarrow [L] = 1.$$

$$\text{17. (15) : } \lim_{x \rightarrow 0} \frac{\tan[e^2]x^4 - \tan[-e^2]x^4}{\sin^4 x}$$

$$= \lim_{x \rightarrow 0} \frac{\tan 7x^4 - \tan(-8)x^4}{\sin^4 x} = \lim_{x \rightarrow 0} \frac{\tan 7x^4 + \tan 8x^4}{\sin^4 x}$$

$$= \frac{7 \lim_{x \rightarrow 0} \frac{\tan 7x^4}{7x^4} + 8 \lim_{x \rightarrow 0} \frac{\tan 8x^4}{8x^4}}{\lim_{x \rightarrow 0} \frac{\sin^4 x}{x^4}} = \frac{7+8}{1} = 15$$

$$\text{18. (5) : } AB = \begin{bmatrix} 1 & \frac{3}{2} \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 4 & -3 \\ -2 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

$\therefore (AB)^1 C_1 = C_1, (AB)^2 C_2 = C_2$ and so on.

Also, $\text{tr}(C_r) = r \cdot 3^r + (r-1) \cdot 3^r = (2r-1) \cdot 3^r$

$$\text{Now, } \sum_{r=1}^{50} \text{tr}((AB)^r C_r) = \text{tr}((AB)^1 C_1) + \text{tr}((AB)^2 C_2) + \dots + \text{tr}((AB)^{50} C_{50}) = S \text{ (Let)}$$

$$\therefore S = \text{tr}(C_1) + \text{tr}(C_2) + \dots + \text{tr}(C_{50})$$

$$S = 1 \cdot 3^1 + 3 \cdot 3^2 + 5 \cdot 3^3 + \dots + 99 \cdot 3^{50}$$

$$3S = 1 \cdot 3^2 + 3 \cdot 3^3 + \dots + 97 \cdot 3^{50} + 99 \cdot 3^{51}$$

Subtracting above equations, we get

$$-2S = 1 \cdot 3 + 2 \cdot 3^2 + 2 \cdot 3^3 + \dots + 2 \cdot 3^{50} - 99 \cdot 3^{51}$$

$$= -3 + 2 \cdot 3 + 2 \cdot 3^2 + \dots + 2 \cdot 3^{50} - 99 \cdot 3^{51}$$

$$= -3 + 2 \cdot \frac{3 \cdot (3^{50}-1)}{3-1} - 99 \cdot 3^{51} = -3 + 3^{51} - 3 - 99 \cdot 3^{51}$$

$$= -6 - 98 \cdot 3^{51} \Rightarrow S = 3 + 49 \cdot 3^{51}$$

$$\therefore a+b = 100$$

$$\text{Hence } \frac{1}{20}(a+b) = 5$$



Challenging PROBLEMS



ON
JEE



Single Option Correct Type

1. If the sum of lengths of the hypotenuse and another side of a right angled triangle is given. The area of the triangle is maximum, then the angle between these is
 (a) $\frac{\pi}{12}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{3}$ (d) $\frac{\pi}{2}$

2. If n arithmetic means are inserted between two sets of numbers $a, 2b$ and $2a, b$, where $a, b \in R$. Suppose that m^{th} arithmetic means between these two sets of numbers is same, then the ratio $a : b$ equals to
 (a) $(n - m + 1) : m$ (b) $(n - m + 1) : n$
 (c) $m : (n - m + 1)$ (d) $n : (n - m + 1)$

3. The cartesian equation of the plane passing through the line of intersection of the planes $\vec{r} \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 1$ and $\vec{r} \cdot (\hat{i} - \hat{j}) + 4 = 0$ and perpendicular to the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 8 = 0$ is
 (a) $3x - 4y + 4z = 5$
 (b) $x - 2y + 4z = 3$
 (c) $5x - 2y - 12z + 47 = 0$
 (d) $2x + 3y + 4 = 0$

4. In a three dimensional co-ordinate system P, Q and R are images of a point $A(a, b, c)$ in the xy , yz and zx planes respectively. If G is the centroid of triangle PQR , then area of triangle AOG is (O is the origin)
 (a) 0 (b) $a^2 + b^2 + c^2$
 (c) $\frac{2}{3}(a^2 + b^2 + c^2)$ (d) None of these

5. Two natural numbers a and b are selected at random. The probability that $a^2 + b^2$ is divisible by 7 is
 (a) $3/8$ (b) $1/7$ (c) $3/49$ (d) $1/49$

6. The number of positive integral solutions of the

$$\text{equation } \begin{vmatrix} y^3 + 1 & y^2 z & y^2 x \\ y z^2 & z^3 + 1 & z^2 x \\ y x^2 & x^2 z & x^3 + 1 \end{vmatrix} = 11 \text{ is}$$

(a) 1 (b) 2 (c) 3 (d) 4

More Than One Option Correct Type

7. Let P be the point on the parabola $y^2 = 8x$ which is at the least distance from the centre C of the circle $x^2 + y^2 - 8x - 32y + 256 = 0$. If Q be the point on the circle dividing the line segment CP internally, then
 (a) x intercept of the normal at P is 12.
 (b) length of $QP = 4(\sqrt{5} - 1)$.
 (c) equation of tangent at P is $x - 2y + 8 = 0$.
 (d) slope of tangent to the circle at $Q = 1/2$.

8. If $2x - y + 1 = 0$ is tangent to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{16} = 1$, then which of the following cannot be the sides of a right angled triangle?
 (a) $a, 4, 2$ (b) $a, 4, 1$
 (c) $2a, 8, 1$ (d) $2a, 4, 1$

9. The eccentric angle of a point on the ellipse $3x^2 + 5y^2 = 15$ at distance 2 units from the origin is
 (a) $\frac{\pi}{4}$ (b) $\frac{3\pi}{4}$
 (c) $\frac{5\pi}{4}$ (d) $\frac{7\pi}{4}$

10. If $\cos \theta + \cos \phi = \alpha$, $\cos 2\theta + \cos 2\phi = \beta$ and $\cos 3\theta + \cos 3\phi = \gamma$, then
 (a) $\cos^2 \theta + \cos^2 \phi = 1 + \frac{\beta}{2}$
 (b) $\cos \theta \cdot \cos \phi = \frac{\alpha^2 - \beta + 2}{2}$
 (c) $2\alpha^3 + \gamma = 3\alpha(1 + \beta)$
 (d) $\alpha + \beta + \gamma = 3\alpha\beta\gamma$

Comprehension Type

Paragraph for Q. No. 11 and 12

If a source of light is placed at the fixed point of a parabola and if the parabola is a reflecting surface, then the ray will bounce back in a line parallel to the axis of the parabola.

- 11.** A ray of light is coming along the line $y = 2$ from the positive direction of x -axis and strikes a concave mirror whose intersection with the x - y plane is a parabola $y^2 = 8x$, then the equation of the reflected ray is
 (a) $2x + 5y = 4$ (b) $3x + 2y = 6$
 (c) $4x + 3y = 8$ (d) $5x + 4y = 10$

- 12.** A ray of light moving parallel to the x -axis gets reflected from a parabolic mirror whose equation is $y^2 + 10y - 4x + 17 = 0$. After reflection, the ray must pass through the point

- (a) $(-2, -5)$ (b) $(-1, -5)$
 (c) $(-3, -5)$ (d) $(-4, -5)$

Paragraph for Q. No. 13 and 14

If $S_1 = 0$ and $S_2 = 0$ are equations of circles intersecting in real and distinct points A and B , then AB is common chord of the circles $S_1 = 0$, $S_2 = 0$ known as radical axis of circles which is \perp to line segment joining the centres of the circles. If $S_1 = x^2 + y^2 + 2gx + 2fy + c = 0$ and $S_2 = x^2 + y^2 + 2g'x + 2f'y + c' = 0$, then equation of radical axis is given by $2x(g - g') + 2y(f - f') + c - c' = 0$. If circles touching each other externally or internally, then point of contact can be determined by section formula. Consider the circles $S_1 = x^2 + y^2 - 4x - 2y + 4 = 0$ and $S_2 = x^2 + y^2 - 12x - 8y + 36 = 0$.

- 13.** The point at which the circles $S_1 = 0$, $S_2 = 0$ touches each other is

- (a) $\left(\frac{8}{5}, \frac{14}{5}\right)$ (b) $\left(\frac{14}{5}, \frac{8}{5}\right)$
 (c) $\left(-\frac{8}{5}, \frac{14}{5}\right)$ (d) $\left(\frac{8}{5}, -\frac{14}{5}\right)$

- 14.** The equation of direct common tangent to the circles $S_1 = 0$ and $S_2 = 0$ is

- (a) $y = 2$ (b) $x = 0$
 (c) $24x - 7y = 16$ (d) $24x + 7y = 16$

Matrix Match Type

- 15.** Match the following :

	Column-I	Column-II
A.	If e be the eccentricity of the hyperbola $36x^2 + 288x - 64y^2 - 128y - 1792 = 0$, then the value of $4e$ equals	P. 7
B.	The number of values of K , such that the line $y = 4x + K$ touches the curve $x^2 + 4y^2 = 4$ is	Q. 3

C.	The eccentricity of an ellipse with centre at $O(0, 0)$ is $\frac{1}{3}$. If one directrix is $x = 6$ and ellipse reduces to $K_1x^2 + K_2y^2 = K_3$, then $K_3 - K_1 - K_2 =$	R. 5
D.	If the foci of an ellipse and the hyperbola whose equations are respectively $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$ and $\frac{25x^2}{144} - \frac{25y^2}{81} = 1$, coincide with each other, then the value of b^2 equals	S. 2

Numerical Value Type

- 16.** If $2\tan^2 x - 5\sec x = 1$ for exactly 7 distinct values of $x \in \left[0, \frac{n\pi}{2}\right], n \in \mathbb{N}$, then the greatest value of n is ____.

- 17.** From a point P outside a circle with centre at C , tangents PA and PB are drawn such that $\frac{1}{(CA)^2} + \frac{1}{(PA)^2} = \frac{1}{16}$, then the length of chord AB is _____.

- 18.** If 7 divides $32^{32^{32}}$, the remainder is ____.

- 19.** There exist positive integers A , B and C with no common factors greater than 1, such that $A \log_{2005} 5 + B \log_{2002} 2 = C$. The sum $A + B + C$ equals to ____.

- 20.** The number of 4-digit numbers that can be formed by using the digits 1, 2, 3, 4, 5, 6, 7, 8 and 9 such that the least digit used is 4, when repetition of digits is allowed, is ____.

SOLUTIONS

1. (c) : $AB + AC = \text{constant} = k$


$\therefore BC^2 = (k - x)^2 - x^2 = k^2 - 2kx$

$\therefore \Delta = \frac{1}{2} BC \cdot AB = \frac{1}{2} x \sqrt{k^2 - 2kx}$

Let $Z = \Delta^2 = \frac{1}{4} x^2 (k^2 - 2kx)$

$= \frac{1}{4} (k^2 x^2 - 2kx^3)$

Z will be max, when $x = k/3$



$$\text{Now, } \cos \theta = \frac{x}{k-x} = \frac{k/3}{k-k/3} = \frac{1}{2}$$

$$\therefore \theta = \pi/3$$

2. (c) : Let A_1, A_2, \dots, A_n be arithmetic means between a and $2b$, then $A_m = a + m\left(\frac{2b-a}{n+1}\right)$

Again, let B_1, B_2, \dots, B_n be arithmetic means

$$\text{between } 2a \text{ and } b, \text{ then } B_m = 2a + m\left(\frac{b-2a}{n+1}\right)$$

Now, $A_m = B_m$ (Given)

$$\Rightarrow a + m\left(\frac{2b-a}{n+1}\right) = 2a + m\left(\frac{b-2a}{n+1}\right)$$

$$\Rightarrow m\left(\frac{b+a}{n+1}\right) = a \Rightarrow \frac{a}{b} = \frac{m}{n-m+1}$$

3. (c) : Equation of any plane passing through the intersection of the planes $\vec{r} \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 1$ and

$$\vec{r} \cdot (\hat{i} - \hat{j}) + 4 = 0 \text{ is } 2x - 3y + 4z - 1 + \lambda(x - y + 4) = 0$$

$$\text{or } (2 + \lambda)x - (3 + \lambda)y + 4z + 4\lambda - 1 = 0$$

The plane is perpendicular to the plane

$$\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 8 = 0 \text{ if } 2(2 + \lambda) + (3 + \lambda) + 4 = 0$$

$$\Rightarrow 11 + 3\lambda = 0 \Rightarrow \lambda = -11/3$$

The required equation of the plane is

$$3(2x - 3y + 4z - 1) - 11(x - y + 4) = 0$$

$$\Rightarrow 5x - 2y - 12z + 47 = 0$$

4. (a) : Point A is $(a, b, c) \Rightarrow$ Images of point A i.e., P, Q and R are $(a, b, -c), (-a, b, c)$ and $(a, -b, c)$ respectively.

\therefore Centroid of triangle PQR is $\left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3}\right)$

$$\Rightarrow G \equiv \left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3}\right)$$

$\Rightarrow A, O, G$ are collinear \therefore Area of triangle AOG is 0.

5. (d) : a, b are of the form

$$a, b \in \{7m, 7m+1, 7m+2, 7m+3, 7m+4, 7m+5, 7m+6\}$$

$$\text{Now, } a^2, b^2 \in \{7m_1, 7m_1+1, 7m_1+4, 7m_1+2, 7m_1+2, 7m_1+4, 7m_1+1\}$$

For required condition, a^2, b^2 must be of the form $7m$.

$$\therefore \text{Required probability} = \frac{1}{49}$$

6. (c) : Multiply R_1, R_2 and R_3 by y, z and x respectively and then take common y, z and x from C_1, C_2, C_3 respectively, then

$$\begin{vmatrix} y^3 + 1 & y^3 & y^3 \\ z^3 & z^3 + 1 & z^3 \\ x^3 & x^3 & x^3 + 1 \end{vmatrix} = 11 \Rightarrow \begin{vmatrix} 1 & 0 & y^3 \\ -1 & 1 & z^3 \\ 0 & -1 & x^3 + 1 \end{vmatrix} = 11$$

(Applying $C_1 \rightarrow C_1 - C_2$ and $C_2 \rightarrow C_2 - C_3$)

$$\Rightarrow 1(x^3 + 1 + z^3) + y^3(1) = 11 \Rightarrow x^3 + y^3 + z^3 = 10$$

So, solutions are $(1, 1, 2), (1, 2, 1)$ or $(2, 1, 1)$

7. (a, b, c, d) : Equation of circle is

$$x^2 + y^2 - 8x - 32y + 256 = 0$$

$$\Rightarrow (x - 4)^2 + (y - 16)^2 = 4^2$$

\therefore Centre $C \equiv (4, 16)$ and radius $R = 4$

Now, equation of normal to the parabola $y^2 = 8x$ is

$$y = mx - 4m - 2m^3 \quad \dots(i)$$

For the least distance from the centre C to parabola the normal must pass through the centre $(4, 16)$ of circle

$$\therefore 16 = 4m - 4m - 2m^3$$

$$\Rightarrow m^3 = (-2)^3 \Rightarrow m = -2$$

\therefore Equation of normal is $y = -2x + 24$ (Using (i))

\therefore Slope of tangent at $Q = 1/2$.

Also, x intercept of normal at P is obtained by putting $y = 0$ in (i), $x = 12$

Now, any point on the parabola $y^2 = 4ax$ is $(am^2, -2am) = (8, 8)$ (using $a = 2, m = -2$)

\therefore Equation of tangent to parabola at P is

$$y - 8 = \frac{1}{2}(x - 8) \Rightarrow x - 2y + 8 = 0$$

Now $PQ = CP - CQ = CP - \text{radius of circle (R)}$

$$= 4\sqrt{5} - 4 = 4(\sqrt{5} - 1)$$

8. (a, b, c) : The line $2x - y + 1 = 0$ be tangent to the

hyperbola $\frac{x^2}{a^2} - \frac{y^2}{16} = 1$ if both roots of the equation

$$\frac{x^2}{a^2} - \frac{(2x+1)^2}{16} = 1 \text{ are equal.}$$

i.e., discriminant of $(16 - 4a^2)x^2 - 4a^2x - 17a^2 = 0$

$$\Rightarrow 16a^4 = 4(-17a^2)(16 - 4a^2) \Rightarrow a = \sqrt{17}/2$$

Thus, we have set of numbers

$$\left(\frac{\sqrt{17}}{2}, 4, 2\right); \left(\frac{\sqrt{17}}{2}, 4, 1\right); (\sqrt{17}, 8, 1) \text{ and } (\sqrt{17}, 4, 1)$$

Only set $(\sqrt{17}, 4, 1)$ represents the sides of a right angled triangle.

9. (a, b, c, d) : We have, $\frac{x^2}{5} + \frac{y^2}{3} = 1$... (i)

\therefore The point (P) on the ellipse is $(\sqrt{5} \cos \theta, \sqrt{3} \sin \theta)$, where θ is eccentric angle.

Given, distance $OP = 2$ units (where O is origin)

$$\therefore 5\cos^2\theta + 3\sin^2\theta = (2)^2 \Rightarrow \cos 2\theta = 0$$

$$\therefore 2\theta = (2n+1)\frac{\pi}{2}, (n \in I) \Rightarrow 2\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}$$

$$\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$$

10. (a, b, c) : $(\cos \theta + \cos \phi)^2 = \alpha^2$... (i)

$$\Rightarrow \cos^2\theta + \cos^2\phi + 2\cos \theta \cos \phi = \alpha^2$$

$$\text{Now, } \cos 2\theta + \cos 2\phi = \beta$$

$$\Rightarrow (2\cos^2\theta - 1) + (2\cos^2\phi - 1) = \beta$$

$$\Rightarrow 2(\cos^2\theta + \cos^2\phi) = \beta + 2$$

$$\Rightarrow \cos^2\theta + \cos^2\phi = \frac{\beta}{2} + 1$$
 ... (ii)

From (i) and (ii), we get $\cos\theta \cdot \cos\phi = \frac{\alpha^2}{2} - \frac{\beta+2}{4}$

$$\text{Also, } \cos 3\theta + \cos 3\phi = \gamma$$

$$\Rightarrow (4\cos^3\theta - 3\cos\theta) + (4\cos^3\phi - 3\cos\phi) = \gamma$$

$$\Rightarrow 4(\cos^3\theta + \cos^3\phi) - 3(\cos\theta + \cos\phi) = \gamma$$

$$\Rightarrow 4[(\cos\theta + \cos\phi)(\cos^2\theta + \cos^2\phi - \cos\theta \cos\phi)] - 3(\cos\theta + \cos\phi) = \gamma$$

$$\Rightarrow 4\left[\alpha\left(\frac{\beta+2}{2} - \frac{1}{2}\left(\alpha^2 - \frac{(\beta+2)}{2}\right)\right)\right] - 3\alpha = \gamma$$

$$\therefore 2\alpha^3 + \gamma = 3\alpha(1 + \beta)$$

11. (c) : Point of intersection of $y = 2$ and $y^2 = 8x$ is

$$P\left(\frac{1}{2}, 2\right)$$
 and focus of the parabola is $S(2, 0)$

$$\therefore \text{Equation of the reflected ray is } y - 0 = \frac{2-0}{1/2-2}(x-2)$$

$$\Rightarrow y = -\frac{4}{3}(x-2) \Rightarrow 4x + 3y = 8$$

12. (b) : $\because y^2 + 10y - 4x + 17 = 0$

$$\Rightarrow (y+5)^2 - 25 - 4x + 17 = 0$$

$$\Rightarrow (y+5)^2 = 4x + 8 \Rightarrow (y+5)^2 = 4(x+2)$$

Let $y+5 = Y$, $x+2 = X$, then $Y^2 = 4X$

Focus is $X = 1$, $Y = 0$, i.e., $(-1, -5)$

After reflection, the ray must pass through focus $(-1, -5)$


13. (b) : Given $S_1 = x^2 + y^2 - 4x - 2y + 4 = 0$

$$\Rightarrow C_1(2, 1)$$
 and $r_1 = 1$

$$S_2 = x^2 + y^2 - 12x - 8y + 36 = 0 \Rightarrow C_2(6, 4)$$
 and $r_2 = 4$

Now, $r_1 + r_2 = 1 + 4 = 5 = C_1C_2$ (distance between the centres of circles).

As $r_1 + r_2 = C_1C_2$ which means circles touch each other externally at a point say P which divides C_1C_2 in the ratio $1 : 4$



The co-ordinates of P

$$= \left(\frac{1 \times 6 + 4 \times 2}{5}, \frac{1 \times 4 + 4 \times 1}{5} \right) = \left(\frac{14}{5}, \frac{8}{5} \right)$$

14. (c) : The direct common tangents meet (intersect) each other at $\left(\frac{2}{3}, 0\right)$, let its slope be m and its equation is given by $y - 0 = m\left(x - \frac{2}{3}\right) \Rightarrow y = mx - \frac{2m}{3}$

As direct common tangents are perpendicular to the lines joining the point of contact and centres of the circles and the distance from point of contact to centres is equal to the radii of circles.

\therefore Distance from $(2, 1)$ to the tangent line $mx - y - \frac{2m}{3} = 0$ is equal to 1.

$$\therefore \frac{2m - 1 - \frac{2m}{3}}{\sqrt{1+m^2}} = 1 \Rightarrow m = 0, m = \frac{24}{7}$$

\therefore Equations of direct common tangents to S_1 and S_2 are $y = 0$ and $y = \frac{24}{7}x - \frac{16}{7} \Rightarrow 24x - 7y = 16$.

15. A \rightarrow R, B \rightarrow S, C \rightarrow Q, D \rightarrow P

$$(A) 36x^2 + 288x - 64y^2 - 128y - 1792 = 0$$

$$\Rightarrow 36(x+4)^2 - 64(y+1)^2 = 2304$$

$$\Rightarrow \frac{(x+4)^2}{64} - \frac{(y+1)^2}{36} = 1 \Rightarrow a^2 = 64, b^2 = 36$$

$$\therefore e = \sqrt{1 + \frac{36}{64}} = \frac{10}{8} = \frac{5}{4} \Rightarrow 4e = 5$$

(B) Given curve is $x^2 + 4y^2 = 4 \Rightarrow \frac{x^2}{4} + \frac{y^2}{1} = 1$

It is an ellipse, where $a^2 = 4$, $b^2 = 1$ and the line $y = mx + K$ will be tangent if $K^2 = a^2m^2 + b^2$

$$\therefore K^2 = 4 \cdot 4^2 + 1 \Rightarrow K = \pm \sqrt{65}$$

\therefore Number of values of $K = 2$




(C) Equation of directrix is given by

$$x = \frac{a}{e} \Rightarrow a = xe \Rightarrow a = 2$$

$$\text{Now, } b^2 = a^2(1 - e^2) = 4\left(1 - \frac{1}{9}\right) = \frac{4 \times 8}{9} = \frac{32}{9}$$

Some of the best lessons are learnt from history!

Introducing Chapterwise Solutions to JEE.
Crack the past to see if you will last.



Tune. Fine tune. Reach the peak of your readiness for JEE with MTG's 43-19 Years Chapterwise Solutions. It is undoubtedly the most comprehensive 'real' question bank, complete with detailed solutions by experts.

Studies have shown that successful JEE aspirants begin by familiarising themselves with the problems that have appeared in past JEEs as early as 2 years in advance.

Making it one of the key ingredients for their success. How about you then?
Get 43+19 Years Chapterwise Solutions to start your rehearsals early.



Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Visit www.mtg.in to buy online!
Call 0124-6601200 or e-mail:info@mtg.in

\therefore The equation of ellipse is $\frac{x^2}{4} + \frac{9y^2}{32} = 1$

$$\Rightarrow 8x^2 + 9y^2 = 32 \Rightarrow K_1x^2 + K_2y^2 = K_3$$

$$\therefore \frac{K_3 - K_1 - K_2}{5} = \frac{32 - 8 - 9}{5} = 3$$

(D) For hyperbola, we have

$$e^2 = \frac{a^2 + b^2}{a^2} = \frac{144 + 81}{144} = \frac{225}{144} \Rightarrow e = \frac{5}{4}$$

$$\text{and } a^2 = \frac{144}{25} \Rightarrow a = \frac{12}{5}$$

$$\therefore \text{Foci are } (\pm ae, 0) = \left(\pm \frac{12}{5} \times \frac{5}{4}, 0 \right) = (\pm 3, 0)$$

Now, for ellipse ($ae = 3$) (foci of an ellipse and hyperbola coincide with each other), $a = 4$

$$\text{Now, } b^2 = a^2(1 - e^2) = a^2 - a^2e^2 = 16 - 9 = 7$$

$$16. (15) : \text{ Now, } \sec x = 3 \Rightarrow \cos x = \frac{1}{3}$$

It gives two values of x in each of $[0, 2\pi]$, $(2\pi, 4\pi]$,


$$(4\pi, 6\pi] \text{ and one value in } 6\pi + \frac{3\pi}{2} = \frac{15\pi}{2}$$

\therefore Greatest value of $n = 15$

$$17. (8) : \tan \theta = \frac{r}{PA}$$

$$\text{Given } \frac{1}{r^2} + \frac{1}{PA^2} = \frac{1}{16}$$

$$\Rightarrow \cot^2 \theta + 1 = \frac{1}{16}$$



$\Rightarrow PA \sin \theta = x = 4 \therefore \text{Length of Chord } AB = 2x = 8$

$$18. (4) : 32 = 2^5 \Rightarrow (32)^{32} = (2^5)^{32}$$

$$= 2^{160} = (3 - 1)^{160} = 3m + 1, m \in N$$

$$\therefore (32)^{32} = (32)^{3m+1} = 2^{5(3m+1)}$$

$$= 2^{3(5m+1)}2^2 = 4 \cdot 8^{5m+1} = 4(7 + 1)^{5m+1}$$

$$= 4(7n + 1), n \in N = 28n + 4$$

\therefore When 7 divides $(32)^{32}$, remainder = 4

$$19. (6) : \text{ Given } A \log_{200} 5 + B \log_{200} 2 = C$$

$$\Rightarrow A \log 5 + B \log 2 = C \log 200 = C \log(5^2 \cdot 2^3) = 2C \log 5 + 3C \log 2$$

Hence, $A = 2C$ and $B = 3C$

For no common factor greater than 1, $C = 1$

So, $A = 2$, $B = 3$. Thus $A + B + C = 6$

20. (671) : Least digit used = 4

\therefore We can use 4, 5, 6, 7, 8, 9. But remember that at least one 4 must be used.

Now, 1st blank can be filled in 6 ways.

2nd blank can be filled in 6 ways.

3rd blank can be filled in 6 ways.

4th blank can be filled in 6 ways.

\therefore 4 blanks can be filled in 6^4 ways. But out of these, some may contain no digit 4 at all. Let us find them.

Each blank can be filled in 5 ways (by 5, 6, 7, 8, or 9)

\therefore 4 blanks can be filled in 5^4 ways (no 4 at all)

\therefore Required Number = $6^4 - 5^4$ (at least one 4) = 671.

Your favourite MTG Books/Magazines available in KARNATAKA at

Renuka Book Stall - Bagalkot Ph: 8183026201

Hema Book Stores - Bangalore Ph: 8041485110

Prakash Sahitya Book Sellers And Publ - Bangalore

Ph: 080-2254950; 9900118800, 9844028710

Sri Book World - Bangalore Ph: 080-2624904, 26608965, 4125679, 9448260596

Sri Sri Ram Book House - Bangalore Ph: 080-41472635, 9449212946, 973881088

Gourav Book Centre - Belgaum Ph: 0831-201929, 4201859, 9341101353

Hema Book World - Bengaluru Ph: 080-40905110, 9482266600, 9945731121

Avenue Book Centre - Bengaluru

Ph: 080-22244753-41474466; 8022244753

Saraswat Book Centre - Bengaluru Ph: 080-23152101, 080-23308673; 9845351101

Sapna Book House Pvt. Ltd. - Bengaluru

Ph: 080-2260088, 40114455; 9901471718, 9844013242

UBS Publishers Distributors Pvt Ltd. - Bengaluru

Ph: 26756671; Mob: 9341621469, 8026756462

Bumblebee Book Depot - Bidar Ph: 9241059982

Mohan Book Depot - Bijapur Ph: 8352253614

Kannadra Book Stall - Bijapur Ph: 9448645098

Shri Hanshankar Books & General Store - Chitradurga Ph: 230069; 9448654882

Lakshmi Agencies - Davanagere Ph: 9844168836, 8192231271, 8192231271

Shri Mahaxay Book Depot - Davanagere Ph: 08192-255591; 8192255591, 9972655736

Nehru Book Depot - Davaregare Ph: 08192-270395; 9886210458, 9740681601

Competitor Book Centre - Dhawrad Ph: 0836-2746514, 24747349, 9802679413

Bharat Book Depot - Dhanwad Ph: 836-2447349; 9945506375, 8382447349

E.S. Bhavikatti & Sons - Gubarga Ph: 9880672425

L.E. Bhavikatti Book House - Gulbarga Ph: 08472-261400, 9880737400

S S Bhavikatti - Gulbarga Ph: 225320, 934139399, 8472225320

Vijay Book Centre - Hubli Ph: 0836-4258401, 2263101; 9538918919

SLV Book House - Mangalore Ph: 9972683044

School Book Company - Mangalore Ph: 0824-249904

Avas Book & Stationery - Moodbidri Ph: 849450761

Suma Books & Stationery - Moodbidri Ph: 9448329074, 8258261264

Diana Book Gallery - Shimoga Ph: 08182-220953; 9884205772, 9844205772

S.L.N. Enterprises - Shimoga Ph: 08182-220953; 9884205772, 9844205772

Namana Book Palace - Tumkur Ph: 0816-2251223, 2277774, 9141490031, 98089244

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call 0124-6601200 for further assistance.

beat the **TIME TRAP**



Duration : 30 minutes

SECTION-I

Single Option Correct Type

- BC is latus rectum of a parabola $y^2 = 4ax$ and A is its vertex. The minimum length of projection of BC on a tangent drawn in portion BAC is
 (a) $\sqrt{2}a$ (b) $2\sqrt{2}a$
 (c) $2a$ (d) $3\sqrt{2}a$
- Least value of the expression

$$\frac{1}{2bx - (x^2 + b^2 + \sin^2 x)}, x \in [-1, 0], b \in [2, 3]$$
 is
 (a) $\frac{1}{4}$ (b) $-\frac{1}{4}$
 (c) $-\frac{1}{8 + \sin^2 1}$ (d) none of these
- If in a right angled triangle ABC, $4\sin A \cos B - 1 = 0$ and $\tan A$ is real, then
 (a) angles are in A.P. (b) angles are in G.P.
 (c) angles are in H.P. (d) none of these
- If $|z| = 2$ and $\frac{z_1 - z_3}{z_2 - z_3} = \frac{z - 2}{z + 2}$, then z_1, z_2 and z_3 will be vertices of a
 (a) equilateral triangle
 (b) acute angled triangle
 (c) right angled triangle
 (d) none of these
- If $a^2 + b^2 - c^2 - 2ab = 0$, then the point(s) of concurrency of family of straight lines $ax + by + c = 0$ lie(s) on the line
 (a) $y = x$ (b) $y = x + 1$
 (c) $y = -x$ (d) $x + y = 1$
- If $f''(x) > 0 \forall x \in R, f'(3) = 0$ and $g(x) = f(\tan^2 x - 2 \tan x + 4), 0 < x < \frac{\pi}{2}$, then $g(x)$ is increasing in
 (a) 1 (b) 2
 (c) 0 (d) none of these
- A vector \vec{r} is equally inclined with the vectors $\vec{a} = \cos \theta \hat{i} + \sin \theta \hat{j}$, $\vec{b} = -\sin \theta \hat{i} + \cos \theta \hat{j}$ and $\vec{c} = \hat{k}$, then angle between \vec{r} and \vec{a} is
 (a) $\cos^{-1}\left(\frac{1}{\sqrt{2}}\right)$ (b) $\cos^{-1}\left(\frac{1}{3}\right)$
 (c) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (d) $\frac{\pi}{2}$
- If the roots of the equation $x^2 + ax + b = 0$ are c and d , then one of the roots of the equation $x^2 + (2c + a)x + c^2 + ac + b = 0$ is
 (a) c (b) $d - c$
 (c) $2c$ (d) $2d$
- Let f be a differentiable function satisfying the condition $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$ for all $x, y (\neq 0) \in R$ and $f(y) \neq 0$. If $f'(1) = 2$, then $f'(x)$ is equal to
 (a) $2f(x)$ (b) $\frac{f(x)}{x}$
 (c) $2xf(x)$ (d) $\frac{2f(x)}{x}$
- The number of solutions of the equation $\cos^{-1} x + \cos^{-1} \sqrt{1-x^2} = \pi$ is
 (a) 1 (b) 2
 (c) 0 (d) none of these

SECTION-II

Numerical Answer Type

11. If $(1+x)^n = \sum_{r=1}^n C_r x^r$, then the value of $C_1 + 2C_2 + 3C_3 + \dots + nC_n$ equals $n \cdot k^{n-1}$. Find k

12. Let $f: (0, \infty) \rightarrow R$ and $F(x) = \int_0^x f(t) dt$.
If $F(x^2) = x^2(1+x)$, then $f(4)$ equals _____.

13. The coefficients of three consecutive terms of $(1+x)^{n+5}$ are in the ratio $5 : 10 : 14$. Then $n = \text{_____}$.

14. If the function $f(x) = x^3 + e^{\frac{x}{2}}$ and $g(x) = f^{-1}(x)$, then the value of $g'(1)$ is _____.

15. Let ω be the complex number $\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$. The number of distinct complex numbers z satisfying $\begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$ is equal to _____.

SOLUTIONS

1. (b) : Let tangent at $P(at^2, 2at)$ makes an angle θ with x -axis, then $\tan \theta = \frac{1}{t}$

Projection of BC on tangent $= BC \sin \theta$
 $= \frac{4a}{\sqrt{1+t^2}} \geq 2a\sqrt{2}$ (as $-1 \leq t \leq 1$).

2. (b) : Given expression will have least value if $2bx - [x^2 + b^2 + \sin^2 x]$ is maximum
 $x^2 + b^2 + \sin^2 x - 2bx$ is minimum
 $(x-b)^2 + \sin^2 x$ is minimum
Now $|x-b|$ and $|\sin x|$ are minimum if $x=0$, $b=2$
So, least value is $-\frac{1}{4}$


3. (a) : Since, $4 \sin A \cos B = 1$, so A and B can not be $\frac{\pi}{2}$
[As if $B = \frac{\pi}{2}$, then $\cos B = 0$ and if $A = \frac{\pi}{2}$, $\tan A$ is not defined]

$$C = \frac{\pi}{2}, B = \frac{\pi}{2} - A \Rightarrow 4 \sin A \cos \left(\frac{\pi}{2} - A \right) = 1$$

$$\Rightarrow \sin^2 A = \frac{1}{4} \Rightarrow \sin A = \frac{1}{2} \Rightarrow A = \frac{\pi}{6} \Rightarrow B = \frac{\pi}{3}$$

So angles are $\frac{4\pi}{6}, \frac{\pi}{3}, \frac{\pi}{2}$ which are in A.P.

4. (c) : Clearly, $\operatorname{Arg} \left(\frac{z-2}{z+2} \right) = \pm \frac{\pi}{2}$



$$\Rightarrow \operatorname{Arg} \left(\frac{z_1 - z_3}{z_2 - z_3} \right) = \pm \frac{\pi}{2}$$

So z_1, z_2, z_3 will be the vertices of a right angled triangle.

5. (c) : $(a-b)^2 - c^2 = 0$

$$\Rightarrow (a-b-c)(a-b+c) = 0$$

$$\text{If } a-b=c \Rightarrow ax+by+a-b=0$$

$$\Rightarrow (x+1)a+b(y-1)=0 \Rightarrow x=-1, y=1$$

$$\text{If } a-b=c \Rightarrow ax+by+b-a=0$$

$$\Rightarrow (x-1)a+(y+1)b=0$$

$$\Rightarrow (x-1)+(y+1)\frac{b}{a}=0 \Rightarrow x=1, y=-1$$

Equation of line passing through both points $(-1, 1)$ and $(1, -1)$ is $y = -x$.

6. (d) : $g'(x) = f'((\tan x - 1)^2 + 3) = (2 \tan x - 2) \sec^2 x$

Since $f''(x) > 0 \Rightarrow f'(x)$ is increasing

So $f'((\tan x - 1)^2 + 3) > f'(3) = 0$

$$\forall x \in \left(0, \frac{\pi}{4} \right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2} \right)$$

Also $(\tan x - 1) > 0 \quad \forall x \in \left(\frac{\pi}{4}, \frac{\pi}{2} \right)$

So, $g(x)$ is increasing in $\left(\frac{\pi}{4}, \frac{\pi}{2} \right)$.

7. (c) : Since $\vec{a} \cdot \vec{b} = 0 = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} \Rightarrow \vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular.

$$\text{Now } \vec{r} = t(\vec{a} + \vec{b} + \vec{c}) = t((\cos \theta - \sin \theta)\hat{i} + (\cos \theta + \sin \theta)\hat{j} + \hat{k})$$

Let angle between \vec{r} and \vec{a} be α , then

$$\cos \alpha = \frac{\vec{r} \cdot \vec{a}}{|\vec{r}| |\vec{a}|} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \alpha = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right)$$

- 8. (b) :** Let $f(x) = x^2 + ax + b$, then
 $x^2 + (2c+a)x + c^2 + ac + b = f(x+c)$

If c, d are the roots of $f(x)$

$$\therefore c^2 + ac + b = 0 \text{ and } c + d = -a$$

Let α, β be the roots of $f(x+c) = 0$

$$\therefore \text{Product of roots, } \alpha\beta = c^2 + ac + b = 0$$

\therefore One root is 0.

Also, sum of roots = $-(2c+a)$

$$\Rightarrow \alpha + \beta = -(2c+a)$$

$$\Rightarrow 0 + \beta = -(2c - d - a) \Rightarrow \beta = d - c$$

Thus roots of $f(x+c) = 0$ will be 0, $(d-c)$.

- 9. (d) :** We have

$$f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)}$$

Replacing x and y both by 1,

$$f(1) = \frac{f(1)}{f(1)} = 1$$

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= f(x) \lim_{h \rightarrow 0} \left\{ \frac{\frac{f(x+h)}{f(x)} - 1}{h} \right\} = f(x) \lim_{h \rightarrow 0} \frac{f\left(\frac{x+h}{x}\right) - 1}{h} \\ &= f(x) \lim_{x \rightarrow h \rightarrow 0} \frac{f\left(1 + \frac{h}{x}\right) - f(1)}{h/x} \\ &= \frac{f(x)}{x} f'(1) = \frac{2f(x)}{x} \quad (\because f'(1) = 2) \end{aligned}$$

- 10. (a) :** $\cos^{-1} \sqrt{1-x^2} = \pi - \cos^{-1} x = \cos^{-1}(-x)$

$$\Rightarrow \sqrt{1-x^2} = -x \Rightarrow x < 0$$

$$\text{Squaring, } 1 - x^2 = x^2 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

Since $x < 0 \therefore x = -\frac{1}{\sqrt{2}}$ is the only solution.

- 11. (2) :** $C_1 + 2C_2 + 3C_3 + \dots + nC_n$

$$= n + 2 \cdot \frac{n(n-1)}{2!} + \frac{3n(n-1)(n-2)}{3!} + \dots + n$$

$$= n \left[1 + (n-1) + \frac{(n-1)(n-2)}{2!} + \dots + 1 \right]$$

$$= n(1+1)^{n-1} = n2^{n-1} \Rightarrow k = 2$$

- 12. (4) :** $F(x) = \int_0^x f(t) dt \Rightarrow F(x^2) = \int_0^{x^2} f(t) dt = x^2(1+x)$

$$\Rightarrow f(x^2) \cdot 2x = 2x + 3x^2 \Rightarrow f(4) = 4$$

- 13. (6) :** Let T_r, T_{r+1}, T_{r+2} be three consecutive terms.

Coefficients of

$$T_r = {}^{n+5}C_{r-1}, T_{r+1} = {}^{n+5}C_r, T_{r+2} = {}^{n+5}C_{r+1}$$

$$\dots(i) \quad \text{Now, } \frac{T_{r+1}}{T_r} = \frac{10}{5}$$

$$\dots(ii) \quad \Rightarrow \frac{{}^{n+5}C_r}{{}^{n+5}C_{r-1}} = 2 \Rightarrow \frac{n+6-r}{r} = 2$$

$$\Rightarrow n+6-r = 2r \Rightarrow 3r = n+6$$

$$\text{Also, } \frac{T_{r+2}}{T_{r+1}} = \frac{14}{10} \Rightarrow \frac{{}^{n+5}C_{r+1}}{{}^{n+5}C_r} = \frac{7}{5}$$

$$\Rightarrow \frac{n+5-r}{r+1} = \frac{7}{5}$$

$$\Rightarrow \frac{n+6}{r+1} = \frac{12}{5} \Rightarrow \frac{3r}{r+1} = \frac{12}{5} \quad [\text{Using (i)}]$$

$$\Rightarrow 15r = 12r + 12 \Rightarrow 3r = 12 \Rightarrow r = 4$$

$$\therefore n = 3(4) - 6 = 6$$

- 14. (2) :** $f(x) = x^3 + e^{x/2} \Rightarrow f'(x) = 3x^2 + \frac{e^{x/2}}{2}$

we have, $g(f(x)) = x$

On differentiating both sides, w.r.t. 'x', we get

$$g'(f(x)) f'(x) = 1$$

Putting $x = 0$ in above equation, we get

$$g'(f(0)) f'(0) = 1$$

$$\text{But } f(0) = 1 \text{ and } f'(0) = \frac{1}{2}$$

$$\therefore g'(1) = \frac{1}{f'(0)} = \frac{1}{1/2} = 2$$

$$15. (1) : \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} z & z & z \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$$

$$\Rightarrow z\{(z+\omega^2)(z+\omega) - 1 - \omega(z+\omega-1) + \omega^2(1-z-\omega^2)\} = 0$$

$$\Rightarrow z^3 = 0$$

$\Rightarrow z = 0$ is the only solution.




MONTHLY TEST DRIVE CLASS XI | ANSWER KEY

- | | | | | |
|-----------|-----------|---------------|----------|-----------|
| 1. (b) | 2. (a) | 3. (c) | 4. (c) | 5. (a) |
| 6. (c) | 7. (b,c) | 8. (b,c) | 9. (a,c) | 10. (c,d) |
| 11. (b,c) | 12. (a,b) | 13. (a,b,c,d) | 14. (c) | |
| 15. (b) | 16. (b) | 17. (785) | 18. (7) | 19. (6) |
| 20. (3) | | | | |

OLYMPIAD CORNER




1. We are considering triangles ABC in space.
- (a) What conditions must be fulfilled by the angles α, β, γ of triangle ABC in order that there exists a point P in space such that $\angle APB, \angle BPC, \angle CPA$ are right angles?
- (b) Let d be the maximum distance among PA, PB, PC and let h be the longest altitude of triangle ABC . Show that $(\sqrt{6}/3)h \leq d \leq h$.
2. Two circles intersect at A and B . P is any point on an arc AB of one circle. The lines PA, PB intersect the other circle at R and S , as shown below. If P' is any other point on the same arc of the first circle and if R', S' are the points in which the lines $P'A, P'B$ intersect the other circle, prove that the arcs RS and $R'S'$ are equal.



3. For any positive integer n , evaluate a_n/b_n , where

$$a_n = \sum_{k=1}^n \tan^2 \frac{k\pi}{2n+1}, b_n = \prod_{k=1}^n \tan^2 \frac{k\pi}{2n+1}.$$


4. There are 9 regions inside the 5 rings of the Olympics. Put a different whole number from 1 to 9 in each so that the sum of the numbers in each ring is the same. What are the largest and the smallest values of this common sum?



5. If m and n are positive integers such that $\frac{m+n}{m^2+mn+n^2} = \frac{4}{49}$, then find $m+n$.
6. An equilateral triangle of side length 2 units is inscribed in a circle. Find the length of a chord of this circle which passes through the midpoints of two sides of this triangle.
7. In a soccer tournament eight teams play each other once, with two points awarded for a win, one point for a draw and zero for a loss. How many points must a team score to ensure that it is in the top four (*i.e.* has more points than at least four other teams)?

SOLUTIONS

1. (a) By Pythagoras theorem, we have $AB^2 = PA^2 + PB^2$, $AC^2 = PA^2 + PC^2$ and $BC^2 = PB^2 + PC^2$.



Hence $AB^2 + AC^2 - BC^2 = 2PA^2 > 0$, thus $\angle BAC < 90^\circ$, *i.e.* $\alpha < 90^\circ$.

Similarly, we get $\beta < 90^\circ$ and $\gamma < 90^\circ$. Conversely, if α, β and γ are all acute angles, we may prove that there exists a point P such that $\angle APB, \angle ABC, \angle CPA$ are right angles.

(b) We may without loss of generality assume that $PA \geq PB \geq PC$, so $PA = d$. Because $AB^2 = AP^2 + BP^2 \geq AP^2 + CP^2 = AC^2$ and $AC^2 = AP^2 + CP^2 \geq BP^2 + CP^2 = BC^2$, we have $AB \geq AC \geq BC$.

Let H be the foot of the perpendicular from A to BC , then AH is the longest altitude of $\triangle ABC$, so $AH = h$. As

$AP \perp BP$ and $AP \perp CP$, AP is perpendicular to the plane of BPC . Thus $AP \perp BC$ and $AP \perp PH$ so that $AP < AH$, i.e., $d < h$ (1)

Because $AP \perp BP$ and $AH \perp BC$, we get BC is perpendicular to the plane of APH . Thus, we have $BC \perp PH$.

Let M be the midpoint of BC , then $PH \leq PM$. As

$\angle BPC = 90^\circ$, we have $PM = BM = MC = \frac{1}{2}BC$. Hence, $2PH \leq 2PM = BC$, so that

$$4PH^2 \leq BC^2 = PB^2 + PC^2 \leq 2PA^2 \quad \dots(2)$$

As $\angle APB = 90^\circ$, we get

$$PH^2 = AH^2 - AP^2 = h^2 - d^2 \quad \dots(3)$$

From (2) and (3), we have

$$4(h^2 - d^2) \leq 2d^2, \text{ i.e., } 2h^2 \leq 3d^2,$$

from which, we have

$$\frac{\sqrt{6}}{3}h \leq d. \quad \dots(4)$$

From (1) and (4) we obtain $\frac{\sqrt{6}}{3}h \leq d < h$, as required.

2. Because opposite angles of a cyclic quadrilateral are supplementary, we have that $\angle PBA = \pi - \angle ABS = \angle ARS$. Similarly $\angle PAB = \angle BSR$. Thus ΔPAB and ΔPSR are similar, from which

$$\frac{PA}{PS} = \frac{PB}{PR} = \frac{AB}{RS}$$

(Notice that this also gives the 'power of the point' result for P , $PA \cdot PR = PB \cdot PS$)

$$\text{Similarly } \frac{P'A}{P'S} = \frac{P'B}{P'R'} = \frac{AB}{RS'}$$

Consider now triangles APS and $AP'S'$. We have $\angle APS = \angle APB = \angle A'PB = \angle AP'S'$, because P , P' lie on the same arc of chord AB of the one circle. From the fact that S and S' lie on the same arc of chord AB of the second circle $\angle AS'P' = \angle ASP$. But then ΔAPS and $\Delta AP'S'$ are similar. Thus

$$\frac{PA}{PS} = \frac{P'A}{P'S'} \text{ So } \frac{AB}{RS} = \frac{AB}{RS'}$$

Thus $RS = RS'$ and the arcs are equal.

3. Using De Moivre's theorem
 $\cos n\theta + i \sin n\theta = (\cos \theta + i \sin \theta)^n$,
one finds easily that

$$\sin n\theta = \sum_{k=0}^{\left[\frac{n-1}{2}\right]} (-1)^k \binom{n}{2k+1} \cos^{n-2k-1} \theta \sin^{2k+1} \theta$$

$$\text{So } \sin(2n+1)\theta = \sum_{k=0}^n (-1)^k \binom{2n+1}{2k+1} \cos^{2n-2k} \theta \sin^{2k+1} \theta$$

$$= \tan \theta \cos^{2n+1} \theta \sum_{k=0}^n (-1)^k \binom{2n+1}{2k+1} \tan^{2k} \theta.$$

$$\text{Thus } \sum_{k=0}^n (-1)^k \binom{2n+1}{2k+1} \tan^{2k} \theta = 0$$

$$\text{For } \theta = \frac{j\pi}{2n+1}, 1 \leq j \leq n$$

$$\text{So } \tan^2 \frac{j\pi}{2n+1}, 1 \leq j \leq n, \text{ are the roots of}$$

$$\sum_{k=0}^n (-1)^k \binom{2n+1}{2k+1} x^k = 0 \text{ and thus also of}$$

$$\sum_{k=0}^n (-1)^k \binom{2n+1}{2k} x^{n-k} = 0. \text{ Since } a_n \text{ and } b_n \text{ are the sum}$$

and product of the roots, respectively, we have

$$a_n = \binom{2n+1}{2} = n(2n+1) \text{ and } b_n = \binom{2n+1}{2n} = 2n+1,$$

$$\text{and so } \frac{a_n}{b_n} = n$$

4. For the five rings, we have

$$a + b = b + c + d = d + e + f = f + g + h \\ = h + i = N. \quad \dots(1)$$

Since we are dealing with the nine non-zero decimal digits, we have $\sum_{j=1}^9 j = 9(10)/2 = 45$. The five regions sum to a common N for $45/5 = 9$ but then one pair must be $9 + 0$ or one triplet $9 + 0 + 0$, which isn't allowed. So $N > 9$. Since $a + b = h + i$, there must be at least two pairs of decimal digits that sum to N . For $10 \leq N \leq 15$, we have

$$N = 9 + a = 8 + (1 + a) = \dots, \text{ for } 1 \leq a \leq 6$$

Your favourite MTG Books/Magazines available in MANIPUR at

Fame Book House - Imphal Ph: 9774220348, 7085556697

Jain Book Shop - Imphal Ph: 9856031157

Job Centre - Imphal Ph: 0385-2440140; 9856700700, 9436204282

P.C. Jain And Co. Ph: 0385-2451758, 2225004; 985604649

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call 0124-6601200 for further assistance.

while $N = 16 = 9 + 17$ and $N = 17 = 9 + 8$ only. So $N \leq 15$.

From (1), $a + b = b + c + d$ or $a = c + d$... (2)

and $h + i = f + g + h$ or $i = f + g$... (3)

The five central digits must equal $45 - 2N$

$$(c + d) + e + (f + g) = a + e + i = 45 - 2N$$

So, we have

N	$2N$	$45 - 2N$	a, e, i
10	20	25	9, 8 - no digit available
11	22	23	9, 8, 6, ...
12	24	21	9, 8, 4, ...
13	26	19	9, 8, 2, ...
14	28	17	9, 7, 1, ...
15	30	15	9, 5, 1, ...

So $11 \leq N \leq 15$.

5. Assume that $m + n = 4k$, $m^2 + mn + n^2 = 49k$. Then $m^2 + 2mn + n^2 = (m + n)^2 = (4k)^2 = 16k^2$,

hence $mn = 16k^2 - 49k$. Since $mn > 0$, from $16k^2 - 49k > 0$ we find that $k > 3$. Since we also have the

$$\text{identity } mn = \left(\frac{m+n}{2}\right)^2 - \left(\frac{m-n}{2}\right)^2,$$

$$\text{where } \left(\frac{m-n}{2}\right)^2 \geq 0,$$

we also find that


$$0 \leq \left(\frac{m+n}{2}\right)^2 - mn = \left(\frac{4k}{2}\right)^2 - (16k^2 - 49k)$$

$= 49k - 12k^2$, from which $k \leq 4$ follows.

Hence $k = 4$ (since it must be an integer), and from $m + n = 16$ and $mn = 6$, $n = 10$, and $m^2 + mn + n^2 = 196$ follow. Indeed,

$$\frac{m+n}{m^2+mn+n^2} = \frac{16}{196} = \frac{4}{49}$$

6.



Let the circumcircle of the equilateral $\triangle ABC$ have centre $O(0, 0)$ and radius r . Join A to D , the mid-point of BC , then AD passes through O and is perpendicular to BC . Draw OB . Let the chord LM cut the sides AB and AC of $\triangle ABC$ at X and Y . Then $LXYM$ is parallel to BC .

$$OD = \tan 30^\circ = \frac{1}{\sqrt{3}}$$

$$\text{Then } D = \left(0, -\frac{1}{\sqrt{3}}\right),$$

$$A = \left(0, \frac{2}{\sqrt{3}}\right) \text{ and}$$

$$B = \left(-1, -\frac{1}{\sqrt{3}}\right) \text{ and the equation of the circle is}$$

$$x^2 + y^2 = \frac{4}{3}.$$

Now X is the mid-point of AB so

$$X = \left(-\frac{1}{2}, \frac{1}{2}\left(\frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}}\right)\right) = \left(-\frac{1}{2}, -\frac{1}{2\sqrt{3}}\right)$$

To find the x -coordinates of L and M , substitute the y -coordinate of X in the equation $x^2 + y^2 = \frac{4}{3}$, i.e.

$$x^2 + \left(\frac{1}{2\sqrt{3}}\right)^2 = \frac{4}{3}$$

$$\Rightarrow x^2 = \frac{4}{3} - \frac{1}{12} = \frac{5}{4}$$

$$\Rightarrow x = \pm \frac{\sqrt{5}}{2}$$

Thus the x -coordinate of L is $-\frac{\sqrt{5}}{2}$ and the x -coordinate of M is $\frac{\sqrt{5}}{2}$, so the length of LM is $\frac{\sqrt{5}}{2} + \frac{\sqrt{5}}{2} = \sqrt{5}$.

7. Since there are 8 teams, there are 7 rounds of four matches and thus a total of $7 \times 8 = 56$ points available. Consider a team with 10 points. It is possible to have 5 teams on 10 points and 3 teams on 2 points when each of the top 5 draws with each other, each of the bottom 3 draws with each other and each of the top 5 wins against each of the bottom 3. So, 10 points does not guarantee a place in the top 4.

Consider a team with 11 points. If this team was fifth, then the number of points gained by the top 5 teams is ≥ 55 . This is impossible as the number of points shared by the bottom 3 teams is then 1, as these 3 teams must have at least $3 \times 2 = 6$ points between them for the games played between themselves. Hence 11 points is sufficient to ensure a place in the top 4. Thus 11 points are required.



YOU ASK WE ANSWER

Do you have a question that you just can't get answered?

Use the vast expertise of our MTG team to get to the bottom of the question. From the serious to the silly, the controversial to the trivial, the team will tackle the questions, easy and tough. The best questions and their solutions will be printed in this column each month.

- Find the angle of intersection of curves, $y = [\sin x] + [\cos x]$ and $x^2 + y^2 = 5$, where $[\cdot]$ denotes the greatest integer function.

(Neeraj, Delhi)

Ans. We know that, $1 \leq |\sin x| + |\cos x| \leq \sqrt{2}$
 $\Rightarrow y = [|\sin x| + |\cos x|] = 1$
 Let P and Q be the points of intersection of given curves.
 Clearly, the given curves meet at points where $y = 1$, so we get $x^2 + 1 = 5$
 $\Rightarrow x = \pm 2$
 $\therefore P(2, 1)$ and $Q(-2, 1)$

On differentiating $x^2 + y^2 = 5$ w.r.t. x , we get

$$2x + 2y \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{x}{y}$$

$$\left(\frac{dy}{dx}\right)_{(2,1)} = -2 \text{ and } \left(\frac{dy}{dx}\right)_{(-2,1)} = 2$$

Clearly, the slope of line $y = 1$ is zero and the slope of the tangents at P and Q are (-2) and (2) , respectively.

Thus, the angle of intersection is $\tan^{-1}(2)$.

- Evaluate: $\int \frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} dx$

(Varnika, U.P.)

Ans. Let

$$\begin{aligned} I &= \int \frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} dx = \int \frac{\frac{\pi}{2} - 2 \cos^{-1} \sqrt{x}}{\frac{\pi}{2}} dx \\ &= \int \left(1 - \frac{4}{\pi} \cos^{-1} \sqrt{x}\right) dx = x - \frac{4}{\pi} \int 1 \cdot \cos^{-1} \sqrt{x} dx \\ &= x - \frac{4}{\pi} \left[\cos^{-1} \sqrt{x} \cdot x - \int x \cdot \frac{-1}{\sqrt{1-(\sqrt{x})^2}} \cdot \frac{1}{2\sqrt{x}} dx \right] \end{aligned}$$

$$\therefore I = x - \frac{4x}{\pi} \cos^{-1} \sqrt{x} - \frac{2}{\pi} \int \frac{\sqrt{x}}{\sqrt{1-x}} dx$$

Put $x = \cos^2 \theta$; then $dx = -2 \cos \theta \cdot \sin \theta d\theta$

$$\therefore \int \frac{\sqrt{x}}{\sqrt{1-x}} dx = \int \frac{\cos \theta}{\sqrt{1-\cos^2 \theta}} (-2 \cos \theta \cdot \sin \theta) d\theta$$

$$= -2 \cos^2 \theta d\theta = - \int (1 + \cos 2\theta) d\theta = - \left\{ \theta + \frac{\sin 2\theta}{2} \right\} + c$$

$$= -\cos^{-1} \sqrt{x} - \sqrt{x} \cdot \sqrt{1-x} + c$$

$$\therefore I = x - \frac{4x}{\pi} \cos^{-1} \sqrt{x} - \frac{2}{\pi} \left\{ -\cos^{-1} \sqrt{x} - \sqrt{x} \sqrt{1-x} + c \right\}$$

$$= x + \frac{2}{\pi} (1-2x) \cos^{-1} \sqrt{x} + \frac{2}{\pi} \sqrt{x(1-x)} + c_1, \text{ where } c_1 = -\frac{2c}{\pi}$$

- Find the general equation of the circle passing through two given points $A(x_1, y_1)$ and $B(x_2, y_2)$.

(Tanvi Sharma, Kerala)

Ans. Let $P(h, k)$ be any point on the circle passing through points $A(x_1, y_1)$ and $B(x_2, y_2)$. Since the angle in the same segment of a circle is always same.

Therefore, $\angle APB = 0$ or $\pi - 0$, where 0 is some angle.

$$\text{Now, } m_1 = \text{Slope of } AP = \frac{k-y_1}{h-x_1},$$

$$\text{and, } m_2 = \text{Slope of } BP = \frac{k-y_2}{h-x_2}$$

$$\tan \theta = \pm \frac{m_1 - m_2}{1 + m_1 m_2} \Rightarrow \tan \theta = \pm \frac{\frac{k-y_1}{h-x_1} - \frac{k-y_2}{h-x_2}}{1 + \frac{k-y_1}{h-x_1} \times \frac{k-y_2}{h-x_2}}$$

$$\Rightarrow \tan \theta = \pm \frac{(h-x_2)(k-y_1) - (h-x_1)(k-y_2)}{(h-x_1)(h-x_2) + (k-y_1)(k-y_2)}$$

$$\Rightarrow (h-x_1)(h-x_2) + (k-y_1)(k-y_2) \\ = \pm \cot \theta \{(h-x_2)(k-y_1) - (h-x_1)(k-y_2)\}$$

Hence, the locus of (h, k) is

$$(x-x_1)(x-x_2) + (y-y_1)(y-y_2) \\ = \pm \cot \theta \{(x-x_2)(y-y_1) - (x-x_1)(y-y_2)\}$$

$$\Rightarrow (x-x_1)(x-x_2) + (y-y_1)(y-y_2) \\ \pm \cot \theta \{x(y_1-y_2) + y(x_2-x_1) + x_1y_2 - x_2y_1\} = 0$$

$$\Rightarrow (x-x_1)(x-x_2) + (y-y_1)(y-y_2) \pm \cot \theta \begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{vmatrix} = 0$$



CONCEPT MAP

SEQUENCES AND SERIES

- Sum of n terms of special series**
- Sum of n natural numbers, $\sum n = \frac{n(n+1)}{2}$
- Sum of squares of n natural numbers, $\sum n^2 = \frac{n(n+1)(2n+1)}{6}$
- Sum of cubes of n natural numbers, $\sum n^3 = \frac{n^2(n+1)^2}{4} = (\sum n)^2$

Basic Properties

- If a constant is added / subtracted / multiplied / divided to each term of an A.P., then the resulting sequence is also an A.P.
- Selection of terms in an A.P.
 - Any three numbers in A.E. can be taken as $a-d, a, a+d$.
 - Any four numbers in A.P. can be taken as $a-3d, a-d, a+d, a+3d$.
- If each term of a G.P. is multiplied / divided by a same non-zero number, then the resulting sequence is also in G.P.
- Selection of terms in a G.P.
 - Any three numbers in G.P. can be taken as ar^2, ar, ar^{-2} .
 - Any four numbers in G.E. can be taken as $ar^3, ar, ar^{-1}, ar^{-3}$.

Arithmetic Mean (A.M.)

- For two numbers a and b , A.M. is $\frac{a+b}{2}$.
- $A_k = a + k\left(\frac{b-a}{n+1}\right), \forall k = 1, 2, \dots, n$

Where A_1, A_2, \dots, A_n are n arithmetic means inserted between two numbers a and b .

Geometric Mean (G.M.)

- For two numbers a and b , G.M. is \sqrt{ab} .
- $G_k = a\left(\frac{b}{a}\right)^{\frac{k-1}{n}}$ $\forall k = 1, 2, \dots, n$

Where G_1, G_2, \dots, G_n are n geometric means inserted between two numbers a and b .

Harmonic Mean (H.M.)

- For two numbers a and b , H.M. is $2ab/(a+b)$.
 - $H_n = \frac{(n+1)ab}{ab+n} \quad \forall n = 1, 2, \dots, n$
- Where H_1, H_2, \dots, H_n are n harmonic means inserted between them.

SERIES

If a_1, a_2, \dots, a_n is a sequence, then the expansion $a_1 + a_2 + \dots + a_n + \dots$ is called the series.

SEQUENCE

A sequence is a function from natural number N (domain) to real numbers (codomain).

Progression

If the terms of a sequence are written under specific conditions, then the sequence is called progression.

Types

Arithmetic Progression (A.P.)

A sequence whose terms increases or decreases by a fixed number.

- n^{th} term : $T_n = a + (n-1)d$, where d = common difference = $T_2 - T_1, d =$ first term.
- n^{th} term from end : $T_n' = l - (n-1)d$, where l = last term
- Sum of n terms : $S_n = \frac{n}{2}[2a + (n-1)d] = \frac{n}{2}[a + l]$

Geometric Progression (G.P.)

A sequence of non-zero numbers for which the ratio of a term to its just preceding term is always constant.

- n^{th} term : $T_n = ar^{n-1}$,
where r (common ratio) = $T_2/T_{1-1}, a$ = first term.
- n^{th} term from end : $T_n' = br^{(n-1)}$, b = last term

- Sum of n terms : $S_n = \begin{cases} a(r^n - 1) & r > 1 \\ \frac{a(1-r^n)}{1-r} & r < 1 \\ 1-r & r=1 \end{cases}$

Note : If $|r| \geq 1$, S_∞ does not exist.

Harmonic Progression (H.P.)

A sequence a_1, a_2, a_3, \dots in which reciprocal of terms form an A.P.

- n^{th} term : $T_n = \frac{1}{\frac{1}{a_1} + (n-1)\left(\frac{1}{a_2} - \frac{1}{a_1}\right)} = \frac{a_1 a_2}{a_2 + (n-1)(a_1 - a_2)}$

Note : No term of an H.P. can be zero.

Class XI Class XII

DEFINITE INTEGRALS AND APPLICATION OF INTEGRALS

Fundamental Theorems of Calculus

- First Fundamental Theorem :** Let $f(x)$ be a continuous function on the closed interval $[a, b]$ and let $A(x)$ be the area function. Then $A'(x) = f(x)$, for all $x \in [a, b]$.
- Second Fundamental Theorem :** Let $f(x)$ be a continuous function on the closed interval $[a, b]$ and $F(x)$ be integral of $f(x)$, then $\int_a^b f(x)dx = [F(x)]_a^b = F(b) - F(a)$

Solving by Substitution

When definite integral is to be solved by substitution, change the lower and upper limits of integration. If substitution is $t = f(x)$ and lower limit of integration is a and upper limit is b , then new lower and upper limits will be $f(a)$ and $f(b)$ respectively.

Limit of Sum

$$\int_a^b f(x)dx = \lim_{h \rightarrow 0} [f(a) + f(a+h) + \dots + f(a+(n-1)h)],$$

where $h = \frac{b-a}{n} \rightarrow 0$ as $n \rightarrow \infty$

Area Under Simple Curves




- Area = $\int_a^b y dx$

$$= \int_a^b f(x)dx \quad (\text{where } b > a)$$

- Area = $\int_a^b x dy$

$$= \int_a^b g(y)dy \quad (\text{where } b > a)$$

- Area = $\int_a^b [f(x) - g(x)] dx$



CONCEPT MAP

Properties

- $\int_a^b f(x)dx = \int_a^b f(t)dt$
- $\int_a^b f(x)dx = - \int_b^a f(x)dx$
In particular $\int_a^a f(x)dx = 0$
- $\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$,
where $a < c < b$
- $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$
- $\int_0^a f(x)dx = \int_0^a f(a-x)dx$
- $\int_a^a f(x)dx = \begin{cases} 0, & \text{if } f(x)=f(a) \\ 2 \int_a^a f(x)dx, & \text{if } f(-x)=f(x) \end{cases}$
- $\int_0^{2a} f(x)dx = \int_0^a f(x)dx + \int_a^{2a} f(2a-x)dx$
- $\int_0^{2a} f(x)dx = \begin{cases} \int_0^a f(x)dx, & \text{if } f(2a-x)=f(x) \\ 0, & \text{if } f(2a-x)=-f(x) \end{cases}$

DEFINITE INTEGRALS

APPLICATION OF INTEGRALS

Area Between Two Curves

- Area = $\int_a^b [f(x) - g(x)] dx$,
 $f(x) \geq g(x)$ in $[a, b]$
- Area = $\int_a^b [f(x) - g(x)] dx$
 $+ \int_a^b [g(x) - f(x)] dx$,
 $g(x) \geq f(x)$ in $[a, b]$
- Area = $\int_a^b f(x)dx + \int_c^b g(x)dx$,
 $y = f(x)$ and $y = g(x)$ intersect at $x = c$ and $x = b$

JEE WORK CUTS

One or More Than One Option(s) Correct Type

- Let A, B and C be three angles such that $A = \frac{\pi}{4}$ and $\tan B \tan C = p$. The set of all possible values of p such that A, B, C are the angles of a triangle contains
 (a) $(-\infty, 0)$ (b) $(0, 1)$
 (c) $(1, 3+2\sqrt{2})$ (d) $[3+2\sqrt{2}, \infty)$
- Let ABC be a triangle with $\angle BAC = 120^\circ$ and $AB \cdot AC = 1$. Also, let AD be the length of the angle bisector of $\angle A$ of the triangle. Then
 (a) Minimum value of AD is $\frac{1}{2}$
 (b) Maximum value of AD is $\frac{1}{2}$
 (c) AD is minimum when ΔABC is isosceles
 (d) AD is maximum when ΔABC is isosceles
- 8 players P_1, P_2, \dots, P_8 of equal strength play in a knockout tournament. Assuming that players in each round are paired randomly, the probability that the player P_1 loses to the eventual winner is
 (a) $\frac{1}{8}$ (b) $\frac{3}{8}$ (c) $\frac{5}{8}$ (d) $\frac{7}{8}$
- If z_1 and z_2 are two complex numbers such that $|z_1| = 2$ and $(1-i)z_2 + (1+i)\bar{z}_2 = 8\sqrt{2}$, then the minimum value of $|z_1 - z_2|$ is
 (a) 1 (b) 2 (c) 3 (d) 4
- Let L_1 and L_2 be the lines $\vec{r} = (2\hat{i} + \hat{j} - \hat{k}) + \lambda(\hat{i} + 2\hat{k})$ and $\vec{r} = (3\hat{i} + \hat{j}) + \mu(\hat{i} + \hat{j} - \hat{k})$. If the plane π which contains L_1 and parallel to L_2 meets the coordinate axes at A, B and C respectively, then the volume of the tetrahedron $OABC$ is
 (a) $\frac{4}{9}$ (b) $\frac{4}{3}$ (c) $\frac{2}{9}$ (d) $\frac{2}{3}$

- Which of the following statement(s) is/are correct?
 (a) Rolle's theorem is applicable to the function $F(x) = 1 - \sqrt[5]{x^6}$ on the interval $[-1, 1]$.
 (b) The domain of definition of the function $F(x) = \frac{\log_4(6-[x]-[x]^2)}{x^2+x-2}$ is $(-3, -2) \cup (-2, 1) \cup (1, 2)$.
 (where $[x]$ denotes the largest integer less than or equal to x)
 (c) The value of a for which the function $F(\theta) = a \sin \theta + \frac{1}{3} \sin 3\theta$ has an extremum at $\theta = \pi/3$ is -2 .
 (d) The value of $\sum_{k=1}^{2010} \frac{\{x+k\}}{2010}$ is $\{x\}$.
 (where $\{x\}$ denotes the fractional part of x).
- Let a, b, c be distinct complex numbers with $|a|=|b|=|c|=1$ and z_1, z_2 be the roots of the equation $az^2 + bz + c = 0$ with $|z|=1$. Let P and Q represent the complex numbers z_1 and z_2 in the argand plane with $\angle POQ = \theta$, $0^\circ < \theta < 180^\circ$ (where O being the origin), then
 (a) $b^2 = ac$; $\theta = \frac{2\pi}{3}$ (b) $\theta = \frac{2\pi}{3}$; $PQ = \sqrt{3}$
 (c) $PQ = 2\sqrt{3}$; $b^2 = ac$ (d) $\theta = \frac{\pi}{3}$; $b^2 = ac$
- Which of the following statement (s) is/are true?
 (a) Maximum value of P such that 3^P divides $100!$ is 48.
 (b) Maximum value of P such that 3^P divides $50!$ is 22.

- (c) Maximum value of P such that 3^P divides 25! is 10.
 (d) none of these
9. In a ΔABC , if $A = (1, 2)$ and internal angle bisectors through B and C are $y = x$ and $y = -2x$. The inradius of ΔABC is equal to
 (a) $\frac{1}{\sqrt{3}}$ (b) $\frac{2}{3}$ (c) $\frac{1}{3}$ (d) $\frac{1}{\sqrt{2}}$
10. If $I_n = \int_0^n x^n \tan^{-1} x dx$ and $a_n I_{n+2} + b_n I_n = c_n$
 $\forall n \in N, n \geq 1$, then
 (a) a_1, a_2, a_3, \dots are in A.P.
 (b) b_1, b_2, b_3, \dots are in A.P.
 (c) c_1, c_2, c_3, \dots are in G.P.
 (d) a_1, a_2, a_3, \dots are in H.P.
11. The equation of a circle is $S_1 \equiv x^2 + y^2 = 1$. The orthogonal tangents to S_1 meet at another circle S_2 and the orthogonal tangents to S_2 meet at the third circle S_3 . Then
 (a) radius of S_2 and S_3 are in the ratio 1 : $\sqrt{2}$
 (b) radius of S_2 and S_3 are in the ratio 1 : 2
 (c) the circles S_1, S_2 and S_3 are concentric
 (d) none of these
12. Tangent is drawn at any point (x_1, y_1) other than the vertex on the parabola $y^2 = 4ax$. If tangents are drawn from any point on this tangent to the circle $x^2 + y^2 = a^2$ such that all the chords of contact pass through a fixed point (x_2, y_2) , then
 (a) x_1, a, x_2 are in G.P.
 (b) $\frac{y_1}{2}, a, y_2$ are in G.P.
 (c) $-4, \frac{y_1}{y_2}, \frac{x_1}{x_2}$ are in G.P.
 (d) $x_1 x_2 + y_1 y_2 = a^2$
13. If $\log_2(5^{2-x} + 1), \log_4(2^{1-x} + 1)$ and 1 are in A.P., then x is equal to
 (a) $\frac{\log 5}{\log 2}$ (b) $\log_2(0.4)$
 (c) $1 - \frac{\log 5}{\log 2}$ (d) $\frac{\log 2}{\log 5}$
14. If $\int \operatorname{cosec} 2x dx = f\{g(x)\} + C$, then
 (a) range of $g(x) = (-\infty, \infty)$
 (b) domain of $f(x) = (-\infty, \infty) - \{0\}$
 (c) $g'(x) = \sec^2 x$
 (d) $f'(x) = \frac{1}{x}$ for all $x \in (0, \infty)$
15. Six persons stand at random in a queue for buying cinema tickets. Individually three of them have only a fifty rupee note each while each of the other three have a hundred rupee note only. The booking clerk has an empty cash box, probability that six persons get tickets without waiting for change is, (cost of one ticket is ₹50 and each person gets one ticket only)
 (a) $\frac{1}{2}$ (b) $\frac{1}{3}$ (c) $\frac{1}{4}$ (d) $\frac{1}{5}$
16. The mean and variance of seven observations are 8 and 16, respectively. If 5 observations are given by 2, 4, 10, 12, 14, then the product of the remaining two observations is
 (a) 45 (b) 48 (c) 40 (d) 49
17. A point on the straight line, $3x + 5y = 15$ which is equidistant from the coordinate axes will lie only in
 (a) 1st, 2nd and 4th quadrants
 (b) 1st and 2nd quadrants
 (c) 4th quadrant
 (d) 1st quadrant
18. The sum of the squares of the lengths of the chords intercepted on the circle, $x^2 + y^2 = 16$, by the lines, $x + y = n$, $n \in N$, where N is the set of all natural numbers, is
 (a) 160 (b) 105 (c) 210 (d) 320
19. If $f(x) = \log_e \left(\frac{1-x}{1+x} \right)$, $|x| < 1$, then $f\left(\frac{2x}{1+x^2}\right)$ is equal to
 (a) $2f(x^2)$ (b) $-2f(x)$ (c) $(f(x))^2$ (d) $2f(x)$
20. If $2y = \left(\cot^{-1} \left(\frac{\sqrt{3} \cos x + \sin x}{\cos x - \sqrt{3} \sin x} \right) \right)^2$, $x \in \left(0, \frac{\pi}{2}\right)$
 then $\frac{dy}{dx}$ is equal to
 (a) $\frac{\pi}{6} - x$ (b) $2x - \frac{\pi}{3}$ (c) $x - \frac{\pi}{6}$ (d) $\frac{\pi}{3} - x$

Comprehension Type

Paragraph for Q. No. 21 to 23

Let \vec{r} is a position vector of a variable point in cartesian OXY plane such that $\vec{r} \cdot (10\hat{j} - 8\hat{i} - \vec{r}) = 40$ and $p_1 = \max \{|\vec{r} + 2\hat{i} - 3\hat{j}|^2\}$, $p_2 = \min \{|\vec{r} + 2\hat{i} - 3\hat{j}|^2\}$. A tangent line is drawn to the curve $y = \frac{8}{x^2}$ at the point A with abscissa 2. The drawn line cuts x -axis at a point B .

21. p_2 is equal to

- (a) 9 (b) $2\sqrt{2}-1$
(c) $6\sqrt{2}+3$ (d) $9-4\sqrt{2}$

22. $p_1 + p_2$ is equal to

- (a) 2 (b) 10 (c) 18 (d) 5

23. $\overline{AB} \cdot \overline{OB}$ is

- (a) 1 (b) 2 (c) 3 (d) 4

Paragraph for Q. No. 24 to 26

For $n \in N$, we put $(1+x+x^2)^n = \sum_{r=0}^{2n} a_r x^r$

24. Value of $2(a_0 + a_1 + \dots + a_{n-1}) + a_n$ is
(a) 2^{2n-1} (b) 3^n (c) $3^{n/2}$ (d) $(3^n-1)/2$
25. If $a_0^2 - a_1^2 + a_2^2 - a_3^2 + \dots + a_{2n}^2 = ka_n$, then k equals
(a) 1 (b) 2 (c) $1/2$ (d) 0

26. If n is not a multiple of 3, and

$$\sum_{r=0}^n (-1)^r a_r \binom{n}{[n/3]} = k \binom{n}{[n/3]}, \text{ where } [x] \text{ denotes the}$$

greatest integer $\leq x$, then k is equal to

- (a) 1 (b) 0 (c) 3 (d) -1

Paragraph for Q. No. 27 and 28

Consider the set of points (x, y) in the plane which satisfy $x^2 + y^2 \leq 100$ and $\sin(x+y) \geq 0$.

27. Let A_1 and A_2 be areas of regions within $x^2 + y^2 \leq 100$ which satisfy $\sin(x+y) > 0$ and $\sin(x+y) < 0$, then
(a) $A_1 > A_2$ (b) $A_1 < A_2$
(c) $A_1 = A_2$ (d) $A_1 = 2A_2$

28. The area of the region $x^2 + y^2 \leq 100$ and $\sin(x+y) \geq 0$ is
(a) 25π (b) 50π (c) 100π (d) 200π

Matrix Match Type

29. For $0 < \theta < \pi/4$, let $x = \sum_{n=0}^{\infty} (\sin \theta)^{2n}, y = \sum_{n=0}^{\infty} (\cos \theta)^{2n}$,

then match the following columns :

Column-I	Column-II
(A) $\sum_{n=0}^{\infty} \sin^{2n} \theta \cos^{2n} \theta$	(P) $\frac{xy^2}{xy^2 - 1}$
(B) $\sum_{n=0}^{\infty} \tan^{2n} \theta$	(Q) $\frac{y}{y-x}$
(C) $\sum_{n=0}^{\infty} \sin^{2n} \theta \cos^{4n} \theta$	(R) $\frac{xy}{xy-1}$
(D) $\sum_{n=0}^{\infty} \cos^{2n} \theta \sin^{4n} \theta$	(S) $\frac{x^2 y}{x^2 y - 1}$

- (a) (A) \rightarrow (Q); (B) \rightarrow (P); (C) \rightarrow (R); (D) \rightarrow (S)

- (b) (A) \rightarrow (S); (B) \rightarrow (P); (C) \rightarrow (Q); (D) \rightarrow (R)

- (c) (A) \rightarrow (R); (B) \rightarrow (Q); (C) \rightarrow (P); (D) \rightarrow (S)

- (d) (A) \rightarrow (S); (B) \rightarrow (Q); (C) \rightarrow (P); (D) \rightarrow (R)

30. Match the following columns :

Column-I	Column-II
(A) If the distance of any point (x, y) from origin is defined as $d(x, y) = 2 x + 3 y $. If perimeter and area of figure bounded by $d(x, y) = 6$ are λ units and μ sq. units respectively, then	(P) (λ, μ) lies on $x^2 - y^2 = 64$
(B) If the vertices of a triangle are $(6, 0), (0, 6)$ and $(6, 6)$. If the distance between circumcentre and orthocentre and distance between circumcentre and centroid are λ unit and μ unit respectively, then	(Q) (λ, μ) lies on $x^2 + y^2 - 6x - 6y = 0$
(C) The ends of the hypotenuse of a right angled triangle are $(6, 0)$ and $(0, 6)$. If the third vertex is (λ, μ) , then	(R) (λ, μ) lies on $x^2 - 16y = 16$
	(S) (λ, μ) lies on $x^2 - y^2 = 16$

- (a) (A) \rightarrow (P); (B) \rightarrow (S); (C) \rightarrow (Q)

- (b) (A) \rightarrow (P,R); (B) \rightarrow (R); (C) \rightarrow (P)

- (c) (A) \rightarrow (P); (B) \rightarrow (Q); (C) \rightarrow (R,S)

- (d) (A) \rightarrow (Q); (B) \rightarrow (P,Q); (C) \rightarrow (R)

31. Consider the circles C_1 of radius a and C_2 of radius b , $b > a$ both lying in the first quadrant and touching the coordinate axes. Find the value of b/a if

Column-I	Column-II
(A) C_1 and C_2 touch each other	(P) $2 + \sqrt{2}$
(B) C_1 and C_2 are orthogonal	(Q) 3
(C) C_1 and C_2 intersect so that the common chord is longest	(R) $2 + \sqrt{3}$
(D) C_2 passes through the centre of C_1	(S) $3 + 2\sqrt{2}$

- (a) (A) \rightarrow (S); (B) \rightarrow (P,S); (C) \rightarrow (Q); (D) \rightarrow (Q)

- (b) (A) \rightarrow (R); (B) \rightarrow (Q); (C) \rightarrow (P,S); (D) \rightarrow (Q)

- (c) (A) \rightarrow (P,Q); (B) \rightarrow (R); (C) \rightarrow (S); (D) \rightarrow (P)

- (d) (A) \rightarrow (S); (B) \rightarrow (R); (C) \rightarrow (Q); (D) \rightarrow (P)



Numerical Answer Type

32. Minimum distance between $y^2 - 4x - 8y + 40 = 0$ and $x^2 - 8x - 4y + 40 = 0$ is $\sqrt{\lambda}$, then value of λ is _____.
33. The value of $\lim_{x \rightarrow \infty} \left(\sqrt{x + \sqrt{x + \sqrt{x}}} - \sqrt{x} \right)$ is _____.
34. Number of pairs of positive integers (p, q) whose L.C.M. is 8100, is "K". Then number of ways of expressing K as a product of two co-prime numbers is _____.
35. In ΔABC , if $A - B = 120^\circ$ and $R = 8r$, then the value of $\frac{1 + \cos C}{1 - \cos C}$ equals (All symbols used have their usual meaning in a triangle) _____.
36. In ΔABC , orthocentre is $(6, 10)$ and circumcentre is $(2, 3)$ and equation of side BC is $2x + y = 17$. Then the radius of the circumcircle of ΔABC is _____.
37. If $f(x)$ and $g(x)$ are periodic functions with periods 7 and 11 respectively, then the period of $F(x) = f(x)g\left(\frac{x}{5}\right) - g(x)f\left(\frac{x}{3}\right)$ is _____.
38. How many different nine digit numbers can be formed from the number 22 33 55 888 by rearranging its digits, so that the odd digits occupy even positions?
39. A straight line L with negative slope passes through the point $(8, 2)$ and cuts the positive coordinate axes at points P and Q . As L varies, the absolute minimum value of $OP + OQ$ is (O is origin) _____.
40. For any real number x , let $[x]$ denote the largest integer less than or equal to x . Let f be a real-valued function defined on the interval $[-10, 10]$ by
- $$f(x) = \begin{cases} x - [x], & \text{if } [x] \text{ is odd} \\ 1 + [x] - x, & \text{if } [x] \text{ is even} \end{cases}$$
- Then the value of $\frac{\pi^2}{10} \int_{-10}^{10} f(x) \cos \pi x \, dx$ is _____.

SOLUTIONS

1. (a, d) : $\tan B \tan C = p$

$$\Leftrightarrow \frac{\cos(B-C)}{\cos(B+C)} = \frac{p+1}{1-p} \Leftrightarrow \cos(B-C) = \frac{p+1}{\sqrt{2}(p-1)}$$

$$\therefore 0 \leq (B-C) < \frac{3\pi}{4} \Leftrightarrow \frac{-1}{\sqrt{2}} < \frac{p+1}{\sqrt{2}(p-1)} \leq 1$$

$$\Leftrightarrow \frac{p+1}{p-1} + 1 = \frac{2p}{p-1} > 0 \text{ and } \frac{(p+1) - \sqrt{2}(p-1)}{\sqrt{2}(p-1)} \leq 0$$

$$\Leftrightarrow p(p-1) > 0 \text{ and } [(\sqrt{2} + 1) - (\sqrt{2} - 1)p](p-1) \leq 0$$

$$\Leftrightarrow p \notin [0, 1] \text{ and } [(\sqrt{2} + 1)^2 - p](p-1) \leq 0$$

$$\Leftrightarrow p \notin [0, 1] \text{ and } p \notin (1, 3 + 2\sqrt{2})$$

$$\Leftrightarrow p \notin (0, 3 + 2\sqrt{2}) \Leftrightarrow p < 0 \text{ or } p \geq 3 + 2\sqrt{2}$$

2. (b, d) : Let $AB = x$. Then $BC^2 = x^2 + \frac{1}{x^2} + 1$

and $\cos B = \frac{2x^2 + 1}{2\sqrt{x^4 + x^2 + 1}}$ i.e., $\tan B = \frac{\sqrt{3}}{2x^2 + 1}$

Also, $\frac{AD}{\sin B} = \frac{x}{\sin(B+60^\circ)}$

$$\Rightarrow AD = \frac{x}{\frac{1 + \sqrt{3}}{2} \cot B} = \frac{x}{x^2 + 1}$$

$$\therefore AD = \frac{1}{x + \frac{1}{x}} \leq \frac{1}{2}, \text{ with equality iff } AB = AC = 1$$

3. (b) : If E_1, E_2, E_3 are the events of P_1 losing to the champion in the 1st, 2nd and 3rd rounds, then the required probability

$$= P(E_1) + P(E_2) + P(E_3) = \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{3}{8}$$

4. (b) : $\operatorname{Re}(1-i)z_2 = 4\sqrt{2} \Rightarrow |(1-i)z_2| \geq 4\sqrt{2}$

$\Rightarrow |z_2| \geq 4 \Rightarrow z_2$ lies either on the circumference or outside the circle with centre at origin and radius 4.

$$\therefore |z_1 - z_2| \geq 2$$

5. (c) : Equation of the plane containing L_1 and parallel

to L_2 is
$$\begin{vmatrix} x-2 & y-1 & z+1 \\ 1 & 0 & 2 \\ 1 & 1 & -1 \end{vmatrix} = 0$$

$$\Rightarrow 2x - 3y - z - 2 = 0$$

This plane meets the axes at $A(1, 0, 0)$, $B(0, -2/3, 0)$ and $C(0, 0, -2)$

$$\therefore \text{Volume of tetrahedron } OABC = \frac{1}{6} |abc| = \frac{2}{9}$$

6. (a, d) : (a) We have, $F(x) = (1-x^{6/5})$

$$\text{Now, } F'(x) = \frac{-6}{5}x^{\frac{1}{5}} \text{ exist } \forall x \in (-1, 1)$$

$$\text{Also, } F(-1) = 0 = F(1)$$

Hence Rolle's theorem is applicable to the function $F(x)$.

(b) For domain of $F(x)$,

$$6 - [x] - [x]^2 > 0 \text{ and } x^2 + x - 2 \neq 0$$

$$\Rightarrow (x+2)(x-1) \neq 0 \Rightarrow x \neq -2, 1$$

$$\text{Now } [x]^2 + [x] - 6 < 0 \Rightarrow ([x] + 3)([x] - 2) < 0$$

$$\Rightarrow -3 < [x] < 2 \Rightarrow -2 \leq x < 2$$

∴ Domain = $(-2, 1) \cup (1, 2)$

$$(c) \text{ We have, } F(0) = a \sin \theta + \frac{1}{3} \sin 3\theta$$

As $F(0)$ has an extremum at $\theta = \frac{\pi}{3}$, so

$$a \cos \theta + \cos 3\theta = 0 \text{ at } \theta = \frac{\pi}{3} \Rightarrow \frac{a}{2} - 1 = 0$$

$$\Rightarrow \frac{a}{2} = 1 \Rightarrow a = 2$$

$$(d) \text{ We have } \sum_{k=1}^{2010} \frac{[x+k]}{2010} = \frac{[x+1]}{2010} + \frac{[x+2]}{2010} + \dots$$

$$+ \frac{[x+2010]}{2010} = \frac{2010[x]}{2010} = [x]$$

$$7. (a, b) : |z_1 + z_2| = \left| \frac{-b}{a} \right| ; z_1 z_2 = \left| \frac{c}{a} \right|^2$$

$$\therefore |z_1 + z_2|^2 = 1 \Rightarrow (z_1 + z_2)(\bar{z}_1 + \bar{z}_2) = 1$$

$$\Rightarrow 2 + \bar{z}_1 z_2 + \bar{z}_2 z_1 = 1 \Rightarrow \frac{(z_1 + z_2)^2}{z_1 z_2} = 1$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{c}{a} \Rightarrow b^2 = ac$$

Now, $z_2 = z_1 e^{i\theta}$, then $|z_1 + z_2| = |z_1| |1 + e^{i\theta}|$

$$\Rightarrow 2 \cos \frac{\theta}{2} = 1 \quad \therefore \theta = \frac{2\pi}{3}$$

$$\therefore PQ = |z_2 - z_1| = \sqrt{3}$$

$$8. (a, b, c) : (a) \left[\frac{100}{3} \right] + \left[\frac{100}{3^2} \right] + \left[\frac{100}{3^3} \right] + \left[\frac{100}{3^4} \right] \\ = 33 + 11 + 3 + 1 = 48$$

$$(b) \left[\frac{50}{3} \right] + \left[\frac{50}{3^2} \right] + \left[\frac{50}{3^3} \right] = 16 + 5 + 1 = 22$$

$$(c) \left[\frac{25}{3} \right] + \left[\frac{25}{3^2} \right] = 8 + 2 = 10$$

9. (d) Let image of A about $y = x$, $y = -2x$ be P and Q .

$$\therefore P = (2, 1), Q = \left(\frac{-11}{5}, \frac{2}{5} \right)$$

Equation of BC is $x - 7y + 5 = 0$

$$\therefore \text{In radius} = 1' \text{ distance from } I(0, 0) \text{ to } BC = \frac{1}{\sqrt{2}}$$

$$10. (a, b) : I_n = \left(\frac{x^{n+1}}{n+1} \tan^{-1} x \right)_0^1 - \int_0^1 \frac{x^{n+1}}{n+1} \frac{1}{1+x^2} dx$$

$$\Rightarrow (n+1)I_n = \frac{\pi}{4} - \int_0^1 \frac{x^{n+3}}{1+x^2} dx$$

$$\therefore (n+1)I_n + (n+3)I_{n+2} = \frac{\pi}{2} - \frac{1}{n+2}$$

$a_n = (n+3) \Rightarrow a_1, a_2, a_3, \dots$ are in A.P.
 $b_n = (n+1) \Rightarrow b_1, b_2, \dots$ are in A.P.

$c_n = \frac{\pi}{2} - \frac{1}{n+2} \Rightarrow c_1, c_2, \dots$ are not in any progression.

11. (a, c) : Orthogonal tangents to a circle meet at the director circle.

$$\therefore S_2 \equiv x^2 + y^2 = 2$$

$$\text{Also, } S_3 \equiv x^2 + y^2 = 4$$

∴ Ratio of radius of S_2 and $S_3 = \sqrt{2} : 2 = 1 : \sqrt{2}$

Also, the three circles are concentric.

12. (b, c, d) : Let $(x_1, y_1) = (at^2, 2at)$.

Tangent at this point is $ty = x + at^2$.

Any point on this tangent is $(h, (h + at^2)/t)$.

The chord of contact of this point with respect to the circle $x^2 + y^2 = a^2$ is

$$hx + \left(\frac{h+at^2}{t} \right) y = a^2 \Rightarrow (aty - a^2) + h \left(x + \frac{y}{t} \right) = 0,$$

which is a family of straight lines passing through the point of intersection of $ty - a = 0$ and $x + \frac{y}{t} = 0$

So, the fixed point is $(-a/t^2, a/t)$.

$$\text{Therefore, } x_2 = -\frac{a}{t^2}, y_2 = \frac{a}{t}$$

$$\text{Clearly, } x_1 x_2 = -a^2, y_1 y_2 = 2a^2$$

$$\text{Also, } \frac{x_1}{x_2} = -t^4 \text{ and } \frac{y_1}{y_2} = 2t^2 \text{ or } 4 \frac{x_1}{x_2} + \left(\frac{y_1}{y_2} \right)^2 = 0,$$

13. (b, c) : From the given condition, we have

$$2 \log_4 (2^{1-x} + 1) = \log_2 (5 \cdot 2^x + 1) + 1$$

$$\Rightarrow \frac{2 \log(2^{1-x} + 1)}{\log 4} = \frac{\log(5 \cdot 2^x + 1)}{\log 2} + 1$$

$$\Rightarrow \log(2^{1-x} + 1) = \log [(5 \cdot 2^x + 1) 2]$$


$$\Rightarrow 2^{1-x} + 1 = 10 \cdot 2^x + 2$$

$$\text{Put } 2^x = y, \text{ so that } \frac{2}{y} + 1 = 10y + 2 \Rightarrow 10y^2 + y - 2 = 0$$

The Best tool for success in **JEE Main**

19 Years' JEE Main Chapterwise-Topicwise Solutions Physics, Chemistry and Mathematics contain not only chapterwise - topicwise questions that have appeared over the last 19 years in JEE Main / AIEEE but also their complete solutions. Needless to say these books are essential for any student to compete successfully in JEE Main.

New



₹ 350



₹ 350



₹ 350

HIGHLIGHTS:

- Chapterwise-Topicwise Questions (Offline & Online) of last 19 years' (2000-2002) of JEE Main / AIEEE
- Chapterwise-Topicwise segregation of questions to help you assess the level of effort required to succeed
- An unmatched series with more than 1000 pages filled with detailed solutions by experts

Visit
www.mtg.in
 for latest offers
 and to buy
 online!



MTG Learning Media (P) Ltd.
 Plot #99, Sector 44, Gurugram – 122 003 (HR)

Available at all leading book shops throughout India.
 For more information or for help in placing your order,
 Call 0124-6601200 or e-mail info@mtg.in

$$\Rightarrow (5y - 2)(2y + 1) = 0 \Rightarrow y = 2/5 \text{ or } y = -1/2.$$

Since $y = 2^x$ cannot be negative,

$$\therefore 2^x = 2/5 = 0.4 \text{ or } x = \frac{\log(2/5)}{\log 2} = 1 - \frac{\log 5}{\log 2}.$$

14. (a, b, c) : $\int \operatorname{cosec} 2x dx = f(g(x)) + C$

$$\Rightarrow \operatorname{cosec} 2x = f'(g(x))g'(x)$$

$$\Rightarrow \frac{1}{2 \tan x} \times \sec^2 x = f'(g(x))g'(x)$$

$$\Rightarrow f(x) = \frac{1}{2x}, g'(x) = \sec^2 x$$

$$\text{Domain } f(x) = (-\infty, \infty) - \{0\}, g'(x) = \sec^2 x$$

$$\text{Range } g(x) = (-\infty, \infty)$$

15. (c) : Here random experiment of arranging 6 persons in a line is $n(S) = 6! = 720$

Let 'a' denotes the person having ₹50 note and 'b' denotes the person having ₹100 note each, since all the six persons, should get ticket. First place should be occupied by a person having ₹50 note and sixth place should be occupied by a person having ₹100 note. So, possible cases are

[a]	[aa]	[bb]	[b]
[a]	[ab]	[ab]	[b]
[a]	[ab]	[ba]	[b]
[a]	[ba]	[ab]	[b]
[a]	[ba]	[ba]	[b]

'a's can arrange among themselves and 'b's can arrange among themselves is $n(E) = 3!3! + 3!3! + 3!3! + 3!3! + 3!3! = 180$ ways

$$\therefore \text{Required probability} = \frac{180}{720} = \frac{1}{4}$$

16. (b) : Let the remaining two observations are x_1 and x_2 .

$$\therefore \text{Mean } (\bar{x}) = \frac{2+4+10+12+14+x_1+x_2}{7} = 8$$

$$\Rightarrow x_1 + x_2 = 14 \quad \dots(i)$$

$$\text{And variance } (\sigma^2) = \frac{\sum x_i^2}{n} - (\bar{x})^2$$

$$\Rightarrow 16 = \frac{4+16+100+144+196+x_1^2+x_2^2}{7} - 64$$

$$\Rightarrow 80 = \frac{460+x_1^2+x_2^2}{7}$$

$$\Rightarrow x_1^2 + x_2^2 = 560 - 460 = 100$$

$$\Rightarrow (x_1 + x_2)^2 - 2x_1 x_2 = 100$$

$$\Rightarrow x_1 x_2 = \frac{14^2 - 100}{2}$$

[Using (i)]

$$\Rightarrow x_1 x_2 = 48$$

17. (b) : Let $P\left(t, \frac{15-3t}{5}\right)$ be any point on the straight line $3x + 5y = 15$.


$$\therefore \left| \frac{15-3t}{5} \right| = |t|$$

$$\Rightarrow \frac{15-3t}{5} = t \text{ or } \frac{15-3t}{5} = -t$$

$$\Rightarrow 15 - 3t = 5t \text{ or } 15 - 3t = -5t$$

$$\Rightarrow 15 = 8t \text{ or } 15 = -2t \Rightarrow t = \frac{15}{8} \text{ or } t = \frac{-15}{2}$$

∴ Point P lies in 1st or 2nd quadrants.




18. (c) : Let p be the perpendicular distance from $O(0, 0)$ to line $x + y = n$.

$$\therefore p = \frac{n}{\sqrt{2}} < 4 \text{ for } n = 1, 2, 3, 4, 5$$

Now, length of chord

$$= 2\sqrt{r^2 - p^2} = 2\sqrt{r^2 - \frac{n^2}{2}}$$

$$= 2\sqrt{16 - \frac{n^2}{2}} = \sqrt{64 - 2n^2}$$



Now, sum of the squares of the lengths of the chords $= 62 + 56 + 46 + 32 + 14 = 210$

19. (d) : Here, $f(x) = \log_e \left(\frac{1-x}{1+x} \right)$

$$\text{Now, } f\left(\frac{2x}{1+x^2}\right) = \log_e \left(\frac{1-\frac{2x}{1+x^2}}{1+\frac{2x}{1+x^2}} \right)$$

$$= \log_e \left(\frac{1+x^2-2x}{1+x^2+2x} \right) = \log_e \left(\frac{1-x}{1+x} \right)^2$$

$$= 2 \log_e \left(\frac{1-x}{1+x} \right) = 2f(x)$$

20. (c) : We have, $2y = \left(\cot^{-1} \left(\frac{\sqrt{3} \cos x + \sin x}{\cos x - \sqrt{3} \sin x} \right) \right)^2$

$$= \left[\cot^{-1} \left(\frac{\frac{\sqrt{3}}{2} \cos x + \frac{1}{2} \sin x}{\frac{1}{2} \cos x - \frac{\sqrt{3}}{2} \sin x} \right) \right]^2 = \left[\cot^{-1} \left(\frac{\sin \left(x + \frac{\pi}{3} \right)}{\cos \left(x + \frac{\pi}{3} \right)} \right) \right]^2$$

$$\begin{aligned}
&= \left[\cot^{-1} \left(\tan \left(x + \frac{\pi}{3} \right) \right) \right]^2 = \left[\frac{\pi}{2} - \tan^{-1} \left(\tan \left(x + \frac{\pi}{3} \right) \right) \right]^2 \\
&= \left[\frac{\pi}{2} - \left(x + \frac{\pi}{3} \right) \right]^2 = \left[\frac{\pi}{6} - x \right]^2 \\
\Rightarrow 2 \frac{dy}{dx} &= 2 \left(\frac{\pi}{6} - x \right) (-1) \quad \Rightarrow \frac{dy}{dx} = x - \frac{\pi}{6}
\end{aligned}$$

(21-23) : Let $\vec{r} = x\hat{i} + y\hat{j}$, $x^2 + y^2 + 8x - 10y + 40 = 0$, centre $C(-4, 5)$, radius = 1, $P_1 = \max\{(x+2)^2 + (y-3)^2\}$, $P_2 = \min\{(x+2)^2 + (y-3)^2\}$

Let $P \equiv (-2, 3)$ and M be any point on circle.

$$CP = 2\sqrt{2}, CM = 1$$

$$P_1 = (CP + CM)^2 = (2\sqrt{2} + 1)^2,$$

$$P_2 = (CP - CM)^2 = (2\sqrt{2} - 1)^2,$$

$$P_1 + P_2 = 18$$

$$\text{Now, slope of } AB = \left(\frac{dy}{dx} \right)_{(2, 2)} = -2$$

Equation of AB is $2x + y = 6$

$$\overline{OA} = 2\hat{i} + 2\hat{j}, \quad \overline{OB} = 3\hat{i}, \quad \overline{AB} = \hat{i} - 2\hat{j},$$

$$\overline{AB} \cdot \overline{OB} = (\hat{i} - 2\hat{j})(3\hat{i}) = 3$$

21. (d) 22. (c) 23. (c)

$$24. (b) : \text{Given, } (1 + x + x^2)^n = \sum_{r=0}^{2n} a_r x^r$$

Putting $x = 1$ in (i), we get

$$a_0 + a_1 + a_2 + \dots + a_{2n} = (1 + 1 + 1)^n = 3^n$$

But $a_r = a_{2n-r}$ for $0 \leq r \leq n - 1$.

$$\therefore 2(a_0 + a_1 + \dots + a_{n-1}) + a_n = 3^n$$

25. (a) : Replacing x by $-1/x$ in (i), we get

$$\left(1 - \frac{1}{x} + \frac{1}{x^2} \right)^n = \sum_{r=0}^{2n} (-1)^r \frac{a_r}{x^r}$$

Now, $a_0^2 - a_1^2 + a_2^2 - a_3^2 + \dots + a_{2n}^2$ = coefficient of the constant term in $(a_0 + a_1 x + \dots + a_{2n} x^{2n})$

$$\left[a_0 - \frac{a_1}{x} + \frac{a_2}{x^2} - \frac{a_3}{x^3} + \dots + \frac{a_{2n}}{x^{2n}} \right]$$

= coefficient of the constant term in

$$(1 + x + x^2)^n \left(1 - \frac{1}{x} + \frac{1}{x^2} \right)^n$$

= coefficient of the constant term in

$$\frac{(1 + x + x^2)^n (x^2 - x + 1)^n}{x^{2n}}$$

= coefficient of x^{2n} in $[(x^2 + 1)^2 - x^2]^n$

$$\begin{aligned}
&= \text{coefficient of } x^{2n} \text{ in } (1 + x^2 + x^4)^n \\
&= \text{coefficient of } y^n \text{ in } (1 + y + y^2)^n = a_n
\end{aligned}$$

26. (b) : $\sum_{r=0}^n (-1)^r a_r {}^n C_r$ = coefficient of the constant term in $[a_0 + a_1 x + a_2 x^2 + \dots + a_{2n} x^{2n}]$

$$\times \left[{}^n C_0 - {}^n C_1 \left(\frac{1}{x} \right) + {}^n C_2 \left(\frac{1}{x} \right)^2 + \dots + (-1)^n {}^n C_n \left(\frac{1}{x} \right)^n \right]$$

$$= \text{coefficient of the constant term in } (1 + x + x^2)^n \left(1 - \frac{1}{x} \right)^n$$

$$= \text{coefficient of } x^n \text{ in } (x^2 + x + 1)^n (x - 1)^n$$

$$= \text{coefficient of } x^n \text{ in } (x^3 - 1)^n$$

= 0 as n is not a multiple of 3.

(27-28) : $\sin(x+y) > 0$

$\Rightarrow x+y \in (0, \pi) \cup (2\pi, 3\pi)$ etc.

and $\sin(x+y) < 0$

$\Rightarrow x+y \in (\pi, 2\pi) \cup (3\pi, 4\pi)$ etc.

Hence, $A_1 = A_2$

and $A_1 + A_2$ = Full circle area

$$= 100\pi$$

Hence, $A_1 = A_2 = 50\pi$

27. (c) 28. (b)

$$29. (c) : \text{Here } x = \frac{1}{1 - \sin^2 \theta} \Rightarrow \cos^2 \theta = \frac{1}{x}$$

$$\text{and } y = \frac{1}{1 - \cos^2 \theta} \Rightarrow \sin^2 \theta = \frac{1}{y}$$

$$(A) \sum_{n=0}^{\infty} \sin^{2n} \theta \cos^{2n} \theta = \frac{1}{1 - \sin^2 \theta \cos^2 \theta}$$

$$= \frac{1}{1 - \frac{1}{xy}} = \frac{xy}{xy - 1}$$


$$(B) \sum_{n=0}^{\infty} \tan^{2n} \theta = \frac{1}{1 - \tan^2 \theta} = \frac{1}{1 - \frac{x}{y}} = \frac{y}{y-x}$$

$$(C) \sum_{n=0}^{\infty} \sin^{2n} \theta \cos^{4n} \theta = \frac{1}{1 - \sin^2 \theta \cos^4 \theta}$$

$$= \frac{1}{1 - \frac{1}{x^2 y}} = \frac{x^2 y}{x^2 y - 1}$$

$$(D) \sum_{n=0}^{\infty} \cos^{2n} \theta \sin^{4n} \theta = \frac{1}{1 - \cos^2 \theta \sin^4 \theta} = \frac{1}{1 - \frac{1}{xy^2}}$$

$$= \frac{xy^2}{xy^2 - 1}.$$



30. (a) : (A) $\because d(x, y) = 2|x| + 3|y| = 6$ (given)

$$\Rightarrow \frac{|x|}{3} + \frac{|y|}{2} = 1$$

∴ Perimeter,

$$\lambda = 4\sqrt{13}$$

and area,

$$\mu = 4 \times \frac{1}{2} \times 3 \times 2 = 12$$

$$\text{Thus, } \frac{\lambda^2}{16} - \mu^2 = 1 \text{ and } \lambda^2 - \mu^2 = 64$$

(B) It is clear that orthocentre is (6, 6)

Circumcentre $\equiv (3, 3)$ and centroid $\equiv (4, 4)$

$$\therefore \lambda = \sqrt{(6-3)^2 + (6-3)^2} \\ = \sqrt{9+9} = 3\sqrt{2}$$

$$\text{and } \mu = \sqrt{(4-3)^2 + (4-3)^2} \\ = \sqrt{1+1} = \sqrt{2}$$

$$\therefore \lambda^2 - \mu^2 = 16$$

(C) Slope of AC \times Slope of BC $= -1$

$$\Rightarrow \left(\frac{\mu-0}{\lambda-6}\right) \times \left(\frac{\mu-6}{\lambda-0}\right) = -1$$

$$\Rightarrow \mu^2 - 6\mu = -\lambda^2 + 6\lambda$$

$$\Rightarrow \lambda^2 + \mu^2 - 6\lambda - 6\mu = 0$$

31. (d) : (A) $C_1 : x^2 + y^2 - 2ax(x + y) + a^2 = 0$,

Centre : (a, a) , radius : a ,

$C_2 : x^2 + y^2 - 2b(x + y) + b^2 = 0$

Centre : (b, b) , radius b

Since C_1 and C_2 touch each other

$$\Rightarrow \sqrt{2}(b-a) = a+b \Rightarrow \frac{b}{a} = (\sqrt{2}+1)^2 = 3+2\sqrt{2}$$

(B) C_1 and C_2 intersect orthogonally

$$\Rightarrow 2(b-a)^2 = a^2 + b^2 \Rightarrow \frac{b}{a} = 2 + \sqrt{3}$$

(C) The common chord is

$$2(b-a)(x+y) = b^2 - a^2.$$


It passes through $(a, a) \Rightarrow b/a = 3$.

(D) C_2 passes through (a, a)

$$\Rightarrow 2a^2 - 4ab + b^2 = 0 \Rightarrow \frac{b}{a} = 2 + \sqrt{2}.$$

32. (2) : Since two parabolas are symmetrical about $y = x$.

Solving $y = x$ and $y^2 - 4x - 8y + 40 = 0$, we get $x^2 - 12x + 40 = 0$ has no real solution.



\therefore They don't intersect.

Point on $(x-4)^2 = 4(y-6)$ is (6, 7) and the corresponding point on $(y-4)^2 = 4(x-6)$ is (7, 6). Minimum distance is $\sqrt{2}$.

33. (0.5) : We have, $\lim_{x \rightarrow \infty} \sqrt{x + \sqrt{x + \sqrt{x}}} - \sqrt{x}$

$$= \lim_{x \rightarrow \infty} \frac{\sqrt{x + \sqrt{x}}}{\sqrt{x + \sqrt{x + \sqrt{x}}} + \sqrt{x}} = \lim_{x \rightarrow \infty} \frac{\sqrt{1 + \frac{1}{\sqrt{x}}}}{\sqrt{1 + \sqrt{\frac{1}{x} + \frac{1}{\sqrt{x^3}}}} + 1} = \frac{1}{2}$$

$$= 0.5$$

34. (2) : L.C.M. (p, q) = $2^2 \cdot 3^4 \cdot 5^2$

Let $p = 2^{a_1} \cdot 3^{b_1} \cdot 5^{c_1}$ and $q = 2^{a_2} \cdot 3^{b_2} \cdot 5^{c_2}$

$$\Rightarrow \max\{a_1, a_2\} = 2 \Rightarrow 5 \text{ ways}$$

$$\Rightarrow \max\{b_1, b_2\} = 4 \Rightarrow 9 \text{ ways}$$

$$\Rightarrow \max\{c_1, c_2\} = 2 \Rightarrow 5 \text{ ways}$$

$\therefore K = 3^2 \cdot 5^2$ can be expressed as $1 \cdot 3^2 \cdot 5^2, 3^2 \cdot 5^2$

35. (15) : $\frac{r}{R} = \cos A + \cos B + \cos C - 1$

$$\Rightarrow \frac{1}{8} = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2} - 1 + \cos C$$

$$\Rightarrow \frac{1}{8} = \sin \frac{C}{2} - 2 \sin^2 \frac{C}{2} \Rightarrow \sin \frac{C}{2} = \frac{1}{4}$$

$$\therefore \cos C = 1 - \frac{1}{8} = \frac{7}{8}. \text{ So, } \frac{1+\cos C}{1-\cos C} = \frac{1+7/8}{1-7/8} = 15$$

36. (5) : Image of orthocentre of ΔABC w.r.t. BC lies on the circle.

37. (1155) : Given, period of $f(x)$ is 7

$$\therefore \text{Period of } f\left(\frac{x}{3}\right) \text{ is } \frac{7}{1/3} = 21$$

and period of $g(x)$ is 11

$$\therefore \text{Period of } g\left(\frac{x}{5}\right) \text{ is } \frac{11}{1/5} = 55$$

$$\text{Now, } T_1 = \text{Period of } f(x) g\left(\frac{x}{5}\right) = 7 \times 55 = 385$$

$$\text{and } T_2 = \text{Period of } g(x) f\left(\frac{x}{3}\right) = 11 \times 21 = 231$$

\therefore Period of $F(x) = \text{L.C.M. of } \{T_1, T_2\}$

= L.C.M. of {385, 231}

$$= 7 \times 11 \times 3 \times 5 = 1155$$

38. (60) : Required number of ways $= \frac{4!}{2!2!} \times \frac{5!}{2!3!}$
 $= 60$

39. (18) : The equation of the line L be $y - 2 = m(x - 8)$,
 $m < 0$. Coordinates of P and Q are $P\left(8 - \frac{2}{m}, 0\right)$ and
 $Q(0, 2 - 8m)$.

$$\text{Now, } OP + OQ = 8 - \frac{2}{m} + 2 - 8m = 10 + \frac{2}{(-m)} + 8(-m)$$

$$\geq 10 + 2\sqrt{\frac{2}{(-m)} \times 8(-m)} \geq 18$$

So, absolute minimum value of $OP + OQ = 18$

40. (4) : We have $f(x) = \begin{cases} x-1, & 1 \leq x < 2 \\ 1-x, & 0 \leq x < 1 \end{cases}$

$f(x)$ is periodic with period 2 and also it is an even function.

Also, $\cos \pi x$ has period 2

$$\therefore I = \int_{-10}^{10} f(x) \cos \pi x \, dx = 2 \int_0^{10} f(x) \cos \pi x \, dx$$

($\because f(x) \cos \pi x$ is an even function)

$$= 2 \times 5 \int_0^2 f(x) \cos \pi x \, dx$$

$$= 10 \left[\int_0^1 (1-x) \cos \pi x \, dx + \int_1^2 (x-1) \cos \pi x \, dx \right]$$

$$= 10(I_1 + I_2)$$

$$I_2 = \int_1^2 (x-1) \cos \pi x \, dx$$

$$I_2 = - \int_0^1 t \cos \pi t \, dt \quad (\text{put } x-1=t)$$

$$I_1 = \int_0^1 (1-x) \cos \pi x \, dx = - \int_0^1 x \cos(\pi x) \, dx$$





$$\therefore I = 10 \left[-2 \int_0^1 x \cos \pi x \, dx \right] = -20 \left[x \frac{\sin \pi x}{\pi} + \frac{\cos \pi x}{\pi^2} \right]_0^1$$

$$= -20 \left[-\frac{1}{\pi^2} - \frac{1}{\pi^2} \right] = \frac{40}{\pi^2}$$

$$\therefore \frac{\pi^2}{10} I = 4$$



AVAILABLE BOUND VOLUMES



buy online at www.mtg.in

of your favourite magazines

Mathematics Today 2019
 (January - December) ₹ 380
 12 issues

Biology Today 2019
 (January - December) ₹ 380
 12 issues

Physics For You 2018
 (January - December) ₹ 380
 12 issues

Chemistry Today 2018
 (January - December) ₹ 380
 12 issues

Mathematics Today 2018
 (January - December) ₹ 380
 12 issues

Biology Today 2018
 (January - December) ₹ 380
 12 issues

How to order : Send money by demand draft/money order. Demand Draft should be drawn in favour of **MTG Learning Media (P) Ltd.** Mention the volume you require along with your name and address. OR buy online from www.mtg.in

Add ₹ 90 as postal charges
 Older issues can be accessed on digital.mtg.in in digital form.

Mail your order to :

Circulation Manager,
 MTG Learning Media (P) Ltd.
 Plot No. 99, Sector 44
 Institutional Area, Gurugram, (HR)
 Tel.: (0124) 6601200
 E-mail: info@mtg.in
 Web: www.mtg.in



CBSE warm-up!

CLASS-XI

TERM-I OBJECTIVE TYPE QUESTIONS*

Series 3

Complex Numbers and Quadratic Equations

MCQs

1. Express $(4 - 3i)^2$ in the standard form.
(a) $-44 - 177i$ (b) $44 - 177i$
(c) $44 + 177i$ (d) $-44 + 177i$
 2. $i^{57} + \frac{1}{i^{125}}$ is equal to
(a) $2i$ (b) $-2i$ (c) 0 (d) 2
 3. Find the multiplicative inverse of $2 - 3i$.
(a) $2/13 + i(3/13)$ (b) $2/13 - i(3/13)$
(c) $2 + 3i$ (d) None of these
 4. Solve: $x^2 + x + 1 = 0$
(a) $\pm i$ (b) $\pm(1/2)i$
(c) $\frac{-1 \pm \sqrt{3}i}{2}$ (d) $\frac{\pm\sqrt{3}i}{2}$
 5. What is the reciprocal of $3 + \sqrt{7}i$?
(a) $\frac{3}{16} - \frac{1}{16}i$ (b) $\frac{3}{16} + \frac{1}{16}i$
(c) $\frac{3}{16} + \frac{\sqrt{7}}{16}i$ (d) $\frac{3}{16} - \frac{\sqrt{7}}{16}i$
 6. The argument of the complex number $\left(\frac{i}{2} - \frac{2}{i}\right)$ is equal to
(a) $\frac{\pi}{4}$ (b) $\frac{3\pi}{4}$ (c) $\frac{\pi}{12}$ (d) $\frac{\pi}{2}$
 7. If $(x + iy)^{1/3} = a + ib$, where $x, y, a, b \in \mathbb{R}$, then
$$\frac{x - y}{a - b} =$$
- (a) $a^2 - b^2$ (b) $-2(a^2 + b^2)$
(c) $2(a^2 - b^2)$ (d) $a^2 + b^2$
 8. $i^2 + i^4 + i^6 + \dots$ ($2n + 1$) terms =
(a) 1 (b) -1 (c) i (d) $-i$
 9. Find the modulus of $\frac{1+i}{1-i}$.
(a) 1 (b) 2 (c) -1 (d) -2
 10. If $1 - i$ is a root of the equation $x^2 + ax + b = 0$, where $a, b \in \mathbb{R}$, then find the value of a .
(a) 2 (b) -2 (c) 3 (d) -3
 11. Simplify: $i^{n+100} + i^{n+50} + i^{n+48} + i^{n+46}$
(a) 0 (b) 1 (c) 2 (d) 3
 12. Express in standard form: $(1 + 2i)^6(4i + 3)^3$
(a) -14225 (b) -15625
(c) -18625 (d) -17525
 13. Find the sum of the complex number $-\sqrt{3} + \sqrt{-2}$ and $2\sqrt{3} - i$.
(a) $\sqrt{3} + i(\sqrt{2} - 1)$ (b) $\sqrt{3} - i(\sqrt{2} - 1)$
(c) $\sqrt{3} - i(\sqrt{2} + 1)$ (d) None of these
 14. Write the additive inverse of $-5 + 4i$.
(a) $5 + 4i$ (b) $5 - 4i$
(c) $5 + 3i$ (d) $5 - 3i$
 15. Find additive inverse of $\frac{3}{\sqrt{2} + i}$.
(a) $\sqrt{2} - i$ (b) $-\sqrt{2} + i$
(c) $\sqrt{2} + i$ (d) $-\sqrt{2} - i$

*Chapterwise practice questions for CBSE Exam Term- I as per the pattern issued by CBSE.

- 16.** Evaluate: $i^{141} + i^{142} + i^{143} + i^{144}$
- 0
 - 1
 - 2
 - 3
- 17.** If $\left(\frac{1+i}{1-i}\right)^n = 1$, then find the least positive integral value of n .
- 0
 - 1
 - 3
 - 4
- 18.** Evaluate : $(-\sqrt{-1})^{4n+3}$, $n \in N$
- 0
 - 1
 - i
 - $-i$
- 19.** Evaluate : $\left[i^{19} + \left(\frac{1}{i}\right)^{25}\right]^2$
- 2
 - 2
 - 4
 - 4
- 20.** What is the smallest positive integer n , for which $(1+i)^{2n} = (1-i)^{2n}$?
- 2
 - 4
 - 6
 - 8
- 21.** Find the multiplicative inverse of $\sqrt{5} + 3i$.
- $\frac{\sqrt{5}}{14} + \frac{3}{\sqrt{14}}i$
 - $-\frac{\sqrt{5}}{14} - \frac{3}{14}i$
 - $\frac{\sqrt{5}}{14} - \frac{3}{14}i$
 - None of these
- 22.** Multiply $3 - 2i$ by its conjugate.
- 13
 - 17
 - 21
 - 23
- 23.** Find the conjugate of $\frac{(3-2i)(2+3i)}{(1+2i)(2-i)}$.
- $\frac{63}{25} + \frac{16}{25}i$
 - $\frac{63}{25} - \frac{16}{25}i$
 - $-\frac{63}{25} - \frac{16}{25}i$
 - None of these
- 24.** If $z = 2 + \sqrt{3}i$, find the value of $z \cdot \bar{z}$.
- 5
 - 7
 - 9
 - 11
- 25.** For every complex number $z = x + iy \neq 0 + i0$; there exists a complex number $z_1 = \frac{x-iy}{x^2 + y^2}$, then value of zz_1 equals
- 0
 - 1
 - 2
 - 3
- 26.** Write the complex number $z = \frac{2+i}{(1+i)(1-2i)}$ in standard form.
- $\frac{1}{2} + \frac{i}{2}$
 - $\frac{1}{2} - \frac{i}{2}$
 - $-\frac{1}{2} + \frac{i}{2}$
 - $-\frac{1}{2} - \frac{i}{2}$
- 27.** Roots of the equation $x^2 - 4x + 13 = 0$ by factorisation methods are
- $7 \pm 11i$
 - $2 \pm 3i$
 - $3 \pm 2i$
 - $5 \pm 7i$
- 28.** Express $(5 - 3i)^3$ in the form of $a + ib$.
- $0 + 198i$
 - $-10 - 198i$
 - $10 - 198i$
 - $10 + 198i$
- 29.** The $\arg\left(\frac{3+i}{2-i} + \frac{3-i}{2+i}\right)$ is equal to
- $\frac{\pi}{2}$
 - 0
 - $\frac{\pi}{4}$
 - $-\frac{\pi}{4}$
- 30.** The real values of x and y for which the following equality hold, are respectively
 $(x^2 + 2xi) - (3x^2 + iy) = (3 - 5i) + (1 + 2iy)$
- 2, 3 or -2, $1/3$
 - 1, 3 or -1, $1/3$
 - 2, $1/3$ or -2, 3
 - 2, $1/3$ or -2, $-1/3$
- 31.** $(-i)(2i)\left(-\frac{1}{8}i\right)^3$ equals
- $0 + i$
 - $0 - i$
 - $\frac{1}{256} + 0i$
 - $0 + \frac{1}{256}i$
- 32.** If $|z^2 - 1| = |z|^2 + 1$, then z lies on
- imaginary axis
 - real axis
 - origin
 - None of these
- 33.** Solve : $x^2 + 3 = 0$
- $\pm\sqrt{3}i$
 - $\pm\sqrt{2}i$
 - $\pm\sqrt{5}i$
 - $\pm\sqrt{7}i$
- 34.** Solve : $x^2 - 14x + 58 = 0$
- $7 \pm 3i$
 - $5 \pm 2i$
 - $7 \pm 2i$
 - $5 \pm 3i$
- 35.** Solve : $5x^2 - 6x + 2 = 0$
- $\frac{3 \pm 2i}{5}$
 - $\frac{3 \pm i}{5}$
 - $\frac{3 \pm 5i}{2}$
 - $\frac{3 \pm i}{9}$
- 36.** Write the roots of the quadratic equation $x^2 + 8 = 0$.
- $\pm 2\sqrt{2}i$
 - $\pm 3\sqrt{3}i$
 - $\pm 3\sqrt{2}i$
 - $\pm 2\sqrt{3}i$
- 37.** Solve for x : $x^2 + 3x + 9 = 0$.
- $\frac{-3 \pm 2\sqrt{3}i}{2}$
 - $\frac{-3 \pm 2\sqrt{2}i}{5}$
 - $\frac{-3 \pm 3\sqrt{3}i}{2}$
 - None of these

38. If $4x + i(3x - y) = 3 + i(-6)$, where x and y are real numbers, then find the values of x and y .
(a) $1/4, 3/4$ (b) $3/4, 33/4$
(c) $33/4, -3/4$ (d) $-3/4, -33/4$

39. If $x = 2 + 5i$, then value of the expression $x^3 - 5x^2 + 33x - 49$ equals
(a) -20 (b) 10
(c) 20 (d) -29
40. If $(x + iy)(p + iq) = (x^2 + y^2)i$, then
(a) $p = x^2, q = y^2$ (b) $p = y, q = x$
(c) $p = x, q = y$ (d) $p = -x, q = y$

41. $\frac{5 + \sqrt{2}i}{1 - \sqrt{2}i}$ equals
(a) $1 - 2\sqrt{2}i$ (b) $1 - \sqrt{2}i$
(c) $1 + 2\sqrt{2}i$ (d) $1 + \sqrt{2}i$

42. Convert $\frac{1+7i}{(2-i)^2}$ into standard form.
(a) $0 + i$ (b) $1 - i$
(c) $-2 - 3i$ (d) $-1 + i$

43. If $z = 2 + i$, then $(z - 1)(\bar{z} - 5) + (\bar{z} - 1)(z - 5)$ is equal to
(a) 2 (b) 7 (c) -1 (d) -4

44. $\left| (1+i) \left(\frac{2+i}{3+i} \right) \right|$ is equal to
(a) $-\frac{1}{2}$ (b) 1 (c) -1 (d) $\frac{1}{2}$

45. If α and β are different complex numbers with $|\beta| = 1$, then $\left| \frac{\beta - \alpha}{1 - \bar{\alpha}\beta} \right|$ is
(a) 0 (b) $3/2$ (c) $1/2$ (d) 1

46. If $2 + i$ is a root of equation $x^3 - 5x^2 + 9x - 5 = 0$, then the other roots are
(a) 1 and $2 - i$ (b) -1 and $3 + i$
(c) 0 and 1 (d) -1 and $i - 2$

47. Find the value of P such that the difference of the roots of the equation $x^2 - Px + 8 = 0$ is 2 .
(a) ± 6 (b) ± 3 (c) ± 5 (d) ± 4

48. If $1 - i$ is a root of the equation $x^2 + ax + b = 0$, where $a, b \in \mathbb{R}$, then find the values of a and b .
(a) $2, 2$ (b) $-2, 2$ (c) $-2, -2$ (d) $1, 2$

49. Number of solutions of the equation $z^2 + |z|^2 = 0$ is
(a) 1 (b) 2
(c) 3 (d) infinitely many

50. The equation of smallest degree with real coefficients having $2 + 3i$ as one of the roots is
(a) $x^2 + 4x + 13 = 0$ (b) $x^2 + 4x - 13 = 0$
(c) $x^2 - 4x + 13 = 0$ (d) $x^2 - 4x - 13 = 0$

CASE BASED

Case-I : Read the following and answer any four questions from 51 to 55 given below.

On Sunday, Kanika was playing a ludo game with her mother. Kanika threw a die on the ludo cardboard. If the die stopped at a point $(-3, 4)$, then answer the following questions.



51. Let the point is represented as z in an argand plane, then find its modulus.
(a) $\sqrt{7}$ (b) 6 (c) $\sqrt{17}$ (d) 5

52. The multiplicative inverse of z is

- (a) $\frac{-3}{25} + 4i$ (b) $\frac{-3}{25} + \frac{4}{25}i$
(c) $\frac{-3}{25} - \frac{4}{25}i$ (d) $-3 + \frac{4}{25}i$

53. If her mother throw a die at $(1, \sqrt{3})$ which is represented by z' in a argand plane, then $z' \cdot z =$

- (a) $1 + i(4 - 3\sqrt{3})$
(b) $(-3 + 4\sqrt{3}) + i(4 - 3\sqrt{3})$
(c) $(-3 - 4\sqrt{3}) + i(4 - 3\sqrt{3})$
(d) $(-3 + 4\sqrt{3}) + i$

54. Find modulus of z' .

- (a) 2 (b) 1 (c) 3 (d) 4

55. The value of conjugate of $z + z'$ is

- (a) $-2 + (4 + \sqrt{3})i$ (b) $-2 + i$
(c) $-2 + 4i$ (d) $-2 - (4 + \sqrt{3})i$

Case-II : Read the following and answer any four questions from 56 to 60 given below.

Rohan and Rohit both are studying in Class XI. During summer vacations their father who is a mathematics teacher gave them a problem of quadratic equation to solve. Rohan found that the sum and product of roots were 3 and 9 respectively, whereas Rohit found that the sum and product of roots were -3 and -9 respectively.

Their father told them, Rohan found a correct product and Rohit found a correct sum.

- 56.** The correct quadratic equation is
 (a) $x^2 - 3x + 9 = 0$ (b) $x^2 + 3x - 9 = 0$
 (c) $x^2 + 3x + 9 = 0$ (d) $x^2 - 9x + 3 = 0$
- 57.** The discriminant of quadratic equation is
 (a) equal to zero (b) less than zero
 (c) greater than zero (d) can't compare
- 58.** The roots of quadratic equation are
 (a) $\frac{-3 \pm 3\sqrt{3}i}{2}$ (b) $\frac{1 \pm \sqrt{3}i}{2}$
 (c) $\frac{-3 \pm \sqrt{3}i}{2}$ (d) $\frac{3 \pm \sqrt{3}i}{2}$
- 59.** If $z_1 = \frac{-3+3\sqrt{3}i}{2}$ and $z_2 = \frac{-3-3\sqrt{3}i}{2}$, then $|z_1 z_2| =$
 (a) 9 (b) 5 (c) 4 (d) 6
- 60.** Complex roots of quadratic equation are always
 (a) negative of each other
 (b) same
 (c) occur in conjugate pair
 (d) can't say

ASSERTION & REASON

Directions (Q.No. 61-70) : In the following questions, a statement of assertion (Statement-I) is followed by a statement of reason (Statement-II). Mark the correct choice as :

- (a) If both Statement-I and Statement-II are true and Statement-II is the correct explanation of Statement-I.
 (b) If both Statement-I and Statement-II are true but Statement-II is not the correct explanation of Statement-I.
 (c) If Statement-I is true but Statement-II is false.
 (d) If Statement-I is false and Statement-II is true.

- 61.** **Statement-I:** Consider $|z_1| = 1$, $|z_2| = 2$ and $|z_3| = 3$. If $|z_1 + 2z_2 + 3z_3| = 6$, then the value of $|z_2 z_3 + 8z_3 z_1 + 27z_1 z_2|$ is 36.

Statement-II : $|z_1 + z_2 + z_3| \leq |z_1| + |z_2| + |z_3|$.

- 62.** **Statement-I :** Let $f(x)$ be a quadratic expression such that $f(0) + f(1) = 0$. If -2 is one of the roots of $f(x) = 0$, then other root is $3/5$.

Statement-II : If α and β are the zeroes of $f(x) = ax^2 + bx + c$, then sum of zeroes is $-b/a$ and product of zeroes is c/a .

- 63.** **Statement-I :** If P and Q are the points in the argand plane representing the complex numbers z_1

and z_2 respectively, then distance $|PQ| = |z_2 - z_1|$.

Statement-II : Locus of the point $P(z)$ satisfying $|z - (2 + 3i)| = 4$ is a straight line.

- 64.** **Statement-I :** If $z \neq 0$ is a complex number such that $\arg z = \pi/4$, then $\operatorname{Re}(z^2) = 0$.

Statement-II : If $z \neq 0$ and $\arg z = \pi/4$, then $\operatorname{Re} z = -\operatorname{Im} z$.

- 65.** **Statement-I :** The equation $3x^2 - 3x + 2 = 0$ has non-real roots.

Statement-II : If a, b, c are real and $b^2 - 4ac \geq 0$, then the roots of the equation $ax^2 + bx + c = 0$ are real and if $b^2 - 4ac < 0$, then roots of $ax^2 + bx + c = 0$ are non-real.

- 66.** **Statement-I :** Expression $\frac{2-\sqrt{-25}}{1-\sqrt{-16}}$ can be expressed in the form of $\frac{22}{17} + \frac{3}{17}i$.

Statement-II : Any number of the form $z = a + ib$, where a, b are real numbers and $i = \sqrt{-1}$ is called complex number.

- 67.** **Statement-I :** The conjugate of $\frac{(1+i)^2}{3-i}$ is $-\frac{1}{5} + \frac{3}{5}i$.

Statement-II : Conjugate of a complex number $z = a + ib$ is denoted by \bar{z} and defined as $\bar{z} = a - ib$.

- 68.** **Statement-I :** The argument of complex number $-2 + 2\sqrt{3}i$ is $\frac{2\pi}{3}$.

Statement-II : Modulus of a complex number $z = a + ib$ is denoted by $|z|$ and defined as $|z| = \sqrt{a^2 + b^2}$.

- 69.** **Statement-I :** If amplitude of $(z - 2 - 3i)$ is $\frac{\pi}{4}$, then locus of z represents a circle.

Statement-II : For any three complex numbers z_1, z_2 and z_3 , $z_1(z_2 + z_3) = z_1 z_2 + z_1 z_3$.

- 70.** **Statement-I :** The real and imaginary part of the complex number $\sqrt{37} + \sqrt{-19}$ are $\sqrt{37}$ and $-\sqrt{19}$ respectively.

Statement-II : Two complex numbers $z_1 = a_1 + ib_1$ and $z_2 = a_2 + ib_2$ are equal if $a_1 = a_2$ and $b_1 = b_2$.

SOLUTIONS

- (a) :** We have, $(4 - 3i)^3$
 $= 4^3 - (3i)^3 - 3 \times 4 \times 3i \times (4 - 3i)$
 $= 64 - 27i^3 - 36i(4 - 3i) = (64 - 108) + i(27 - 144)$
 $= (-44 - 117i)$
 $\therefore (4 - 3i)^3 = (-44 - 117i)$

2. (c) : Since $i^{4n} = 1$ and $i^{4n+1} = i$ ($n \in N$)

$$\therefore i^{57} + \frac{1}{i^{125}} = i + \frac{1}{i} = i - i = 0$$

$$\begin{aligned} \text{3. (a) : Let } z &= 2 - 3i, \text{ then } z^{-1} = \frac{1}{2-3i} = \frac{2+3i}{(2-3i)(2+3i)} \\ &= \frac{2+3i}{2^2 - (3i)^2} = \frac{2+3i}{13} = \frac{2}{13} + \frac{3}{13}i \end{aligned}$$

4. (c) : Here, $b^2 - 4ac = 1^2 - 4 \times 1 \times 1 = 1 - 4 = -3$

Therefore, the solutions are given by

$$x = \frac{-1 \pm \sqrt{-3}}{2 \times 1} = \frac{-1 \pm \sqrt{3}i}{2}$$

5. (d) : Reciprocal of $z = \frac{\bar{z}}{|z|^2}$

$$\text{Therefore, reciprocal of } 3 + \sqrt{7}i = \frac{3 - \sqrt{7}i}{16} = \frac{3}{16} - \frac{\sqrt{7}}{16}i$$

$$\text{6. (d) : Since } \left(\frac{i}{2} - \frac{2}{i}\right) = \frac{i}{2} - \frac{2i}{i^2} = \frac{i}{2} + 2i = \frac{5}{2}i$$

$$\text{So, argument of } \left(\frac{5}{2}i\right) = \tan^{-1}\left(\frac{5/2}{0}\right) = \frac{\pi}{2}.$$

7. (b) : $(x + iy)^{1/3} = a + ib \Rightarrow x + iy = (a + ib)^3$

$$\Rightarrow x + iy = a^3 - ib^3 + 3ia^2b - 3ab^2 = a^3 - 3ab^2 + i(3a^2b - b^3)$$

$$\therefore x = a^3 - 3ab^2 \text{ and } y = 3a^2b - b^3$$

$$\text{So, } \frac{x}{a} - \frac{y}{b} = a^2 - 3b^2 - 3a^2 + b^2$$

$$= -2a^2 - 2b^2 = -2(a^2 + b^2)$$

8. (b) : The first $2n$ terms will cancel in pairs and the last term is $(i^2)^{2n+1} = i^{4n} \cdot i^2 = -1$

9. (a) : Let $z = \frac{1+i}{1-i}$.

$$\text{Then, } |z| = \left| \frac{1+i}{1-i} \right| = \frac{|1+i|}{|1-i|} = \frac{\sqrt{1^2 + 1^2}}{\sqrt{1^2 + (-1)^2}} = \frac{\sqrt{2}}{\sqrt{2}} = 1$$

10. (b) : Given $1 - i$ is a root of the equation $x^2 + ax + b = 0$, where $a, b \in R$, therefore $1 + i$ will also be the root of the given equation.

Now, sum of roots $= (1 - i) + (1 + i) = -a/1 \Rightarrow a = -2$

$$\begin{aligned} \text{11. (a) : Given expression} &= i^{n+100} + i^{n+50} + i^{n+48} + i^{n+46} \\ &= i^n(i^{100} + i^{50} + i^{48} + i^{46}) \\ &= i^n[(i^2)^{50} + (i^2)^{25} + (i^2)^{24} + (i^2)^{23}] \\ &= i^n[(-1)^{50} + (-1)^{25} + (-1)^{24} + (-1)^{23}] \\ &= i^n[1 - 1 + 1 - 1] = i^n \cdot 0 = 0 \end{aligned}$$

12. (b) : We have, $(1 + 2i)^6(4i + 3)^3$

$$\begin{aligned} &= ((1 + 2i)^3)^2(4i + 3)^3 = (1 + 8i^3 + 6i + 12i^2)^2 \\ &= (1 - 12 + 6i - 8i)^2 (-64i + 108i + 27 - 144) \\ &= (-11 - 2i)^2(44i - 117) = 1936i^2 - 13689 = -15625 \end{aligned}$$

13. (a) : Let $z_1 = -\sqrt{3} + \sqrt{-2} = -\sqrt{3} + i\sqrt{2}$

and $z_2 = 2\sqrt{3} - i$

$$\text{Then } z_1 + z_2 = (-\sqrt{3} + i\sqrt{2}) + (2\sqrt{3} - i)$$

$$= (-\sqrt{3} + 2\sqrt{3}) + i(\sqrt{2} - 1) = \sqrt{3} + i(\sqrt{2} - 1)$$

14. (b) : Let $z = -5 + 4i$

The additive inverse of z is $-z$ i.e. $-(-5 + 4i) = 5 - 4i$.

$$\text{15. (b) : Let } z = \frac{3}{\sqrt{2}+i} = \frac{3(\sqrt{2}-i)}{2+1}$$

\therefore Additive inverse of z is $-z$ i.e. $-\sqrt{2} + i$.

$$\begin{aligned} \text{16. (a) : We have, } &i^{141} + i^{142} + i^{143} + i^{144} \\ &= i^{140}[i + i^2 + i^3 + i^4] \\ &= (i^4)^{35}[i - 1 - i + 1] \quad [\because i^2 = -1, i^3 = -i, i^4 = 1] \\ &= 0. \end{aligned}$$

17. (d) : We have, $\left(\frac{1+i}{1-i}\right)^n = 1$

$$\Rightarrow \left(\frac{1+i}{1-i} \times \frac{1+i}{1+i}\right)^n = 1$$

$$\Rightarrow \left(\frac{1-1+2i}{1+1}\right)^n = 1 \Rightarrow \left(\frac{2i}{2}\right)^n = 1 \Rightarrow i^n = 1$$

$\Rightarrow n$ is multiple of 4.

\therefore The least positive integral value of n is 4.

$$\begin{aligned} \text{18. (c) : } &(-\sqrt{-1})^{4n+3} = (-i)^{4n+3} \quad [\because \sqrt{-1} = i] \\ &= (-i)^{4n}(-i)^3 = 1(-1)(-i) \quad [\because i^4 = 1, i^3 = -i] \\ &= i \end{aligned}$$

19. (d) : Given expression

$$\begin{aligned} &\left[i^{19} + \left(\frac{1}{i}\right)^{25} \right]^2 = \left[i^{18} \cdot i + \left(\frac{1}{i}\right)^{25} \right]^2 \\ &= [(i^2)^9 \cdot i + (-i)^{25}]^2 = [(-1)^9 \cdot i - i^{25}]^2 = [-i - i^{24} \cdot i]^2 \\ &= [-i - (i^2)^{12} \cdot i]^2 = [-i - (-1)^{12} i]^2 \\ &= (-i - i)^2 = (-2i)^2 = 4i^2 = -4 \end{aligned}$$

20. (a) : We have, $(1 + i)^{2n} = (1 - i)^{2n}$

$$\Rightarrow \left(\frac{1+i}{1-i}\right)^{2n} = 1 \Rightarrow \left(\frac{1+i}{1-i} \times \frac{1+i}{1+i}\right)^{2n} = 1$$

$$\Rightarrow \left(\frac{1-1+2i}{1+1}\right)^{2n} = 1 \Rightarrow \left(\frac{2i}{2}\right)^{2n} = 1$$

$\Rightarrow (i)^{2n} = 1 \Rightarrow 2n$ is a multiple of 4.

\therefore The least positive integral value of n is 2.

21. (c) : Let $z = \sqrt{5} + 3i$

$$\bar{z} = \sqrt{5} - 3i \text{ and } |z|^2 = (\sqrt{5})^2 + (3)^2 = 14$$

$$z^{-1} = \frac{\bar{z}}{|z|^2} = \frac{\sqrt{5} - 3i}{14} = \frac{\sqrt{5}}{14} - \frac{3}{14}i$$

22. (a) : Let $z = 3 - 2i$

$$\bar{z} = 3 + 2i$$

$$\therefore z \cdot \bar{z} = (3 - 2i)(3 + 2i) = 9 - 4i^2 = 9 + 4 = 13$$

23. (a) : Let $z = \frac{(3-2i)(2+3i)}{(1+2i)(2-i)} = \frac{6+9i-4i+2}{2-i+4i+2}$

$$\begin{aligned} &= \frac{12+5i}{4+3i} = \left(\frac{12+5i}{4+3i} \right) \times \left(\frac{4-3i}{4-3i} \right) \\ &= \frac{48-36i+20i+15}{4^2-(3i)^2} = \frac{63-16i}{16+9} = \frac{63}{25} - \frac{16}{25}i \end{aligned}$$

$$\therefore \text{Conjugate of } z = \frac{63}{25} + \frac{16}{25}i.$$

24. (b) : $z = 2 + \sqrt{3}i$

$$\therefore \bar{z} = 2 - \sqrt{3}i$$

$$z \cdot \bar{z} = (2 + \sqrt{3}i)(2 - \sqrt{3}i) = 4 - 3i^2 = 4 + 3 = 7$$

25. (b) : $zz_1 = (x+iy) \left(\frac{x-iy}{x^2+y^2} \right)$

$$= (x+iy) \left(\frac{x}{x^2+y^2} - i \frac{y}{x^2+y^2} \right)$$

$$= \left(\frac{x^2}{x^2+y^2} + \frac{y^2}{x^2+y^2} \right) + i \left(\frac{-xy}{x^2+y^2} + \frac{xy}{x^2+y^2} \right)$$

$$= 1 + i \cdot (0) = 1$$

26. (a) : We have,

$$z = \frac{2+i}{(1+i)(1-2i)} = \frac{2+i}{3-i} = \frac{(2+i)(3+i)}{(3-i)(3+i)}$$

$$= \frac{5+5i}{10} = \frac{1}{2} + i \frac{1}{2}$$

27. (b) : We have, $x^2 - 4x + 13 = 0$

$$\therefore x = \frac{4 \pm \sqrt{16 - 4(13)}}{2} = \frac{4 \pm \sqrt{16 - 52}}{2}$$

$$= \frac{4 \pm \sqrt{-36}}{2} = \frac{4 \pm 6i}{2} = 2 \pm 3i$$

28. (b) : $(5 - 3i)^3 = 125 - 225i - 135 + 27i = -10 - 198i$

29. (b) : $\arg \left(\frac{3+i}{2-i} + \frac{3-i}{2+i} \right) = \arg \left(\frac{6+5i+i^2+6-5i+i^2}{5} \right)$

$$= \arg \left(\frac{10}{5} \right) = \arg(2) = 0$$

30. (a) : The given equality can be rewritten as

$$(x^4 - 3x^2) + i(2x - y) = 4 + i(2y - 5)$$

$$\Rightarrow x^4 - 3x^2 = 4, 2x - y = 2y - 5$$

$$\Rightarrow (x^2 - 4)(x^2 + 1) = 0 \Rightarrow x = \pm 2 \quad (\because x^2 \neq -1)$$

\therefore At $x = 2, y = 3$ and at $x = -2, y = 1/3$

$$31. (d) : (-i)(2i) \left(-\frac{1}{8}i \right)^3 = 2 \times \frac{1}{8 \times 8 \times 8} \times i^5$$

$$= \frac{1}{256}(i^2)^2 i = \frac{1}{256}i = 0 + \frac{1}{256}i$$

32. (a) : Let $z = x + iy$. Then, $|z^2 - 1| = |z|^2 + 1$

$$\Rightarrow |x^2 - y^2 - 1 + i2xy| = |x + iy|^2 + 1$$

$$\Rightarrow (x^2 - y^2 - 1)^2 + 4x^2y^2 = (x^2 + y^2 + 1)^2$$

$$\Rightarrow 4x^2 = 0 \Rightarrow x = 0$$

Hence z lies on y -axis or imaginary axis.

33. (a) : Given, $x^2 + 3 = 0 \Rightarrow x^2 = -3$

$$\Rightarrow x = \pm \sqrt{-3} = \pm \sqrt{3}i$$

$$34. (a) : x = \frac{-(-14) \pm \sqrt{(-14)^2 - 4(1)(58)}}{2}$$

$$= \frac{14 \pm \sqrt{196 - 232}}{2} = \frac{14 \pm 6i}{2} = 7 \pm 3i$$

35. (b) : Given, $5x^2 - 6x + 2 = 0$

$$D = (-6)^2 - 4(5)(2) = 36 - 40 = -4$$

$$x = \frac{-(-6) \pm \sqrt{-4}}{2(5)} = \frac{6 \pm 2i}{10} = \frac{3 \pm i}{5}$$

36. (a) : Given, $x^2 + 8 = 0 \Rightarrow x^2 = -8$

$$\Rightarrow x = \pm \sqrt{-8} = \pm 2\sqrt{2}i$$

Your favourite MTG Books/Magazines available in BIHAR at

Ganthalaya - Bhagalpur Ph: 0641-2301413, 7654885581, 9304288600

Kishore Pustak Bhandar - Bhagalpur Ph: 0641-2409338, 9431274924

Kiran Book Depot - Gaya Ph: 0631-2222213, 2430863, 9430056741

Gyan Ganga Ltd - Patna Ph: 0612-2578394, 9304826650, 9771492969

Manoj New Agency - Patna Ph: 0612-2206151, 2362169, 9905271351

Nova Publisher & Distributors - Patna Ph: 0612-3259655, 2666404, 9334112031, 7488409037

Saraswati Pustak Bhandar - Patna Ph: 8757594019, 9334606425, 9304976881

Sash Pustak Bhandar - Patna Ph: 9095024722, 9386961203

Competitor Book Centre - Purnea Ph: 7654559152

Chaurasia Book Centre - Purnea Ph: 064-54240214, 9005717044, 9573644800

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

37. (c) : Given, $x^2 + 3x + 9 = 0$

$$x = \frac{-3 \pm \sqrt{9 - 4(9)}}{2} = \frac{-3 \pm \sqrt{-27}}{2} = \frac{-3 \pm 3\sqrt{3}i}{2}$$

38. (b) : We have, $4x + i(3x - y) = 3 + i(-6)$

Equating the real and imaginary parts, we get
 $4x = 3$, $3x - y = -6$

On solving above questions, we get

$$x = \frac{3}{4} \text{ and } y = \frac{33}{4}.$$

39. (a) : Given $x = 2 + 5i$

$$\Rightarrow x - 2 = 5i$$

$$\Rightarrow x^2 - 4x + 29 = 0$$

Now, $x^3 = 5x^2 + 33x - 49$

$$= x(x^2 - 4x + 29) - 1(x^2 - 4x + 29) - 20 = -20$$

40. (b) : Since, $x^2 + y^2 = (x + iy)(x - iy)$

$$\therefore (p + iq) = \left(\frac{x^2 + y^2}{x + iy} \right)i = (x - iy)i = y + ix$$

$$\therefore p = y, q = x$$

41. (c) : We have, $\frac{5 + \sqrt{2}i}{1 - \sqrt{2}i} = \frac{5 + \sqrt{2}i}{1 - \sqrt{2}i} \times \frac{1 + \sqrt{2}i}{1 + \sqrt{2}i}$

$$= \frac{5 + 5\sqrt{2}i + \sqrt{2}i - 2}{1 - (\sqrt{2})^2} = \frac{3(1 + 2\sqrt{2}i)}{3} = 1 + 2\sqrt{2}i$$

42. (d) : We have, $z = \frac{1+7i}{(2-i)^2} = \frac{1+7i}{(4+i^2-4i)}$

$$= \frac{1+7i}{3-4i} \times \frac{3+4i}{3+4i}$$

$$= \frac{3+4i+21i+28i^2}{9-16i^2} = \frac{3-28+25i}{25} = -1+i$$

43. (d) : $(z-1)(\bar{z}-5) + (\bar{z}-1)(z-5)$

$$= 2\operatorname{Re}[(z-1)(\bar{z}-5)] \quad (\because z_1\bar{z}_2 + z_2\bar{z}_1 = 2\operatorname{Re}(z_1\bar{z}_2))$$

$$= 2\operatorname{Re}[(1+i)(-3-i)] = 2(-2) = -4 \quad (\text{Given } z = 2 + i)$$

$$\text{44. (b) : } \frac{|1+i||2+i|}{|3+i|} = \frac{\sqrt{2}\sqrt{5}}{\sqrt{10}} = 1$$

45. (d) : $|\beta| = 1 \Rightarrow |\beta|^2 = 1 \Rightarrow \beta\bar{\beta} = 1$

$$\therefore \left| \frac{\beta - \alpha}{1 - \bar{\alpha}\beta} \right| = \left| \frac{\beta - \alpha}{\beta\bar{\beta} - \bar{\alpha}\beta} \right| = \left| \frac{\beta - \alpha}{\beta(\bar{\beta} - \bar{\alpha})} \right| \\ = \frac{|\beta - \alpha|}{|\beta||\bar{\beta} - \bar{\alpha}|} = \frac{|\beta - \alpha|}{1|\beta - \alpha|} = \frac{|\beta - \alpha|}{|\beta - \alpha|} = 1$$

46. (a) : Since complex roots always occur in conjugate pair.

So, $2 - i$ is also a root of given equation.

Given equation is $x^3 - 5x^2 + 9x - 5 = 0$.

$x = 1$ satisfies this equation.

\therefore Other roots are $(2 - i)$ and 1 .

47. (a) : Let α, β be the roots of the equation $x^2 - Px + 8 = 0$.

Then, $|\alpha - \beta| = 2$, $\alpha + \beta = P$ and $\alpha\beta = 8$.

Now, $(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$

$$\therefore (2)^2 = P^2 - 32 \Rightarrow P^2 - 32 = 4 \Rightarrow P = \pm 6$$

48. (b) : Since complex roots always occur in conjugate pair.

\therefore Other root is $1 + i$.

$$\text{Sum of roots} = \frac{-a}{1} = (1 - i) + (1 + i) \Rightarrow a = -2$$

$$\text{Product of roots} = \frac{b}{1} = (1 - i)(1 + i) \Rightarrow b = 2$$

49. (d) : We have, $z^2 + |z|^2 = 0$

$$\Rightarrow x^2 - y^2 + i2xy + x^2 + y^2 = 0 \quad [\text{Taking } z = x + iy]$$

$$\Rightarrow 2x^2 + i2xy = 0$$

$$\Rightarrow 2x^2 = 0 \text{ and } 2xy = 0$$

[Equating real and imaginary part]

$$\Rightarrow x = 0 \text{ and } xy = 0$$

Thus, $x = 0$ and y can have any real value.

Hence infinitely many solutions.

50. (c) : Since complex roots always exist in conjugate pair, so other root will be $2 - 3i$.

Now, $S = \text{sum of roots} = 2 + 3i + 2 - 3i = 4$

and $P = \text{product of roots} = (2 + 3i)(2 - 3i) = 4 + 9 = 13$

So, required equation will be

$$x^2 - Sx + P = 0 \Rightarrow x^2 - 4x + 13 = 0$$

51. (d) : Clearly, $z = -3 + 4i$

$$\therefore |z| = \sqrt{(-3)^2 + 4^2} = \sqrt{9+16} = \sqrt{25} = 5$$

52. (c) : The multiplicative inverse of $-3 + 4i$ is

$$\frac{1}{-3+4i} = \frac{-1}{3-4i} \times \frac{3+4i}{3+4i} = \frac{-(3+4i)}{25} = \frac{-3}{25} - \frac{4}{25}i$$

53. (c) : $\because z' = 1 + \sqrt{3}i$

$$\therefore z' \cdot z = (1 + \sqrt{3}i)(-3 + 4i) = 1(-3 + 4i) + \sqrt{3}i(-3 + 4i)$$

$$= -3 + 4i - 3\sqrt{3}i + 4\sqrt{3}i^2$$

$$= (-3 - 4\sqrt{3}) + i(4 - 3\sqrt{3})$$

$$\text{54. (a) : } |z'| = \sqrt{1+3} = \sqrt{4} = 2$$

$$\text{55. (d) : } z + z' = -3 + 4i + 1 + \sqrt{3}i = -2 + (4 + \sqrt{3})i$$

Thus, its conjugate is $-2 - (4 + \sqrt{3})i$

56. (c) : The required quadratic equation is given by

$$x^2 - (\text{sum of roots})x + \text{Product of roots} = 0$$

$$\Rightarrow x^2 - 3x + 9 = 0$$

57. (b) : Here, $a = 1, b = 3, c = 9$

$$\text{Discriminant} = b^2 - 4ac$$

$$= 3^2 - 4 \cdot 1 \cdot 9 = 9 - 36 = -27 < 0$$

58. (a) : $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
 $\Rightarrow x = \frac{-3 \pm \sqrt{-27}}{2} = \frac{-3 \pm 3\sqrt{3}i}{2}$

59. (a) : $|z_1 z_2| = |\text{Product of roots}| = |9| = 9$

60. (c) : Complex roots of quadratic equation are of the form $a \pm ib$.
 Thus, occur in conjugate pair.

61. (b) : $|z_2 z_3 + 8z_3 z_1 + 27z_1 z_2| = \left| z_1 z_2 z_3 \left(\frac{1}{z_1} + \frac{8}{z_2} + \frac{27}{z_3} \right) \right|$
 $= |z_1||z_2||z_3| \left| \frac{1}{z_1} + \frac{8}{z_2} + \frac{27}{z_3} \right| = |z_1||z_2||z_3| \left| \frac{\bar{z}_1}{|\bar{z}_1|^2} + \frac{8\bar{z}_2}{|\bar{z}_2|^2} + \frac{27\bar{z}_3}{|\bar{z}_3|^2} \right|$
 $= |z_1||z_2||z_3| \left| \frac{\bar{z}_1}{1} + \frac{8\bar{z}_2}{4} + \frac{27\bar{z}_3}{9} \right| = |z_1||z_2||z_3| \left| \bar{z}_1 + 2z_2 + 3z_3 \right|$
 $= |z_1||z_2||z_3||z_1 + 2z_2 + 3z_3| = 1 \times 2 \times 3 \times 6 = 36$

62. (b) : Since, $x = -2$ is a root of $f(x) = 0$.
 $\therefore f(x) = (x+2)(ax+b)$

But $f(0) + f(1) = 0 \Rightarrow 2b + 3a + 3b = 0 \Rightarrow -\frac{b}{a} = \frac{3}{5}$


63. (c) : Statement-I is a standard result.

We have, $|z - (2+3i)| = 4$

\Rightarrow Distance of $P(z)$ from the point $(2, 3)$ is equal to 4.
 \Rightarrow Locus of P is a circle with centre at $(2, 3)$ and radius 4.

64. (c) : Let $z = x + iy$.

Since $\arg z = \frac{\pi}{4}$, therefore $\angle XOP = \frac{\pi}{4}$



$\Rightarrow \tan(\angle XOP) = 1$
 $\Rightarrow MP = OM \Rightarrow y = x \neq 0 (\because z \neq 0)$
 Hence $z^2 = (x+yi)^2 = (x+xi)^2$
 $= x^2 + i^2 x^2 + 2x^2i$
 $= x^2 - x^2 + 2x^2i = 2x^2i$
 $\Rightarrow \operatorname{Re}(z^2) = 0$

We have seen that if $\arg z = \frac{\pi}{4}$ and $z \neq 0$
 then $\operatorname{Re} z = \operatorname{Im} z$.

65. (a) : We have, $3x^2 - 3x + 2 = 0$
 Here, $b^2 - 4ac = 9 - 4(3)(2) = 9 - 24 = -15 < 0$
 \therefore Roots are non-real.

66. (a) : We have, $\frac{2 - \sqrt{-25}}{1 - \sqrt{-16}} = \frac{2 - 5i}{1 - 4i} = \frac{2 - 5i}{1 - 4i} \times \frac{1 + 4i}{1 + 4i}$
 $= \frac{(2+20)i + (8-5)}{1 - 16i^2} = \frac{22 + 3i}{17} = \frac{22}{17} + \frac{3}{17}i$

Also, Statement-II is a standard result.

67. (d) : Clearly statement-II is true.

Let $z = \frac{(1+i)^2}{3-i} = \frac{1+2i+i^2}{3-i} = \frac{2i}{3-i} \times \frac{3+i}{3+i}$
 $= \frac{6i+2i^2}{9-i^2} = \frac{6i-2}{10} = -\frac{1}{5} + \frac{3}{5}i \therefore \bar{z} = -\frac{1}{5} - \frac{3}{5}i$

68. (b) : $\tan \alpha = \left| \frac{2\sqrt{3}}{-2} \right| = \sqrt{3} = \tan \frac{\pi}{3} \Rightarrow \alpha = \frac{\pi}{3}$

Clearly, $\operatorname{Re}(z) < 0$ and $\operatorname{Im}(z) > 0$. So, the point representing z lies in the second quadrant.

$\therefore \arg(z) = \pi - \alpha = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$.

69. (d) : Let $z = x + iy$. Then, $z - 2 - 3i = (x + iy) - 2 - 3i = (x - 2) + i(y - 3)$

Now, $\tan \frac{\pi}{4} = \frac{y-3}{x-2} \Rightarrow 1 = \frac{y-3}{x-2}$

$\Rightarrow x - y + 1 = 0$, which is a straight line.

70. (d) : Let $z = \sqrt{37} + \sqrt{-19} = \sqrt{37} + \sqrt{19 \times (-1)}$
 $= \sqrt{37} + \sqrt{19} \sqrt{-1} = \sqrt{37} + i\sqrt{19}$

$\therefore \operatorname{Re}(z) = \sqrt{37}$ and $\operatorname{Im}(z) = \sqrt{19}$



Your favourite MTG Books/Magazines available in UTTARAKHAND at

National Book House - Dehradun Ph: 0135-2659430; 9897830283

Army Book Depot - Dehradun Ph: 0135-2756683; 9897927092

Om Vidyा Educational Book - Dehradun Ph: 9897833882

Career Zone - Haldwani Ph: 05946-262051; 9412128075, 9412383167

World Vision Publication - Haldwani Ph: 9927932200, 9027240169

Diamond Stationers - Haridwar Ph: 0133-4252043; 9358398035, 9359763348


Consul Book Deptt - Nainital Ph: 0592-235164; 9412084105

Indra Book Emporium - Roorkee Ph: 0133-276105; 8126293314

Cambridge Book Deptt - Roorkee Ph: 272341, 272345; 971910955

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call 0124-6601200 for further assistance.

MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 80

Series 3 : Trigonometric Function

Time Taken : 60 Min.

Only One Option Correct Type

- In a ΔABC if $(\sqrt{3}-1)a = 2b$, $A = 3B$, then C is
 - (a) 60°
 - (b) 120°
 - (c) 30°
 - (d) 45°
- If $\sin 6\theta + \sin 4\theta + \sin 2\theta = 0$, then the general value of θ is
 - (a) $\frac{n\pi}{4}, n\pi \pm \frac{\pi}{3}$
 - (b) $\frac{n\pi}{4}, n\pi \pm \frac{\pi}{6}$
 - (c) $\frac{n\pi}{4}, 2n\pi \pm \frac{\pi}{3}$
 - (d) $\frac{n\pi}{4}, 2n\pi \pm \frac{\pi}{6}$
- The value of $\cot 16^\circ \times \cot 44^\circ + \cot 44^\circ \times \cot 76^\circ - \cot 76^\circ \times \cot 16^\circ$ is
 - (a) 4
 - (b) 1
 - (c) 3
 - (d) 0
- If $\tan^2\alpha \tan^2\beta + \tan^2\beta \tan^2\gamma + \tan^2\gamma \tan^2\alpha + 2\tan^2\alpha \tan^2\beta \tan^2\gamma = 1$, then the value of $\sin^2\alpha + \sin^2\beta + \sin^2\gamma$ is
 - (a) 0
 - (b) -1
 - (c) 1
 - (d) None of these
- The solution of the equation $\log_{\cos x} \sin x + \log_{\sin x} \cos x = 2$ is given by
 - (a) $x = 2n\pi + \frac{\pi}{4}, n \in I$
 - (b) $x = n\pi + \frac{\pi}{2}, n \in I$
 - (c) $x = n\pi + \frac{\pi}{8}, n \in I$
 - (d) None of these
- In any triangle, with usual notations the minimum value of $r_1 r_2 r_3 / r^3$ is equal to
 - (a) 1
 - (b) 9
 - (c) 27
 - (d) None of these

One or More Than One Option(s) Correct Type

- If $A + B = \frac{\pi}{3}$ and $\cos A + \cos B = 1$, then which of the following is/are true?
 - (a) $\cos(A - B) = \frac{1}{3}$
 - (b) $|\cos A - \cos B| = \sqrt{\frac{2}{3}}$
 - (c) $\cos(A - B) = -\frac{1}{3}$
 - (d) $|\cos A - \cos B| = \frac{1}{2\sqrt{3}}$
- If $\sin(\alpha + \beta) = 1$ and $\sin(\alpha - \beta) = 1/2$, where $\alpha, \beta \in [0, \pi/2]$, then
 - (a) $\tan(\alpha + 2\beta) = -\sqrt{3}$
 - (b) $\tan(2\alpha + \beta) = -1/\sqrt{3}$
 - (c) $\tan(\alpha + 2\beta) = \sqrt{3}$
 - (d) $\tan(2\alpha + \beta) = 1/\sqrt{3}$
- If sides of triangle ABC are a, b and c such that $2b = a + c$, then
 - (a) $\frac{b}{c} > \frac{2}{3}$
 - (b) $\frac{b}{c} > \frac{1}{3}$
 - (c) $\frac{b}{c} < 2$
 - (d) $\frac{b}{c} < \frac{1}{2}$



BUY ONLINE

Now you can buy


MTG Books & Magazines

Log on to : www.mtg.in

10 Very Similar Practice Tests **JEE Main**

Visit
www.mtg.in
for latest offers
and to buy
online!

₹ 200/-



HIGHLIGHTS

- 10 Very Similar Practice Tests as per the latest pattern of NTA JEE Main (60 MCQs+ 30 Numerical Value Type Questions)
- OMR sheet provided at the end of each test
- Detailed solutions of each practice test included

Available at all leading book shops throughout India. To buy online visit www.mtg.in.
For more information or for help in placing your order,
call 0124-6601200, 1800-10-38673 (Toll Free) or e-mail info@mtg.in

Also available on
Flipkart, Snapdeal, Amazon,
Infibeam on cash on delivery

10. If $\sin x + \cos x + \tan x + \cot x + \sec x + \cosec x = 7$ and $\sin 2x = a - b\sqrt{7}$, then
 (a) $a = 8$ (b) $b = 22$
 (c) $a = 22$ (d) $b = 8$

11. If $\cos(\theta - \alpha), \cos\theta, \cos(\theta + \alpha)$ are in H.P., then $\cos\theta \sec \frac{\alpha}{2}$ is equal to
 (a) -1 (b) $-\sqrt{2}$
 (c) $\sqrt{2}$ (d) 2

12. If A is the area and $2s$ is the sum of the sides of a triangle, then
 (a) $A \leq \frac{s^2}{4}$ (b) $A \leq \frac{s^2}{3\sqrt{3}}$
 (c) $A < \frac{s^2}{\sqrt{3}}$ (d) None of these

13. In a ΔABC ,
 (a) $\sin A \sin B \sin C \leq \frac{3\sqrt{3}}{8}$
 (b) $\sin^2 A + \sin^2 B + \sin^2 C \leq \frac{9}{4}$
 (c) $\sin A \sin B \sin C$ is always positive
 (d) $\sin^2 A + \sin^2 B \leq 1 + \cos C$

Comprehension Type

Paragraph for Q. No. 14 and 15

The sides of a triangle ABC are $7, 8, 6$ the smallest angle being ' C '.

14. The length of the median from vertex C is
 (a) $\frac{\sqrt{95}}{4}$ (b) $\frac{\sqrt{95}}{2}$
 (c) $\sqrt{\frac{95}{2}}$ (d) $\frac{\sqrt{95}}{3}$
15. The length of the internal bisector of angle C is
 (a) $\sqrt{30}$ (b) $\frac{14}{5}\sqrt{6}$
 (c) $\frac{14}{5}$ (d) $2\sqrt{6}$

Matrix Match Type

16. Match the following:

	Column-I	Column-II
P.	Max $\{5 \sin \theta + 3 \sin(0 - \alpha), \theta \in R\} = 7$, then the set of possible values of α is	1. $2n\pi + 3\pi/4, n \in Z$
Q.	If $x \neq \frac{(2n+1)\pi}{2}$ and $(\cos x)^{(\sin^2 x - 3 \sin x + 2)} = 1$, then $x =$	2. $2n\pi \pm \frac{\pi}{3}, n \in Z$
R.	If $\sqrt{(\sin x)^2 + 2^{1/4}} \cos x = 0$, then $x =$	3. $2n\pi + \cos^{-1}(1/3), n \in Z$
S.	If $\log_5 \tan x = (\log_5 4)$ ($\log_4(3 \sin x)$), then $x =$	4. $2n\pi, n \in Z$

P	Q	R	S
(a) 1	3	4	2
(b) 2	4	1	3
(c) 4	2	3	1
(d) 4	1	2	3

Numerical Answer Type

17. If $\frac{1}{\cos 290^\circ} + \frac{1}{\sqrt{3} \sin 250^\circ} = \lambda$, then the value of $9\lambda^4 + 81\lambda^2 + 97$ must be _____.
18. Consider a ΔABC , in which the sides are $a = (n+1)$, $b = (n+2)$, $c = n$ with $\tan C = 4/3$, then the value of $\Delta/12$ is _____.
19. If $3 \sin x + 4 \cos x = 5$, then the value of $90 \tan^2(x/2) - 60 \tan(x/2) + 16$ is equal to _____.
20. If $k \left[\tan^6 \frac{\pi}{9} - 33 \tan^4 \frac{\pi}{9} + 27 \tan^2 \frac{\pi}{9} \right] = 9$, then the value of k is _____.



Keys are published in this issue. Search now! ☺

SELF CHECK

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.



CBSE

warm-up!

CLASS-XII

TERM-I OBJECTIVE TYPE QUESTIONS*

Series 2

Inverse Trigonometric Functions

MCQs

- Find the principal value of $\sec^{-1}(2)$.
 - $\frac{\pi}{2}$
 - $\frac{\pi}{4}$
 - $\frac{\pi}{3}$
 - 0
- $\sin^{-1}\left(\frac{-1}{2}\right) =$
 - $\frac{\pi}{3}$
 - $-\frac{\pi}{3}$
 - $\frac{\pi}{6}$
 - $-\frac{\pi}{6}$
- Evaluate : $\cos\left\{-\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \frac{\pi}{6}\right\}$
 - 0
 - 1
 - $\frac{1}{2}$
 - $\frac{1}{\sqrt{2}}$
- $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right) =$
 - $\frac{\pi}{2}$
 - $\frac{2\pi}{3}$
 - $\frac{\pi}{4}$
 - $\frac{5\pi}{6}$
- If $\theta = \tan^{-1}a$, $\phi = \tan^{-1}b$ and $ab = -1$, then $|\theta - \phi|$ is equal to
 - 0
 - $\frac{\pi}{4}$
 - $\frac{\pi}{2}$
 - None of these
- Domain of $\cos^{-1}[x]$ (where $[.]$ denotes G.I.F.) is
 - $[-1, 2]$
 - $[-1, 2)$
 - $(-1, 2]$
 - None of these
- Evaluate : $\operatorname{cosec}\left\{\cot^{-1}\left(\frac{4}{3}\right)\right\}$
 - $\frac{3}{4}$
 - $\frac{5}{3}$
 - $\frac{3}{5}$
 - $\frac{4}{3}$
- $\cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right) + 4\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$ is equal to
 - $\frac{\pi}{6}$
 - $\frac{\pi}{3}$
 - $\frac{4\pi}{3}$
 - $\frac{3\pi}{4}$
- Evaluate :

$$\cot^{-1}(1) + \operatorname{cosec}^{-1}(\sqrt{2}) + \sec^{-1}(2)$$
 - $\frac{\pi}{6}$
 - $\frac{5\pi}{6}$
 - $\frac{\pi}{3}$
 - $\frac{2\pi}{3}$
- Find the value of $\cot\left[\sin^{-1}\left\{\cos(\tan^{-1}1)\right\}\right]$.
 - 0
 - $\pi/4$
 - 1
 - does not exist.
- Range of $f(x) = \cos^{-1}x + \sec^{-1}x$ is
 - $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$
 - $\left[\frac{\pi}{4}, \frac{3\pi}{4}\right]$
 - $\left\{\frac{\pi}{4}, \frac{3\pi}{4}\right\}$
 - None of these
- Evaluate : $\tan^{-1}\left(\tan\frac{3\pi}{4}\right)$
 - $\frac{\pi}{2}$
 - $\frac{\pi}{4}$
 - $-\frac{\pi}{4}$
 - $-\frac{\pi}{2}$
- $\cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right)$ is equal to
 - $\frac{\pi}{4}$
 - $\frac{\pi}{6}$
 - $\frac{\pi}{3}$
 - $\frac{2\pi}{3}$
- Find the domain of $\sec^{-1}(2x+1)$.
 - $(-\infty, -1]$
 - $[0, \infty)$
 - $R - (-1, 1)$
 - None of these

*Chapterwise practice questions for CBSE Exam Term- I as per the pattern issued by CBSE.

15. $\tan^{-1}(-\sqrt{3}) + \sec^{-1}(-2) - \operatorname{cosec}^{-1}\left(\frac{2}{\sqrt{3}}\right)$ is equal to
 (a) $\frac{5\pi}{6}$ (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{3}$ (d) 0
16. If $x, y, z \in [-1, 1]$ such that $\cos^{-1}x + \cos^{-1}y + \cos^{-1}z = 0$, then find the value of $x + y + z$.
 (a) 0 (b) 1 (c) 2 (d) 3
17. $\cos^{-1}\left(\frac{-1}{2}\right) + 2\sin^{-1}\left(\frac{-1}{2}\right)$ is equal to
 (a) $\frac{\pi}{3}$ (b) $\frac{2\pi}{3}$ (c) $\frac{3\pi}{4}$ (d) $\frac{5\pi}{8}$
18. If $\cos^{-1}\alpha + \cos^{-1}\beta + \cos^{-1}\gamma = 3\pi$, then $\alpha(\beta + \gamma) + \beta(\gamma + \alpha) + \gamma(\alpha + \beta)$ equals
 (a) 0 (b) 1 (c) 6 (d) 12
19. $\sec^{-1}\left(\frac{-2}{\sqrt{3}}\right)$ equals to
 (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{3}$ (c) $\frac{5\pi}{6}$ (d) $\frac{2\pi}{3}$
20. If $\cot^{-1}\left(-\frac{1}{5}\right) = x$, then the value of $\sin x$ is
 (a) $\frac{1}{\sqrt{26}}$ (b) $\frac{-1}{\sqrt{26}}$ (c) $\frac{5}{\sqrt{26}}$ (d) $\frac{-5}{\sqrt{26}}$
21. The principal value of $\cos^{-1}\left\{\sin\left(\cos^{-1}\frac{1}{2}\right)\right\}$ is
 (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{2}$
 (c) $\frac{\pi}{6}$ (d) none of these
22. The domain of $\operatorname{cosec}^{-1}(3 - 2x)$ is
 (a) $(-\infty, -1]$ (b) $[2, \infty)$
 (c) $R - (-1, 1)$ (d) None of these
23. Evaluate : $\sin^{-1}(\sin(-600^\circ))$.
 (a) 0 (b) $\frac{\pi}{3}$ (c) $\frac{2\pi}{3}$ (d) $-\frac{\pi}{3}$
24. If $\cos^{-1}x + \cos^{-1}y + \cos^{-1}z = \pi$, then $x^2 + y^2 + z^2 + 2xyz =$
 (a) 0 (b) 1
 (c) -1 (d) None of these
25. Evaluate : $\sin\left(2\cot^{-1}\left(-\frac{5}{12}\right)\right)$.
 (a) $\frac{1}{169}$ (b) $-\frac{1}{169}$ (c) $\frac{120}{169}$ (d) $-\frac{120}{169}$
26. $\sin\left(2\tan^{-1}\frac{1}{3}\right) + \cos(\tan^{-1}2\sqrt{2}) =$
 (a) $\frac{13}{15}$ (b) $\frac{15}{13}$ (c) $\frac{14}{15}$ (d) $\frac{15}{14}$
27. If $\tan^{-1}\frac{4}{3} = \theta$, find the value of $\cos \theta$.
 (a) $\frac{3}{5}$ (b) $\frac{4}{5}$ (c) $\frac{5}{3}$ (d) $\frac{5}{4}$
28. The principal value of $\sin^{-1}\left(\sin\frac{2\pi}{3}\right)$ is
 (a) $\frac{\pi}{2}$ (b) $-\frac{\pi}{2}$ (c) $\frac{\pi}{3}$ (d) $-\frac{\pi}{3}$
29. Evaluate :
 $\cot^{-1}(-\sqrt{3}) + \tan^{-1}(1) + \sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$
 (a) $\frac{\pi}{4}$ (b) $\frac{5\pi}{4}$ (c) $\frac{\pi}{6}$ (d) $\frac{5\pi}{6}$
30. The value of $\tan^{-1}\left(\tan\frac{5\pi}{6}\right)$ is
 (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{3}$ (c) $-\frac{\pi}{6}$ (d) $\frac{\pi}{3}$
31. The value of $\cos^{-1}\left(\cos\frac{13\pi}{6}\right)$ is
 (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{6}$ (d) $\frac{2\pi}{3}$
32. Find the principal value of $\tan^{-1}\left(\tan\frac{7\pi}{6}\right)$.
 (a) $\frac{\pi}{3}$ (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{6}$ (d) $\frac{5\pi}{6}$
33. The value of $\cos^2\left(\frac{1}{2}\cos^{-1}\left(\frac{3}{5}\right)\right)$ is
 (a) $\frac{4}{5}$ (b) $\frac{5}{4}$ (c) $\frac{3}{4}$ (d) $\frac{4}{3}$
34. The value of $\sin\left(\cot^{-1}\frac{3}{4}\right)$ is
 (a) $\frac{4}{5}$ (b) $\frac{5}{4}$ (c) $\frac{3}{4}$ (d) $\frac{4}{3}$
35. The value of $\tan\left(\frac{\pi}{4} + \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right)$
 $+ \tan\left(\frac{\pi}{4} - \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right)$ is
 (a) $\frac{a}{b}$ (b) $\frac{2b}{a}$ (c) $\frac{b}{a}$ (d) $\frac{b}{2a}$
36. The number of triplets (x, y, z) satisfies the equation
 $\sin^{-1}x + \sin^{-1}y + \sin^{-1}z = \frac{3\pi}{2}$ is
 (a) 1 (b) 2 (c) 0 (d) infinite

37. If $6\sin^{-1}(x^2 - 6x + 8.5) = \pi$, then x is equal to
 (a) 1 (b) 2 (c) 3 (d) 8

38. If $\sin^{-1}(x^2 - 7x + 12) = 0$, then $x =$
 (a) -2 (b) 4 (c) -3 (d) 5

39. If $\tan^{-1}(x^2 - 4x + 4) = 0$, then x equals to
 (a) 2 (b) 4 (c) 3 (d) 5

40. If $\tan^{-1}(\cot\theta) = 20$, then θ is equal to

- (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{4}$
 (c) $\frac{\pi}{6}$ (d) None of these

41. Which of the following is the principal value branch of $\sec^{-1}x$?

- (a) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (b) $(0, \pi)$
 (c) $[0, \pi]$ (d) $[0, \pi] - \left\{\frac{\pi}{2}\right\}$

42. Evaluate : $\cos\left(2\cos^{-1}\left(\frac{2}{5}\right)\right)$

- (a) $\frac{13}{25}$ (b) $\frac{17}{25}$ (c) $-\frac{13}{25}$ (d) $-\frac{17}{25}$

43. Evaluate : $\sin\left[\frac{\pi}{3} + \sin^{-1}\left(\frac{1}{2}\right)\right]$.
 (a) $\sqrt{3}/2$ (b) 1/2 (c) 0 (d) 1

44. $\cot^{-1}(-\sqrt{3})$ is equal to
 (a) $\frac{5\pi}{6}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{4}$

45. The value of $\tan^{-1}(1) + \tan^{-1}(0) + \tan^{-1}(-1)$ is equal to

- (a) π (b) $\frac{5\pi}{4}$
 (c) $\frac{\pi}{2}$ (d) None of these

46. The solution set of the equation

- $\tan^{-1}x - \cot^{-1}x = \cos^{-1}(2 - x)$ is
 (a) $[0, 1]$ (b) $[-1, 1]$
 (c) $[1, 3]$ (d) None of these

47. $\cos^{-1}[\cos(2\cot^{-1}(\sqrt{3}))] =$
 (a) $\frac{2\pi}{3}$ (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{3}$

48. If $2(\sin^{-1}x)^2 - 5(\sin^{-1}x) + 2 = 0$, then $x =$
 (a) $\pi/6$ (b) $\pi/3$ (c) 2 (d) 1/2

49. The principal solution of $\cos^{-1}\left(\cos\left(\frac{9\pi}{4}\right)\right)$ is

- (a) $\frac{7\pi}{4}$ (b) $-\frac{\pi}{4}$ (c) $\frac{9\pi}{4}$ (d) $\frac{\pi}{4}$


50. The principal solution of $\sin^{-1}\left(\sin\left(\frac{5\pi}{3}\right)\right)$ is

- (a) $\frac{4\pi}{3}$ (b) $\frac{5\pi}{3}$ (c) $-\frac{5\pi}{3}$ (d) $-\frac{\pi}{3}$

CASE BASED

Case I : Read the following and answer any four questions from 51 to 55 given below.

Two men on either side of a temple, which is 30 metres high above the stairs, observe its top at the angles of elevation α and β respectively (as shown in the figure below). The distance between the two men is $40\sqrt{3}$ metres and the distance between the first person A and the temple is $30\sqrt{3}$ metres.



51. $\angle CAB = \alpha =$

- (a) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$ (b) $\sin^{-1}\left(\frac{1}{2}\right)$
 (c) $\sin^{-1}(2)$ (d) $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$

52. $\angle CAB = \alpha =$

- (a) $\cos^{-1}\left(\frac{1}{5}\right)$ (b) $\cos^{-1}\left(\frac{2}{5}\right)$
 (c) $\cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$ (d) $\cos^{-1}\left(\frac{4}{5}\right)$

53. $\angle BCA = \beta =$

- (a) $\tan^{-1}\left(\frac{1}{2}\right)$ (b) $\tan^{-1}(2)$
 (c) $\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (d) $\tan^{-1}(\sqrt{3})$


54. $\angle ABC =$

- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$

55. Domain and range of $\cos^{-1} x$ are respectively
 (a) $(-1, 1), (0, \pi)$ (b) $[-1, 1], (0, \pi)$
 (c) $[-1, 1], [0, \pi]$ (d) $(-1, 1), \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

Case II : Read the following and answer any four questions from 56 to 60 given below.

The Government of India is planning to fix a hoarding board at the face of a building on the road of a busy market for awareness of Covid-19 protocol. Ram, Robert and Rahim are the three engineers who are working on this project. "A" is considered as a person, viewing the hoarding board, 20 metres away from the building and standing at the edge of a pathway nearby. Ram, Robert and Rahim suggested to the firm to place the hoarding board at three different locations namely, C, D and E. "C" is at the height of 10 metres from the ground level. For the viewer A, the angle of elevation of "D" is double the angle of elevation of "C". The angle of elevation of "E" is triple the angle of elevation of "C" for the same viewer.



56. Measure of $\angle CAB =$
 (a) $\tan^{-1}(2)$ (b) $\tan^{-1}\left(\frac{1}{2}\right)$
 (c) $\tan^{-1}(1)$ (d) $\tan^{-1}(3)$
57. Measure of $\angle DAB =$
 (a) $\tan^{-1}\left(\frac{3}{4}\right)$ (b) $\tan^{-1}(3)$
 (c) $2\tan^{-1}\left(\frac{1}{2}\right)$ (d) $\tan^{-1}(4)$
58. Measure of $\angle EAB =$
 (a) $\tan^{-1}(11)$ (b) $\tan^{-1}3$
 (c) $\tan^{-1}\left(\frac{2}{11}\right)$ (d) $3\tan^{-1}\left(\frac{1}{2}\right)$
59. If A' is another viewer, which is at a distance of 5 m from A as shown in figure, then the difference between $\angle CAB$ and $\angle CA'B$ is

- (a) $\tan^{-1}(1/2)$ (b) $\tan^{-1}(1/2) - \tan^{-1}(2/5)$
 (c) $\tan^{-1}\left(\frac{2}{5}\right)$ (d) $\tan^{-1}\left(\frac{11}{21}\right)$
60. Domain and range of $\tan^{-1} x$ are respectively
 (a) $R^+, \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (b) $R^-, \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
 (c) $R, \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (d) $R, \left(0, \frac{\pi}{2}\right)$

ASSERTION & REASON

Directions (Q.No. 61-70) : In the following questions, a statement of assertion (Statement-I) is followed by a statement of reason (Statement-II). Mark the correct choice as :

- (a) If both Statement-I and Statement-II are true and Statement-II is the correct explanation of Statement-I.
 (b) If both Statement-I and Statement-II are true but Statement-II is not the correct explanation of Statement-I.
 (c) If Statement-I is true but Statement-II is false.
 (d) If Statement-I is false but Statement-II is true.

61. Statement-I : Principal value of

$$\sin^{-1}\left(\sin\left(\frac{2\pi}{3}\right)\right) \text{ is } \frac{\pi}{3}.$$

Statement-II : Principal value branch of arc sin function is $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

62. Statement-I : Range of $f(x) = \sin^{-1}x + \tan^{-1}x + \sec^{-1}x$ is $\left[\frac{\pi}{4}, \frac{3\pi}{4}\right]$.
 Statement-II : $f(x) = \sin^{-1}x + \tan^{-1}x + \sec^{-1}x$ is defined for all $x \in [-1, 1]$.

63. Statement-I : Number of roots of the equation $\cot^{-1}x + \cos^{-1}2x + \pi = 0$ is zero.

Statement-II : Range of $\cot^{-1}x$ and $\cos^{-1}x$ is $(0, \pi)$ and $[0, \pi]$, respectively.

64. Statement-I : Range of $f(x) = \cot^{-1}(2x - x^2)$ is $(0, \pi)$.
 Statement-II : $\cot^{-1}x$ is defined for all $x \in R$.

65. Statement-I : The domain for

$$f(x) = \sin^{-1}(1+x^2)$$

is $\{0, 1\}$.
 Statement-II : $\sin^{-1}x$ is defined only if $x \in [-1, 1]$.

66. Statement-I : The value of $\sin(2\tan^{-1}(0.75))$, is 0.96.
 Statement-II : Range of $\sin^{-1}(x)$ is $[0, \pi]$.




A must have series to score high in CBSE Board Term-I

**100
PERCENT**

**Exam
Ready**

CBSE BOARD TERM-I
Objective Question Bank



Price
₹ 125/-
each

HIGHLIGHTS OF THE SERIES

CHAPTERWISE

RECAP NOTES • MCQs • CASE / EXTRACT / SOURCE BASED MCQs • ASSERTION & REASONING MCQs

Available at all leading book shops throughout India. To buy online visit www.mtg.in
For more information or for help in placing your order, call 0124-6601200 or email info@mtg.in

67. **Statement-I :** The value of $\cos\left(\frac{\pi}{3} - \cos^{-1}\frac{1}{2}\right)$ is 0.

Statement-II : If $y = \cos x$, then $x = \cos^{-1}y$, where $x \in [0, \pi]$.

68. **Statement-I :** The principal solution of

$$4\tan^{-1}\left(\frac{1}{\sqrt{3}}\right) \text{ is } \frac{2\pi}{3}.$$

Statement-II : $\tan^{-1}x$ is defined for all $x \in R$.

69. **Statement-I :** The domain of function defined by $f(x) = \sin^{-1}\sqrt{x-1}$ is $[1, 2]$.

Statement-II : $\sin^{-1}x$ is defined for all $x \in R$.

70. **Statement-I :** The domain of $\sin^{-1}x + \cos^{-1}x + \tan^{-1}x$ is R .

Statement-II : $\text{Dom } (f+g)(x) = \text{Dom } f(x) \cap \text{Dom } g(x)$

$$\Rightarrow \theta - \phi = \frac{\pi}{2} \text{ or } \phi - \theta = \frac{\pi}{2}$$

$$\Rightarrow |\theta - \phi| = \frac{\pi}{2}$$

6. **(b) :** Clearly, $-1 \leq [x] \leq 1$

$$\Rightarrow -1 \leq x < 2 \Rightarrow x \in [-1, 2)$$

$$7. \quad \text{(b) : Let } \theta = \cot^{-1}\left(\frac{4}{3}\right) \Rightarrow \cot \theta = \frac{4}{3}$$

$$\text{Now, cosec}\left[\cot^{-1}\left(\frac{4}{3}\right)\right] = \text{cosec } \theta = \sqrt{1+\cot^2 \theta}$$

$$= \sqrt{1+\frac{16}{9}} = \frac{5}{3}$$

$$8. \quad \text{(c) : } \cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right) + 4\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

$$= \frac{\pi}{3} + 2 \cdot \frac{\pi}{6} + 4 \cdot \frac{\pi}{6} = \frac{\pi}{3} + \frac{\pi}{3} + \frac{2\pi}{3} = \frac{4\pi}{3}$$

9. **(b) :** $\cot^{-1}(1) + \text{cosec}^{-1}(\sqrt{2}) + \sec^{-1}(2)$

$$= \frac{\pi}{4} + \frac{\pi}{4} + \frac{\pi}{3} = \frac{5\pi}{6}$$

10. **(c) :** We have, $\cot[\sin^{-1}\{\cos(\tan^{-1}1)\}]$

$$= \cot\left[\sin^{-1}\left(\cos\frac{\pi}{4}\right)\right] = \cot\left(\sin^{-1}\frac{1}{\sqrt{2}}\right) = \cot\frac{\pi}{4} = 1$$

11. **(d) :** Given $f(x) = \cos^{-1}x + \sec^{-1}x$

Domain of $\cos^{-1}x = [-1, 1]$

Domain of $\sec^{-1}x = (-\infty, \infty) - (-1, 1)$

Domain of $f(x) = [-1, 1] \cap [(-\infty, \infty) - (-1, 1)] = [-1, 1]$

Now, $f(-1) = \cos^{-1}(-1) + \sec^{-1}(-1) = \pi + \pi = 2\pi$

and $f(1) = \cos^{-1}(1) + \sec^{-1}(1) = 0 + 0 = 0$

\therefore Range of $f(x)$ is $[0, 2\pi]$.

$$12. \quad \text{(c) : } \tan^{-1}\left(\tan\frac{3\pi}{4}\right) = \tan^{-1}\left\{\tan\left(\pi - \frac{\pi}{4}\right)\right\}$$

$$= \tan^{-1}\left(-\tan\frac{\pi}{4}\right) = \tan^{-1}\left\{\tan\left(-\frac{\pi}{4}\right)\right\} = -\frac{\pi}{4}$$

$$13. \quad \text{(d) : The value of } \cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3} + 2 \times \frac{\pi}{6}$$

$$= \frac{\pi}{3} + \frac{\pi}{3} = \frac{2\pi}{3}$$

14. **(d) :** $\sec^{-1}(2x+1)$ is defined if

$$2x+1 \geq 1 \text{ or } 2x+1 \leq -1$$

$$\Rightarrow 2x \geq 0 \text{ or } 2x \leq -2 \Rightarrow x \geq 0 \text{ or } x \leq -1$$

$$\Rightarrow x \in (-\infty, -1] \cup [0, \infty)$$

Hence, the domain of $\sec^{-1}(2x+1)$ is $(-\infty, -1] \cup [0, \infty)$

SOLUTIONS

1. **(c) :** Let $\sec^{-1}(2) = 0 \Rightarrow \sec \theta = 2 = \sec \frac{\pi}{3}$

$$\Rightarrow \theta = \frac{\pi}{3} \in \left[0, \pi\right] - \left\{\frac{\pi}{2}\right\}$$

\therefore Principal value of $\sec^{-1}(2)$ is $\frac{\pi}{3}$.

2. **(d) :** Let $\sin^{-1}\left(\frac{-1}{2}\right) = 0 \Rightarrow \sin \theta = -\frac{1}{2} = -\sin \frac{\pi}{6} = \sin\left(-\frac{\pi}{6}\right)$

$$\Rightarrow \theta = -\frac{\pi}{6} \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

\therefore Principal value of $\sin^{-1}\left(\frac{-1}{2}\right)$ is $\left(-\frac{\pi}{6}\right)$.

3. **(b) :** $\cos\left[-\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \frac{\pi}{6}\right] = \cos\left[-\frac{\pi}{6} + \frac{\pi}{6}\right]$

$$= \cos 0 = 1$$

4. **(d) :** Let $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right) = 0$

$$\Rightarrow \cos \theta = -\frac{\sqrt{3}}{2} = -\cos \frac{\pi}{6} = \cos\left(\pi - \frac{\pi}{6}\right) = \cos \frac{5\pi}{6}$$

$$\Rightarrow \theta = \frac{5\pi}{6} \in [0, \pi] \quad \therefore \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right) = \frac{5\pi}{6}$$

5. **(c) :** Given, $\theta = \tan^{-1}a$ and $\phi = \tan^{-1}b$ and $ab = -1$

$$\therefore \tan \theta \tan \phi = ab = -1$$

$$\Rightarrow \tan \theta = -\cot \phi \text{ or } \tan \phi = -\cot \theta$$

$$\Rightarrow \tan \theta = \tan\left(\frac{\pi}{2} + \phi\right) \text{ or } \tan \phi = \tan\left(\frac{\pi}{2} + \theta\right)$$



15. (d) : The value of

$$\tan^{-1}(-\sqrt{3}) + \sec^{-1}(-2) - \cosec^{-1}\left(\frac{2}{\sqrt{3}}\right) = \frac{-\pi}{3} + \frac{2\pi}{3} - \frac{\pi}{3} = 0$$

16. (d) : We have, $x, y, z \in [-1, 1]$

$$\Rightarrow -1 \leq x \leq 1, -1 \leq y \leq 1, -1 \leq z \leq 1$$

$$\Rightarrow 0 \leq \cos^{-1}x \leq \pi, 0 \leq \cos^{-1}y \leq \pi, 0 \leq \cos^{-1}z \leq \pi$$

$$\therefore \cos^{-1}x + \cos^{-1}y + \cos^{-1}z = 0 \quad [\text{Given}]$$

$$\Rightarrow \cos^{-1}x = 0, \cos^{-1}y = 0 \text{ and } \cos^{-1}z = 0$$

$$\Rightarrow x = y = z = 1. \text{ Hence, } x + y + z = 3.$$

17. (a) : Principal value of $\cos^{-1}\left(\frac{-1}{2}\right)$ is $\frac{2\pi}{3}$

and principal value of $\sin^{-1}\left(\frac{-1}{2}\right)$ is $\left(-\frac{\pi}{6}\right)$.

$$\therefore \cos^{-1}\left(\frac{-1}{2}\right) + 2\sin^{-1}\left(\frac{-1}{2}\right)$$

$$= \frac{2\pi}{3} + \left(2 \times \frac{-\pi}{6}\right) = \frac{2\pi}{3} - \frac{\pi}{3} = \frac{\pi}{3}$$

$$18. (c) : \cos^{-1}\alpha + \cos^{-1}\beta + \cos^{-1}\gamma = 3\pi$$

$$\therefore 0 \leq \cos^{-1}x \leq \pi$$

$$\Rightarrow \cos^{-1}\alpha = \cos^{-1}\beta = \cos^{-1}\gamma = \pi \Rightarrow \alpha = \beta = \gamma = -1$$

$$\therefore \alpha(\beta + \gamma) + \beta(\gamma + \alpha) + \gamma(\alpha + \beta)$$

$$= -1(-1 - 1) + (-1)(-1 - 1) + (-1)(-1 - 1)$$

$$= 2 + 2 + 2 = 6$$

19. (c) : Let $\sec^{-1}\left(\frac{-2}{\sqrt{3}}\right) = \theta$

$$\Rightarrow \sec\theta = \frac{-2}{\sqrt{3}} = -\sec\left(\frac{\pi}{6}\right) = \sec\left(\pi - \frac{\pi}{6}\right) = \sec\frac{5\pi}{6}$$

$$\Rightarrow \theta = \frac{5\pi}{6} \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$$

$$\therefore \text{Principal value of } \sec^{-1}\left(\frac{-2}{\sqrt{3}}\right) \text{ is } \frac{5\pi}{6}.$$

20. (c) : Given, $\cot^{-1}\left(-\frac{1}{5}\right) = x$

$$\Rightarrow \cot x = -\frac{1}{5}, 0 < x < \pi$$

$$\therefore \cosec x = \sqrt{1 + \cot^2 x} = \sqrt{1 + \left(-\frac{1}{5}\right)^2} = \frac{\sqrt{26}}{5}$$

$$\Rightarrow \sin x = \frac{5}{\sqrt{26}} \quad [\because \cosec x \text{ is positive}]$$

21. (c) : We know that $\cos^{-1}\frac{1}{2} = \frac{\pi}{3}$.

$$\therefore \cos^{-1}\left\{\sin\left(\cos^{-1}\frac{1}{2}\right)\right\} = \cos^{-1}\left(\sin\frac{\pi}{3}\right)$$

$$= \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6} \quad \left[\because \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}\right]$$

22. (d) : For the domain of $\cosec^{-1}(3 - 2x)$, $|3 - 2x| \geq 1$

$$\Rightarrow 3 - 2x \leq -1 \text{ or } 3 - 2x \geq 1$$

$$\Rightarrow -2x \leq -4 \text{ or } -2x \geq -2 \Rightarrow x \geq 2 \text{ or } x \leq 1.$$

\therefore Domain of $\cosec^{-1}(3 - 2x)$ is $(-\infty, 1] \cup [2, \infty)$.

$$23. (b) : \sin^{-1}(\sin(-600^\circ)) = \sin^{-1}\left\{\sin\left(-600 \times \frac{\pi}{180}\right)\right\}$$

$$= \sin^{-1}\left\{\sin\left(-\frac{10\pi}{3}\right)\right\} = \sin^{-1}\left(-\sin\frac{10\pi}{3}\right)$$

$$= \sin^{-1}\left\{-\sin\left(3\pi + \frac{\pi}{3}\right)\right\} = \sin^{-1}\left\{-\left(-\sin\frac{\pi}{3}\right)\right\}$$

$$= \sin^{-1}\left(\sin\frac{\pi}{3}\right) = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}$$

24. (b) : Let $\cos^{-1}x = \alpha, \cos^{-1}y = \beta$ and $\cos^{-1}z = \gamma$

$$\Rightarrow \cos\alpha = x, \cos\beta = y \text{ and } \cos\gamma = z$$

$$\text{Given, } \cos^{-1}x + \cos^{-1}y + \cos^{-1}z = \pi$$

$$\Rightarrow \alpha + \beta + \gamma = \pi \quad \therefore \alpha + \beta = \pi - \gamma$$

$$\therefore \cos(\alpha + \beta) = \cos(\pi - \gamma)$$

$$\Rightarrow \cos\alpha\cos\beta - \sin\alpha\sin\beta = -\cos\gamma$$

$$\Rightarrow xy - \sqrt{1-x^2}\sqrt{1-y^2} = -z$$

$$\Rightarrow xy + z = \sqrt{1-x^2}\sqrt{1-y^2}$$

Squaring both sides, we get

$$x^2y^2 + z^2 + 2xyz = 1 - x^2 - y^2 + x^2y^2$$

$$\Rightarrow x^2 + y^2 + z^2 + 2xyz = 1.$$

25. (d) : Let $\cot^{-1}\left(-\frac{5}{12}\right) = x, 0 < x < \pi$

$$\Rightarrow \cot x = -\frac{5}{12}, 0 < x < \pi.$$

For these values of x , $\cosec x > 0$.

MONTHLY TEST DRIVE CLASS XII | ANSWER KEY

- | | | | | |
|-------------|------------|-------------|------------|-------------|
| 1. (b) | 2. (c) | 3. (d) | 4. (a) | 5. (d) |
| 6. (d) | 7. (a,b,c) | 8. (b,c) | 9. (b,c,d) | 10. (b,c,d) |
| 11. (a,c,d) | 12. (c) | 13. (a,c,d) | 14. (a) | 15. (b) |
| 16. (a) | 17. (144) | 18. (0) | 19. (2) | 20. (4) |

$$\therefore \cosec x = \sqrt{1 + \cot^2 x} = \sqrt{1 + \left(-\frac{5}{12}\right)^2} = \frac{13}{12}$$

$$\Rightarrow \sin x = \frac{12}{13}.$$

Now, $\cos x = \frac{\cos x}{\sin x} \cdot \sin x = \cot x \cdot \sin x$

$$= \left(-\frac{5}{12}\right) \frac{12}{13} = -\frac{5}{13}.$$

$$\therefore \sin\left(2 \cot^{-1}\left(-\frac{5}{12}\right)\right) = \sin 2x = 2 \sin x \cos x = -\frac{120}{169}.$$

26. (c) : Let $\tan^{-1} \frac{1}{3} = \alpha \Rightarrow \tan \alpha = \frac{1}{3}$.

$$\sin\left(2 \tan^{-1} \frac{1}{3}\right) = \sin 2\alpha = \frac{2 \tan \alpha}{1 + \tan^2 \alpha} = \frac{2 \times \frac{1}{3}}{1 + \left(\frac{1}{3}\right)^2} = \frac{3}{5}$$

Let $\tan^{-1} 2\sqrt{2} = \beta \Rightarrow \tan \beta = 2\sqrt{2}$.

$$\sec \beta = \sqrt{1 + \tan^2 \beta} = \sqrt{1 + (2\sqrt{2})^2} = 3 \Rightarrow \cos \beta = \frac{1}{3}.$$

$$\Rightarrow \cos(\tan^{-1} 2\sqrt{2}) = \cos \beta = \frac{1}{3}.$$

$$\therefore \sin\left(2 \tan^{-1} \frac{1}{3}\right) + \cos(\tan^{-1} 2\sqrt{2}) = \frac{3}{5} + \frac{1}{3} = \frac{14}{15}.$$

27. (a) : Given, $\tan^{-1} \frac{4}{3} = \theta$, where $\theta \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$

$$\therefore \tan \theta = \frac{4}{3}.$$

We know that $\cos \theta > 0$, when $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

$$\therefore \cos \theta = \frac{1}{\sec \theta} = \frac{1}{\sqrt{1 + \tan^2 \theta}} = \frac{1}{\sqrt{1 + \frac{16}{9}}} = \frac{3}{5}$$

28. (c) : Since, $\sin \frac{2\pi}{3} = \sin\left(\pi - \frac{\pi}{3}\right) = \sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$

$$\therefore \sin^{-1}\left(\sin \frac{2\pi}{3}\right) = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{3}.$$

Hence, the principal value of $\sin^{-1}\left(\sin \frac{2\pi}{3}\right)$ is $\frac{\pi}{3}$.

29. (b) : $\cot^{-1}(-\sqrt{3}) + \tan^{-1}(1) + \sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$

$$= \left(\pi - \frac{\pi}{6}\right) + \frac{\pi}{4} + \frac{\pi}{6} = \frac{5\pi}{6} + \frac{\pi}{4} + \frac{\pi}{6} = \frac{5\pi}{4}$$

30. (c) : $\tan^{-1}\left(\tan\left(\pi - \frac{\pi}{6}\right)\right)$

$$= \tan^{-1}\left(-\tan\left(\frac{\pi}{6}\right)\right) = \tan^{-1}\left(\frac{-1}{\sqrt{3}}\right) = -\frac{\pi}{6}$$

31. (c) : $\cos^{-1} \cos\left(2\pi + \frac{\pi}{6}\right) = \cos^{-1}\left(\cos\left(\frac{\pi}{6}\right)\right)$

$$= \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}$$

32. (c) : $\tan^{-1}\left(\tan \frac{7\pi}{6}\right) = \tan^{-1}\left\{\tan\left(\pi + \frac{\pi}{6}\right)\right\}$

$$= \tan^{-1}\left\{\tan\left(\frac{\pi}{6}\right)\right\} = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$$

33. (a) : Let $\cos^{-1} \frac{3}{5} = \theta \Rightarrow \cos \theta = \frac{3}{5}$

$$\therefore \cos^2\left\{\left(\frac{1}{2}\right) \cos^{-1}\left(\frac{3}{5}\right)\right\} = \cos^2\left(\frac{\theta}{2}\right)$$

$$= \frac{\cos \theta + 1}{2} = \frac{\frac{3}{5} + 1}{2} = \frac{4}{5}$$

34. (a) : Let $\cot^{-1} \frac{3}{4} = \theta \Rightarrow \cot \theta = \frac{3}{4}$

Now, $\cosec \theta = \sqrt{1 + \cot^2 \theta} = \sqrt{1 + \frac{9}{16}} = \frac{5}{4}$

$$\therefore \sin \theta = \frac{4}{5} \quad \therefore \sin\left(\cot^{-1}\left(\frac{3}{4}\right)\right) = \sin \theta = \frac{4}{5}$$

35. (b) : Let $\cos^{-1} \frac{a}{b} = \theta \Rightarrow \cos \theta = \frac{a}{b}$

So, we have, $\tan\left(\frac{\pi}{4} + \frac{1}{2}\theta\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\theta\right)$

$$= \frac{1 + \tan \theta / 2}{1 - \tan \theta / 2} + \frac{1 - \tan \theta / 2}{1 + \tan \theta / 2}$$

$$= \frac{(1 + \tan \theta / 2)^2 + (1 - \tan \theta / 2)^2}{1 - \tan^2 \theta / 2} = \frac{2(1 + \tan^2 \theta / 2)}{1 - \tan^2 \theta / 2}$$

$$= \frac{2}{\cos \theta} = \frac{2}{a/b} = \frac{2b}{a}$$

36. (a) : We have, $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$

$$\therefore -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}, -\frac{\pi}{2} \leq \sin^{-1} y \leq \frac{\pi}{2}$$



and $\frac{-\pi}{2} \leq \sin^{-1} z \leq \frac{\pi}{2}$

\therefore The above condition will satisfy if

$$\sin^{-1}x = \sin^{-1}y = \sin^{-1}z = \frac{\pi}{2} \Rightarrow x = y = z = 1$$

37. (b) : We have, $6 \sin^{-1}(x^2 - 6x + 8.5) = \pi$

$$\Rightarrow \sin^{-1}(x^2 - 6x + 8.5) = \frac{\pi}{6}$$

$$\Rightarrow x^2 - 6x + 8.5 = \sin \frac{\pi}{6} = \frac{1}{2}$$

$$\Rightarrow x^2 - 6x + 8 = 0$$

$$\Rightarrow (x-4)(x-2) = 0 \Rightarrow x = 4 \text{ or } x = 2$$

38. (b) : $\sin^{-1}(x^2 - 7x + 12) = 0$

$$\Rightarrow x^2 - 7x + 12 = \sin 0$$

$$\Rightarrow x^2 - 7x + 12 = 0 \Rightarrow (x-4)(x-3) = 0 \Rightarrow x = 4, 3$$

39. (a) : We have, $\tan^{-1}(x^2 - 4x + 4) = 0$

$$\Rightarrow x^2 - 4x + 4 = \tan 0 \Rightarrow x^2 - 4x + 4 = 0$$

$$\Rightarrow (x-2)^2 = 0 \Rightarrow x = 2$$

40. (c) : $\tan^{-1}(\cot \theta) = 20 \Rightarrow \cot \theta = \tan 20$

$$\Rightarrow \cot \theta = \cot \left(\frac{\pi}{2} - 20 \right) \Rightarrow \theta = \frac{\pi}{2} - 20$$

$$\Rightarrow 30 = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{6}$$

41. (d) : Principal value branch of $\sec^{-1} x$ is $[0, \pi] - \left[\frac{\pi}{2} \right]$

$$42. (d) : \cos \left(2 \cos^{-1} \left(\frac{2}{5} \right) \right) = \cos 2x, \text{ where } x = \cos^{-1} \frac{2}{5}$$

$$= 2 \cos^2 x - 1 = 2 \left(\frac{2}{5} \right)^2 - 1 \quad \left(\because \cos x = \frac{2}{5} \right)$$

$$= \frac{2 \times 4}{25} - 1 = \frac{8 - 25}{25} = -\frac{17}{25}$$

$$43. (d) : \sin \left(\frac{\pi}{3} + \sin^{-1} \left(\frac{1}{2} \right) \right)$$

$$= \sin \left(\frac{\pi}{3} + \frac{\pi}{6} \right) = \sin \frac{\pi}{2} = 1$$

44. (a) : Let $\cot^{-1}(-\sqrt{3}) = \theta \Rightarrow \cot \theta = -\sqrt{3} = -\cot \frac{\pi}{6}$

$$= \cot \left(\pi - \frac{\pi}{6} \right) = \cot \frac{5\pi}{6} \Rightarrow \theta = \frac{5\pi}{6} \in (0, \pi)$$

\therefore Principal value of $\cot^{-1}(-\sqrt{3})$ is $\frac{5\pi}{6}$.

45. (a) : $\tan^{-1}(1) + \tan^{-1}(0) + \tan^{-1}(-1)$

$$= \frac{\pi}{4} + \pi - \frac{\pi}{4} = \pi$$

46. (c) : Since, $\tan^{-1} x$ and $\cot^{-1} x$ exists for all $x \in R$ and $\cos^{-1}(2-x)$ exists, if $-1 \leq 2-x \leq 1 \Rightarrow 1 \leq x \leq 3$

$\therefore \tan^{-1} x - \cot^{-1} x = \cos^{-1}(2-x)$ is possible only if $1 \leq x \leq 3$

Thus, the solution of given equation is $[1, 3]$.

47. (d) : We have, $\cos^{-1} [\cos(2 \cot^{-1}(\sqrt{3}))]$

$$= \cos^{-1} \left[\cos 2 \left(\frac{\pi}{6} \right) \right]$$

$$= \cos^{-1} \left(\cos \left(\frac{\pi}{3} \right) \right) = \cos^{-1} \left(\frac{1}{2} \right) = \frac{\pi}{3}$$

48. (a) : $2(\sin^{-1} x)^2 - 5(\sin^{-1} x) + 2 = 0$

$$\Rightarrow \sin^{-1} x = \frac{5 \pm \sqrt{25-16}}{4}$$

$$\Rightarrow \sin^{-1} x = \frac{1}{2}, \sin^{-1} x = 2$$

$$\Rightarrow x = \frac{\pi}{6} \text{ is only solution}$$

$[\because \frac{-\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}, \sin^{-1} x = 2 \text{ is not possible}]$

49. (d) : In $[0, \pi]$,

$$\cos^{-1} \left(\cos \left(\frac{9\pi}{4} \right) \right) = \cos^{-1} \left(\cos \left(2\pi + \frac{\pi}{4} \right) \right)$$

$$= \left(\cos^{-1} \left(\cos \frac{\pi}{4} \right) \right) = \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) = \frac{\pi}{4}$$


50. (d) : In $\left[\frac{-\pi}{2}, \frac{\pi}{2} \right]$,

$$\sin^{-1} \left(\sin \left(\frac{5\pi}{3} \right) \right) = \sin^{-1} \left(\sin \left(2\pi - \frac{\pi}{3} \right) \right)$$

$$= \sin^{-1} \left(\sin \left(\frac{-\pi}{3} \right) \right) = \frac{-\pi}{3}$$

51. (b) : We have, $BD = 30 \text{ m}$,

$AC = 40\sqrt{3} \text{ m}$, $AD = 30\sqrt{3} \text{ m}$,



Clearly, $BD \perp AC$

∴ In right $\triangle ADB$,

$$AB^2 = AD^2 + BD^2 = (30\sqrt{3})^2 + (30)^2 = 3600$$

$$\Rightarrow AB = 60 \text{ m}$$

$$\text{Now, } \sin \alpha = \frac{DB}{AB} = \frac{30}{60} = \frac{1}{2}$$

$$\Rightarrow \alpha = \sin^{-1}\left(\frac{1}{2}\right) \Rightarrow \angle CAB = \alpha = \sin^{-1}\left(\frac{1}{2}\right)$$

$$52. \text{ (c)} : \text{In right } \triangle ADB, \cos \alpha = \frac{AD}{AB} = \frac{30\sqrt{3}}{60} = \frac{\sqrt{3}}{2}$$
$$\Rightarrow \angle CAB = \alpha = \cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$$

$$53. \text{ (d)} : \text{Clearly, } CD = AC - AD = 40\sqrt{3} - 30\sqrt{3} = 10\sqrt{3} \text{ m}$$

$$\therefore \tan \beta = \frac{BD}{CD} = \frac{30}{10\sqrt{3}} = \frac{3}{\sqrt{3}} = \sqrt{3}$$

$$\Rightarrow \angle BCA = \beta = \tan^{-1}(\sqrt{3})$$

$$54. \text{ (c)} : \sin \alpha = \frac{1}{2} \Rightarrow \alpha = \frac{\pi}{6}; \tan \beta = \sqrt{3} \Rightarrow \beta = \frac{\pi}{3}$$

$$\therefore \angle ABC = \pi - (\alpha + \beta) = \pi - \left(\frac{\pi}{6} + \frac{\pi}{3}\right) = \frac{\pi}{2}$$

55. (c) : Domain and range of $\cos^{-1}x$ are $[-1, 1]$ and $[0, \pi]$ respectively.

56. (b) : Let $\angle CAB = \alpha$, then $\angle DAB = 2\alpha$ and $\angle EAB = 3\alpha$
In right $\triangle CAB$,

$$\tan \alpha = \frac{BC}{AB} = \frac{10}{20} = \frac{1}{2} \Rightarrow \angle CAB = \alpha = \tan^{-1}\left(\frac{1}{2}\right)$$

$$57. \text{ (c)} : \angle DAB = 2\alpha = 2\tan^{-1}(1/2)$$

$$58. \text{ (d)} : \angle EAB = 3\alpha = 3\tan^{-1}(1/2)$$

59. (b) : In right $\triangle CA'B$,

$$\tan \angle CA'B = \frac{BC}{A'B} = \frac{10}{25} = \frac{2}{5} \Rightarrow \angle CA'B = \tan^{-1}\left(\frac{2}{5}\right)$$

$$\therefore \text{Required difference} = \tan^{-1}\left(\frac{1}{2}\right) - \tan^{-1}\left(\frac{2}{5}\right)$$

60. (c) : Domain and range of $\tan^{-1}x$ are R and $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ respectively.

61. (a) : Let $y = \sin^{-1}(\sin(\pi - \pi/3))$

$$= \sin^{-1}\left(\sin \frac{\pi}{3}\right) = \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{3}$$
$$\left[\because \sin^{-1} : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \right]$$

62. (c) : We have, $f(x) = \sin^{-1}x + \tan^{-1}x + \sec^{-1}x$

Clearly, domain of $f(x)$ is given by $x = \pm 1$.

Thus, the range is $\{f(-1), f(1)\}$, i.e., $\left\{-\frac{\pi}{4}, \frac{3\pi}{4}\right\}$.

63. (a) : Reason is correct, from which we can say $\cot^{-1}x + \cos^{-1}2x = -\pi$ is not possible. Hence, both the Statements are correct and Statement-II is the correct explanation of Statement-I.

64. (d) : Let $\theta = \cot^{-1}(2x - x^2)$, where $\theta \in (0, \pi)$

$$\Rightarrow \cot \theta = 2x - x^2, \text{ where } \theta \in (0, \pi)$$

$$= 1 - (1 - 2x + x^2), \text{ where } \theta \in (0, \pi)$$

$$= 1 - (1 - x)^2, \text{ where } \theta \in (0, \pi)$$

$$\Rightarrow \cot \theta \leq 1, \text{ where } \theta \in (0, \pi)$$

$$\Rightarrow \frac{\pi}{4} \leq \theta < \pi \Rightarrow \text{Range of } f(x) \text{ is } \left[\frac{\pi}{4}, \pi\right)$$

65. (d) : $\sin^{-1}(x)$ is defined if $-1 \leq x \leq 1$.

But $1 + x^2 \geq 1$ and equality holds when $x = 0$

∴ Domain of $f(x)$ is $\{0\}$.

66. (c) : Let $2\tan^{-1}(0.75) = 0 \Rightarrow 0.75 = \tan\left(\frac{\theta}{2}\right)$

$$\therefore \sin(2\tan^{-1}(0.75))$$

$$= \sin 0 = \frac{2 \tan \theta / 2}{1 + \tan^2 \theta / 2} = \frac{2 \times 0.75}{1 + (0.75)^2} = \frac{1.50}{1.5625} = 0.96$$

67. (d) : Clearly, Statement-II is correct statement.

$$\text{Now, consider } \cos\left(\frac{\pi}{3} - \cos^{-1}\left(\frac{1}{2}\right)\right) = \cos\left(\frac{\pi}{3} - \frac{\pi}{3}\right) \\ = \cos 0 = 1$$

∴ Statement-I is wrong statement.

68. (b) : We have, $4\tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = 4 \cdot \frac{\pi}{6} = \frac{2\pi}{3}$.

∴ Statement-I is correct statement.

Thus, Statement-I and Statement-II are correct statements but Statement-II is not the correct explanation of Statement-I.

69. (c) : Clearly, $f(x)$ will be defined, if $0 \leq x - 1 \leq 1$

$$\Rightarrow 1 \leq x \leq 2$$

∴ Domain of $f(x)$ is $[1, 2]$.

∴ Statement-I is true statement.

But, Statement-II is false statement.

70. (a) : Let $f(x) = \sin^{-1}x + \cos^{-1}x + \tan^{-1}x$. Then

$$\text{Dom } f = [-1, 1] \cap [-1, 1] \cap R = [-1, 1].$$

∴ Both Statements are correct and Statement-II is the correct explanation of Statement-I.



MONTHLY TEST DRIVE



This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.

Total Marks : 80

Series 3 : Matrices and Determinants

Time Taken : 60 Min.

Only One Option Correct Type

1. Let $\Delta(x) = \begin{vmatrix} (x-2) & (x-1)^2 & x^3 \\ (x-1) & x^2 & (x+1)^3 \\ x & (x+1)^2 & (x+2)^3 \end{vmatrix}$, then the coefficient of x in $\Delta(x)$ is
- (a) -3
 - (b) -2
 - (c) -1
 - (d) 0

2. Let $A + 2B = \begin{bmatrix} 1 & 2 & 0 \\ 6 & -3 & 3 \\ -5 & 3 & 1 \end{bmatrix}$ and $2A - B = \begin{bmatrix} 2 & -1 & 5 \\ 2 & -1 & 6 \\ 0 & 1 & 2 \end{bmatrix}$.

Then $\text{tr}(A) - \text{tr}(B)$ has the value equal to

- (a) 0
- (b) 1
- (c) 2
- (d) 3

3. Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}$. If $10A^{10} + \text{adj}(A^{10}) = B$, then $b_1 + b_2 + b_3 + b_4$ is equal to
- (a) 91
 - (b) 92
 - (c) 111
 - (d) 112

4. If a, b, c are in G.P. and $(x - \alpha)$ is a factor of $ax^2 + 2bx + c = 0$, then value of the determinant

$$\begin{vmatrix} a & b & a\alpha+b \\ b & c & b\alpha+c \\ a\alpha+b & b\alpha+c & 0 \end{vmatrix}$$

is

- (a) 0
- (b) 1
- (c) 2
- (d) 3

5. If $A^5 = O$ such that $A^n \neq I$ for $1 \leq n \leq 4$, then $(I - A)^{-1}$ is equal to
- (a) A^4
 - (b) A^3
 - (c) $I + A$
 - (d) None of these

6. If $D = \text{diag}(d_1, d_2, d_3, \dots, d_n)$, where $d_i \geq 0 \forall i$, then D^{-1} equals
- (a) D
 - (b) $\text{diag}(d_1^n, d_2^n, \dots, d_n^n)$
 - (c) I_n
 - (d) $\text{diag}(d_1^{-1}, d_2^{-1}, d_3^{-1}, \dots, d_n^{-1})$

One or More than One Option(s) Correct Type

7. If $f(x) = \begin{vmatrix} a & -1 & 0 \\ ax & a & -1 \\ ax^2 & ax & a \end{vmatrix}$, then $f(2x) - f(x)$ is divisible by
- (a) x
 - (b) a
 - (c) $2a + 3x$
 - (d) x^2

PUZZLE CORNER

ANSWER - AUGUST 2021



1-	2-	2-	3+	1	2-
6	4	5	2	1	3
5	6	3	2+	2	1
2-	4+	1	2	11+	6
4				3	16+
2	10x	5	1	3	4
9x	3	2	10+	5-	1-
			6	4	5
1	3	4	6	3-	2

8. If $A + B + C = \pi$, $e^{i\theta} = \cos\theta + i \sin\theta$ and

$$z = \begin{vmatrix} e^{2iA} & e^{-iC} & e^{-iB} \\ e^{-iC} & e^{2iB} & e^{-iA} \\ e^{-iB} & e^{-iA} & e^{2iC} \end{vmatrix}, \text{ then}$$

- (a) $\operatorname{Re}(z) = 4$ (b) $\operatorname{Im}(z) = 0$
 (c) $\operatorname{Re}(z) = -4$ (d) $\operatorname{Im}(z) = -1$

9. Let ω be a complex cube root of unity with $\omega \neq 1$ and $P = [p_{ij}]$ be a $n \times n$ matrix with $p_{ij} = \omega^{i+j}$. Then $P^2 \neq 0$, when $n =$

- (a) 57 (b) 55 (c) 58 (d) 56

10. Let $a, \lambda, \mu \in R$. Consider the system of linear equations $ax + 2y = \lambda$, $3x - 2y = \mu$. Which of the following statement(s) is/are correct?

- (a) If $a = -3$, then the system has infinitely many solutions for all values of λ and μ .
 (b) If $a \neq -3$, then the system has a unique solution for all values of λ and μ .
 (c) If $\lambda + \mu = 0$, then the system has infinitely many solutions for $a = -3$.
 (d) If $\lambda + \mu \neq 0$, then the system has no solution for $a = -3$.

11. Let $A = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix}$ be a matrix. If $A^{10} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then

- (a) number of divisors of a is 11
 (b) a is an odd integer
 (c) $(a + b + d)$ is an even integer
 (d) $a + d$ is a multiple of 13

12. The value of θ for which the system of linear equations in x, y, z given as

$$(\sin 30)x - y + z = 0$$

$$(\cos 20)x + 4y + 3z = 0$$

$$2x + 7y + 7z = 0$$

has a non-trivial solution, is/are

- (a) $\frac{n\pi}{2}$ (b) $\frac{2n\pi}{3}$
 (c) $n\pi + (-1)^n \frac{\pi}{6}$ (d) None of these

13. If A is symmetric and B is skew symmetric matrix, then which of the following is/are not true?

- (a) ABA^T is symmetric matrix
 (b) $AB^T + BA^T$ is symmetric matrix
 (c) $(A + B)(A - B)$ is skew symmetric
 (d) $(A + I)(B - I)$ is skew symmetric

ATTENTION
COACHING
INSTITUTES :
a great offer from
MTG

CLASSROOM STUDY MATERIAL

MTG offers "Classroom Study Material" for JEE (Main & Advanced), NEET and FOUNDATION MATERIAL for Class 6, 7, 8, 9, 10, 11 & 12 **with YOUR BRAND NAME & COVER DESIGN.**


This study material will save your lots of money spent on teachers, typing, proof-reading and printing. Also, you will save enormous time. Normally, a good study material takes 2 years to develop. But you can have the material printed with your logo delivered at your doorstep.

Profit from associating with MTG Brand – the most popular name in educational publishing for JEE (Main & Advanced)/NEET

Order sample chapters on Phone/Fax/e-mail.

Phone : 0124-6601200 09312680856

e-mail : sales@mtg.in | www.mtg.in



Comprehension Type

Paragraph for Q. No. 14 and 15

If $x > m, y > n, z > r$ ($x, y, z > 0$) such that $\begin{vmatrix} x & n & r \\ m & y & r \\ m & n & z \end{vmatrix} = 0$.

14. The value of $\frac{x}{x-m} + \frac{y}{y-n} + \frac{z}{z-r}$ is

- (a) 2 (b) -4
 (c) 0 (d) -1

15. The greatest value of $\frac{xyz}{(x-m)(y-n)(z-r)}$ is

- (a) 27 (b) 8/27
 (c) 64/27 (d) None of these

Matrix Match Type

16. Match the following:

	Column-I	Column-II
P.	If ω is a non real cube root of unity, then a root of the following equation $\begin{vmatrix} x+1 & \omega & \omega^2 \\ \omega & x+\omega^2 & 1 \\ \omega^2 & 1 & x+\omega \end{vmatrix} = 0$ is	1. 0
Q.	If $A = \begin{bmatrix} 2 & 4 & 5 \\ 4 & 8 & 10 \\ -6 & -12 & -15 \end{bmatrix}$, then rank of A is	2. -2
R.	Suppose that system of equations $x = cy + bz, y = az + cx, z = bx + ay$ has non trivial solution. Then $a^2 + b^2 + c^2 + 2abc =$	3. 1

S.	$\begin{vmatrix} a+1 & a+2 & a+4 \\ a+3 & a+5 & a+8 \\ a+7 & a+10 & a+14 \end{vmatrix}$	4. 4
----	---	------

P	Q	R	S
(a) 1	3	3	2
(b) 2	4	3	1
(c) 4	2	3	1
(d) 4	1	2	3

Numerical Answer Type

17. In a ΔABC , if $\begin{vmatrix} 1 & a & b \\ 1 & c & a \\ 1 & b & c \end{vmatrix} = 0$, then the value of $64(\sin^2 A + \sin^2 B + \sin^2 C)$ must be _____.

18. Let S be the set which contains all possible values of l ,

$$m, n, p, q, r \text{ for which } A = \begin{bmatrix} l^2 - 3 & p & 0 \\ 0 & m^2 - 8 & q \\ r & 0 & n^2 - 15 \end{bmatrix}$$

be a non singular idempotent matrix. Then the sum of all the elements of the set S is _____.

19. If A, B and C are $n \times n$ matrices and $\det(A) = 2$, $\det(B) = 3$ and $\det(C) = 5$, then the value of $[\det(A^2BC^{-1})]$ (where $[\cdot]$ represents greatest integer function) is _____.

20. If $M = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$ and $M^2 - \lambda M - I_2 = O$, λ must be _____.



Keys are published in this issue. Search now! ☺

SELF CHECK

Check your score! If your score is

> 90% EXCELLENT WORK ! You are well prepared to take the challenge of final exam.

90-75% GOOD WORK ! You can score good in the final exam.

74-60% SATISFACTORY ! You need to score more next time.

< 60% NOT SATISFACTORY! Revise thoroughly and strengthen your concepts.

No. of questions attempted
 No. of questions correct
 Marks scored in percentage

Now, savings of up to ₹920* with MTG's magazine subscription plans!

*On cover price of ₹40/-.

Our new offers are here!

Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to www.mtg.in now to subscribe online.

About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines (**Physics For You**, **Chemistry Today**, **Mathematics Today** & **Biology Today**) ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

For JEE
(Main &
Advanced),
NEET and
BOARDS



Over 1.2 Cr readers. Since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team comprising experts and scholars from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement – confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET/AIIMS toppers
- Bonus: Exposure to competition at a global level, with questions from International Olympiads & Contests

Lifetime Subscription Plan for teachers, and special schemes and offers available for libraries and coaching institutes.
SMS MTG to 8800255334 to learn more.

SUBSCRIPTION FORM

Confirm your choice by placing tick-marks in relevant boxes.

Plan 1:	<input type="checkbox"/> Physics	27 months	15 months	9 months
Individual magazines P, C, M, B	<input type="checkbox"/> Chemistry	₹850	₹500	₹300
	<input type="checkbox"/> Mathematics	(save ₹ 230)	(save ₹ 100)	(save ₹ 60)
	<input type="checkbox"/> Biology			

Plan 2:	<input type="checkbox"/> PCM	₹2500	₹1400	₹900
Combo of 3	<input type="checkbox"/> PCB	(save ₹ 740)	(save ₹ 400)	(save ₹ 180)

Plan 3:	<input type="checkbox"/> PCMB Combo	₹3400	₹1900	₹1200
		(save ₹ 920)	(save ₹ 500)	(save ₹ 240)

Courier Charges Add to your subscription amount for quicker & reliable delivery	<input type="checkbox"/> ₹600	<input type="checkbox"/> ₹450	<input type="checkbox"/> ₹240
---	-------------------------------	-------------------------------	-------------------------------

Recommended by (Optional)

Name of your teacher _____
Teacher's Mobile #

Note: Magazines are despatched by Book Post on 4th of every month (each magazine separately).

E-mail subscription@mtg.in. Visit www.mtg.in to subscribe online. Call (0)1800-10-38673 for more info.
Get digital editions of MTG Magazines on <http://digital.mtg.in/>

Enclose Demand Draft favouring
MTG Learning Media (P) Ltd, payable at New Delhi.
Mail this Subscription Form to Subscription Dept.,
MTG Learning Media (P) Ltd, Plot 99, Sector 44, Gurgaon-122 003 (H.R.).

Find all your **QUESTIONS** here, before getting them in your exam.


By not studying
MORE but
ACCURATE...



Our Books



₹800/-



₹800/-



₹800/-

JEE 2021 exams are nearing and all the students must be in the stage of preparation with full dedication and hard work. But have you put your knowledge to test? With our 39 years of experience, we would suggest you to put yourself through MTG's JEE Champion Series so that you can be sure about your level of preparation and find your core strengths and capabilities. The best part is you study at a pace you're comfortable with. Because it's all chapters, topicwise.

WWW.JEEBOOKS.IN



JEE (MAIN) 2021

PHASE-III

OUTSTANDING RESULTS YET AGAIN!

7
State
Toppers

3
100
Percentilers

388
99+
Percentilers

1451
95+
Percentilers



PRAVAR KATARIA
5 YR. CLASSROOM

100

Percentile



ANMOL ARICHWAL
4 YR. CLASSROOM

OUR STATE TOPPERS

DELHI

HARYANA

UTTAR PRADESH

CHANDIGARH

GUJARAT

PUNJAB

KERALA



100
PERCENTILE



100
PERCENTILE



100
PERCENTILE



99.99
PERCENTILE



99.99
PERCENTILE



99.99
PERCENTILE



99.99
PERCENTILE

PRAVAR KATARIA
5 YR. CLASSROOM

ANMOL ARICHWAL
4 YR. CLASSROOM

PAL AGGARWAL
1 YR. DISTANCE

CHAITANYA AGGARWAL
2 YR. DISTANCE

PARTH PATEL
2 YR. DISTANCE

VAIBHAV BAJAJ
5 YR. CLASSROOM

SREEHARI C
2 YR. CLASSROOM

SUBJECT-WISE
PERCENTILE SCORERS

PHYSICS
51

CHEMISTRY
16

MATHS
15

Though every care has been taken to publish the result correctly, yet the Institute shall not be responsible for error, if any.

ADMISSIONS OPEN



Hybrid Course for Power Step (JEE) / Medical (NEET) with Learning Device (Tablet) | Repeater / XII Passed Batches



Flexibility to shift
to classroom at no
additional cost



Continue your learning
with Tablet (8") 10.1"
4G | WiFi Enabled



Pre-activated
SIM (if opted
for SIM card)



Easy financing /
No Cost EMI
options available

WWW.JEEBOOKS.IN

Download Akash App



CALL (TOLL-FREE)

1800-102-2727

VISIT

akash.ac.in

