Complex Analysis Lecture Notes

Hand written summary from lectures

Acknowledgment

Special thanks to my professor MR.BAKIR FARHI, who gave the lectures and explanations, this work wouldn't exist without his teaching. here is the link to his website:

http://farhi.bakir.free.fr/home/index-fr.html

Disclaimer

These notes were written in real-time during the lectures, this is not the final version, yet. so they may contain:

- Incomplete or incorrect information.
- Typos, transcription mistakes, or missing content.
- Interpretations or notations that reflect my own understanding. at the moment

Please double check anything important with official material or trusted sources.

If you spot an error feel free	e to open an issue or submit a pull request, or contact me via gmail:
	kara.abderahmane@nhsm.edu.dz
Notes on Contribution :	
	ative effort. students who contribute by reporting errors or helping
	be credited in the next page as contributors in future versions, your s improve this document for everyone.
neip is appreciated and neip	s improve this document for everyone.

My Github Page Last Update: 2025-10-11			
Main Drawer		Haddar Noureddine	
News Reporter		Hammiche Ismail	
	ARMWRESTLING4EVER		

Contents 1 Power Series 2 5 10 1.2.1 1.3.1 13 1.3.2 16 23 1.5.1 24 1.5.2 28 2 The Cauchy integral formula on a circle and applications 31

Chapter 1

Power Series

Lecture 1

08:06 AM Mon, Sep 29 2025

Definition 1.0.1 (Power Series): A power series is a formal series of the form $\sum_{n=0}^{\infty} a_n z^n$, where $a_n \in \mathbb{C}$ for all $n \in \mathbb{N}_0$.

More generally, given $z_0 \in \mathbb{C}$, a power series centered at z_0 is a formal series of the form:

$$\sum_{n=0}^{\infty} a_n (z-z_0)^n,$$

where $a_n \in \mathbb{C} \quad (\forall n \in \mathbb{N}_0)$

Remark 🐿

The set of all complex power series (centered at 0) is denoted by $\mathbb{C}[[z]]$. More generally, given $z_0 \in \mathbb{C}$, the set of all complex power series centered at z_0 is denoted by $\mathbb{C}[[z-z_0]]$.

Operations on Formal Power Series:

Given $z_0 \in \mathbb{C}$, we equip $\mathbb{C}[[z-z_0]]$. with the following operations:

① **Additions:** For all $(a_n)_{n\in\mathbb{N}_0}$, $(b_n)_{n\in\mathbb{N}_0}\subset\mathbb{C}$:

$$\sum_{n=0}^{\infty} a_n (z-z_0)^n + \sum_{n=0}^{\infty} b_n (z-z_0)^n = \sum_{n=0}^{\infty} (a_n + b_n) (z-z_0)^n.$$

2 Multiplication

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \times \sum_{n=0}^{\infty} b_n (z - z_0)^n = \sum_{n=0}^{\infty} c_n (z - z_0)^n,$$

where, $c_n := \sum_{k=0}^n a_k b_{n-k}$ for all $n \in \mathbb{N}_0$. Also $(c_n)_{n \in \mathbb{N}}$ is called the covolution of the two sequences $(a_n)_{n \in \mathbb{N}_0}$ and $(b_n)_{n \in \mathbb{N}_0}$.

③ Scalar Multiplication: For all $\lambda \in \mathbb{C}$, and all $(a_n)_{n \in \mathbb{N}_0} \subset \mathbb{C}$:

$$\lambda \sum_{n=0}^{\infty} a_n (z-z_0)^n = \sum_{n=0}^{\infty} (\lambda a_n) (z-z_0)^n.$$

It's straightforward to verify that $\mathbb{C}[[z-z_0]]$ equipped with these operations forms a commutative algebra over \mathbb{C} . The Multiplicative identity is the constant power series:

$$1 = 1 + 0 \cdot (z - z_0) + 0 \cdot (z - z_0)^2 + \dots$$

Definition 1.0.2 (Domain of Convergence) : The domain of convergence of a power series $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ is the set of all points $z \in \mathbb{C}$ for which the series converge. The structure of this domain is very specific. Its a disk (possibly with some points in its boundary) centered at z_0 .

Proposition 1.0.1 (Abel's Lemma): Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series and let $z_1 \in \mathbb{C} \setminus \{z_0\}$. Suppose that the sequence $\{a_n(z_1-z_0)^n\}_{n\in\mathbb{N}_0}$ is bounded. Then, the power series in question converges absolutely (so converges) for every $z\in\mathbb{C}$, such that:

$$|z-z_0|<|z_1-z_0|$$

Proof. By hypothesis, $\exists M > 0$ such that $\forall n \in \mathbb{N}_0$:

$$|a_n(z_1-z_0)^n| \le M$$

Then, for all $z \in \mathbb{C}$ such that $|z - z_0| < |z_1 - z_0|$ we have:

$$|a_n(z - z_0)^n| = \underbrace{|a_n(z_1 - z_0)^n|}_{\leq M} \cdot \underbrace{\left|\frac{z - z_0}{z_1 - z_0}\right|^n}_{\leq 1}$$

$$\leq M \underbrace{\left|\frac{z - z_0}{z_1 - z_0}\right|^n}_{\leq 1}.$$

Since $\left|\frac{z-z_0}{z_1-z_0}\right| < 1$ then the geometric series

$$\sum_{n=0}^{\infty} M \left| \frac{z - z_0}{z_1 - z_0} \right|^n \text{ Converges }.$$

Thus, the series $\sum_{n=0}^{\infty} |a_n(z-z_0)^n|$ also converges, that is $\sum_{n=0}^{\infty} a_n(z-z_0)^n$ is absolutely convergent.

Corollary 1.0.2 : Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series which converges at some $z=z_1 \in \mathbb{C} \setminus \{z_0\}$. Then the power series in question converges absolutely (so converges), for every $z \in \mathbb{C}$ such that:

 $\hat{z_0}$

$$|z - z_0| < |z - z_1|$$

Proof. $\sum_{n=0}^{\infty} a_n (z_1 - z_0)^n$ converges implies that $a_n (z - z_0)^n \to 0$ as $n \to +\infty$, which implies that the sequence $\{a_n (z_1 - z_0)^n\}_{n \ge 0}$ is bounded. *Proposition 1.0.1* permits us to conclude the required result.

Theorem 1.0.3 (Radius of Convergence) : Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series. Then there exists a unique $R \in [0, \infty]$, called the radius of convergence with the following properties:

- ① The power series converges absolutely for every $z \in \mathbb{C}$ satisfying $|z z_0| < R$.
- ② The power series diverges for every $z \in \mathbb{C}$ satisfying $|z-z_0| > R$. The disk $D(z_0,R)=\{z\in\mathbb{C}: |z-z_0|< R\}$ is called the disk of convergence.

Proof. Define the set $A \subset \mathbb{R}_{\geq 0}$ of nonegative real numbers for which the sequence $\{|a_n| r^n\}_{n \in \mathbb{N}_0}$ is bounded.

$$A:=\left\{ r\geq 0: \quad \sup_{n\in\mathbb{N}_0}\left|a_n\right|r^n<\infty
ight\}$$

we have $A \neq \emptyset$ because $0 \in A$. Define $R := \sup A \in [0, \infty]$, we now show that R has the stated properties.

- Let $z \in D(z_0, R)$. By definition of the supremum, there exists $r \in A$, (i.e., $|a_n| r^n$ is bounded) such that $|z z_0| < r \le R$. Since $|z z_0| < r$ and $\{|a_n| r^n\}_{n \ge 0}$ is bounded, then by Abel's lemma, we deduce that the series $\sum_{n=0}^{\infty} a_n (z z_0)^n$ converges absolutely.
- **◆**② Let $z ∈ \mathbb{C}$ such that $|z z_0| > R$, suppose for contradictions that the power series converges at z. Then by the *Corollary 1.0.2*, it would converge absolutely for any ω with $|ω z_0| < 1$

 $|z - z_0|$. In particular, for any r such that:

$$R < r < |z - z_0|$$

the series would converge at points on the circle $C(z_0, r)$, implying $r \in A$. This contradicts the fact that $R = \sup A$. Therefore, the power series diverges.

→ The Uniqueness of R:

If another $R' \in [0, \infty]$ satisfies the same properties, a point z such that $|z - z_0|$ lies between R and R' would lead to a contradiction regarding the convergence or divergence of the power series.

1.1 Formulas for Calculating the Radius of Convergence

Proposition 1.1.1 (Hadamard's Formula) : Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series centered at $z_0 \in \mathbb{C}$. Denote by R its radius of convergence. Then:

$$\frac{1}{R} = \lim_{n \to \infty} \sup \sqrt[n]{|a_n|}$$

with the convention $\frac{1}{0} = \infty$ and $\frac{1}{\infty} = 0$

Proof. Let $L := \lim_{n \to \infty} \sup |a_n|^{\frac{1}{n}} \in [0, \infty]$. We must show that $R = \frac{1}{L}$. Let $z \in \mathbb{C} \setminus \{z_0\}$, we distinguish three cases:

•• ① If L = 0. In this case, we have:

$$0 \le \lim_{n \to \infty} \inf |a_n|^{\frac{1}{n}} \le \lim_{n \to \infty} \sup |a_n|^{\frac{1}{n}} = 0$$

Thus, $\lim_{n\to\infty}\inf|a_n|^{\frac{1}{n}}=\lim_{n\to\infty}\sup|a_n|^{\frac{1}{n}}=0$. This implies that $\lim_{n\to\infty}|a_n|^{\frac{1}{n}}$ exists and equals to 0, so for all n sufficiently large, we have:

$$|a_n|^{\frac{1}{n}} < \frac{1}{2|z-z_0|};$$

That is,

$$|a_n(z-z_0)^n|<\frac{1}{2^n}.$$

Since the geometric series $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges then the series $\sum_{n=0}^{\infty} |a_n(z-z_0)^n|$ converges $\forall z \in \mathbb{C}$, thus $R = +\infty = \frac{1}{L}$

• ② If $L = +\infty$, we have $L = \lim_{n \to \infty} \sup |a_n|^{\frac{1}{n}} = +\infty$ is equivalent to the fact that the sequence $\left\{|a_n|^{\frac{1}{n}}\right\}_{n \in \mathbb{N}}$ is bounded. Therefore, the sequence:

$$|a_n(z-z_0)^n|^{\frac{1}{n}} = |a_n|^{\frac{1}{n}} |z-z_0|$$

is also unbounded. This implies that $|a_n(z-z_0)^n|$ is unbounded, thus $|a_n(z-z_0)^n|$ does not converge to 0 as $n \to \infty$. Hence $\sum_{n=0}^{\infty} a_n(z-z_0)^n$ diverges. Hence R=0.

- •• ③ If $L \in (0, \infty)$. Let $z \in \mathbb{C}$. We consider two subcases:
 - If $|z-z_0| < \frac{1}{L}$. Choose r such that $|z-z_0| < r < \frac{1}{L}$, thus $L < \frac{1}{r}$. By defintion of a $\lim_{n\to\infty} \sup$, for all n sufficiently large we have:

$$|a_n|^{\frac{1}{n}}<\frac{1}{r},$$

which implies that:

$$|a_n(z-z_0)^n| < \underbrace{\left(\frac{|z-z_0|}{r}\right)^n}_{\leq 1}.$$

Since $\left|\frac{z-z_0}{r}\right| < 1$, the geometric series $\sum_{n=0}^{\infty} \left|\frac{z-z_0}{r}\right|^n$ converges. By comparison, the power series $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ converges absolutely.

2 If $(|z-z_0| > \frac{1}{L})$. In this case, we have:

$$\lim_{n \to \infty} \sup |a_n (z - z_0)^n|^{\frac{1}{n}} = \lim_{n \to \infty} \sup \left(|a_n|^{\frac{1}{n}} |z - z_0| \right)$$
$$= L |z - z_0| > 1$$

Thus, $\{a_n(z-z_0)^n\}_{n\in\mathbb{N}}$ is unbounded, hence $|a_n(z-z_0)^n|$ does not converge to zero as $n\to\infty$, implying that $\sum_{n=0}^{\infty}a_n(z-z_0)^n$ diverges. Therefore:

$$R=\frac{1}{L}$$
.

Lecture 2

08:00 AM Mon, Oct 06 2025

Proposition 1.1.2 (Ratio Test Formula) : Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series. Suppose that the limit

$$\alpha = \lim_{n \to \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists (i.e., $\in [0, \infty]$). Then the radius of convergence R of the power series in question is $R = \alpha$.

Proof. We use the d'Allembert rule for the series

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \qquad (z \in \mathbb{C} \setminus \{z_0\}).$$

Let $z \in \mathbb{C} \setminus \{z_0\}$. we have:

$$\lim_{n \to \infty} \left| \frac{a_{n+1}(z - z_0)^{n+1}}{a_n(z - z_0)^n} \right| = \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \cdot |z - z_0|$$
$$= |z - z_0| \cdot \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right|$$
$$= \frac{|z - z_0|}{\alpha}$$

By the d'Allembert rule, we have:

 \implies The series $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ converges if

$$\frac{|z-z_0|}{\alpha} < 1 \quad \text{i.e.} \quad |z-z_0| < \alpha.$$

 \implies The series $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ diverges if

$$\frac{|z-z_0|}{\alpha} > 1 \quad \text{i.e.} \quad |z-z_0| > \alpha.$$

Hence $R = \alpha$.

Example: Determine the radius of convergence of the power series $\sum_{n=0}^{\infty} \frac{z_n}{n!}$ where $z_0 = 0$.

1 st METHOD: (BY HADAMARD FORMULA)

We must compute $\lim_{n\to\infty} \sup\left(\frac{1}{n!}\right)^{\frac{1}{n}}$. By the stirling formula, we have that:

$$n! \sim_{+\infty} n^n e^{-n} \sqrt{2\pi n}$$
.

Thus we get:

$$(n!)^{\frac{1}{n}} \sim_{+\infty} ne^{-1} (2\pi n)^{\frac{1}{2n}}.$$

Thus

$$\left(\frac{1}{n!}\right)^{\frac{1}{n}} \sim_{+\infty} \frac{e}{n} (2\pi n)^{-\frac{1}{2n}} \to 0 \text{ as } n \to +\infty.$$

Thus $R = \frac{1}{0} = +\infty$.

This means that the power series $\sum_{n=0}^{\infty} \frac{z^n}{n!}$ converges for all $z \in \mathbb{C}$.

2 nd METHOD:

We use *Proposition 2* . we have:

$$\lim_{n \to \infty} \left| \frac{\frac{1}{n!}}{\frac{1}{(n+1)!}} \right| = \lim_{n \to \infty} \frac{(n+1)!}{n!}$$
$$= \lim_{n \to \infty} (n+1) = +\infty.$$

Thus $R = +\infty$

1.2 Analytic Functions


Definition 1.2.1: Let Ω be a non empty open subset of $\mathbb C$ and let $z_0 \in \Omega$.

Let $f: \Omega \longrightarrow \mathbb{C}$ be a map. then:

1. f is said to be analytic at z_0 if there exists r>0 and a complex sequence $(a_n)_{n\in\mathbb{N}_0}$ such that $D(z_0,r)\subset\Omega$ and:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (\forall z \in D(z_0, r)).$$

2. f is said to be analytic on Ω if its analytic at every point of Ω .



Example:

1. Every complex polynomial is analytic on \mathbb{C} . Indeed, let $P \in \mathbb{C}[\mathbb{Z}]$, and $z_0 \in \mathbb{C}$. since $P(z+z_0) \in \mathbb{C}[\mathbb{Z}]$, we can write:

$$P(z+z_0)=\sum_{n=0}^d a_n z^n \quad (d\in\mathbb{N}_0).$$

Substituting z by $(z - z_0)$, we get:

$$P(z) = \sum_{n=0}^{d} a_n (z - z_0)^n,$$

which is a power series centered at z_0 with infinite randius of convergence. Thus, P is analytic at z_0 . Since z_0 was arbitrary, P is analytic on \mathbb{C} .

2. The function $z \longrightarrow \frac{1}{z}$ is analytic on $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. Indeed, let $z_0 \in \mathbb{C}^*$ arbitrary. For $z \in D(z_0, |z_0|)$, we have:

$$\left|\frac{z-z_0}{z_0}\right|<1.$$

We can write

$$\frac{1}{z} = \frac{1}{z_0 + (z - z_0)}$$

$$= \frac{1}{z_0} \cdot \frac{1}{1 + \frac{z - z_0}{z_0}}$$

$$= \frac{1}{z_0} \cdot \sum_{n=0}^{\infty} (-1)^n \left(\frac{z - z_0}{z_0}\right)^n$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{z_0^{n+1}} (z - z_0)^n,$$

which is a power series centered at z_0 , valid on $D(z_0, |z_0|)$. Hence $z \longrightarrow \frac{1}{z}$ is analytic at z_0 . Since $z_0 \in \mathbb{C}^*$ was arbitrary, then $z \longrightarrow \frac{1}{z}$ is analytic on \mathbb{C}^* .

1.2.1 Properties of Analytic Functions

Proposition 1.2.1: Let Ω be a non empty open subset of \mathbb{C} and let $z_0 \in \Omega$. If $f,g:\Omega \to \mathbb{C}$ are analytic at z_0 , then the same is for (f+g) and $(f \cdot g)$. Moreover, if f and g are represented by power series with radii of convergence R_f and R_g respectively then (f+g) and $(f \cdot g)$ are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P and P are represented by power series with radii of convergence P are represented by P and P are represented by P and P are represented by P are represented by P and P are represented by P are represented by P and P are represented by P are represented by P and P are r

Proof. Exercise.

Corollary 1.2.2: Let Ω be a non empty open subset of \mathbb{C} and let $f,g:\Omega\longrightarrow\mathbb{C}$. If f and g are both analytic on Ω , then the same is for (f+g) and $(f\cdot g)$.

Proposition 1.2.3 (Analyticity \Longrightarrow **Continuity)**: Let Ω be a non empty open subset of $\mathbb C$ and let $z_0 \in \Omega$. Let also $f: \Omega \longrightarrow \mathbb C$ be a map. If f is analytic at z_0 then f is continuous at z_0

Proof. Suppose that f is analytic at z_0 then there exists R > 0 and a complex sequence $(a_n)_{n \in \mathbb{N}_0}$ such that $D(z_0, R) \subset \Omega$ and:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (\forall z \in D(z_0, R))$$

In particular, $f(z_0) = a_0$. Thus for all $z \in D(z_0, R)$ we have:

$$f(z) - f(z_0) = \sum_{n=1}^{\infty} a_n (z - z_0)^n$$

$$= (z - z_0) \sum_{n=1}^{\infty} a_n (z - z_0)^{n-1}$$

$$= (z - z_0) \sum_{n=0}^{\infty} a_{n+1} (z - z_0)^n$$
 (1)

By the Hadamard formula, we see that the power series $\sum_{n=0}^{\infty} a_{n+1}(z-z_0)^n$ has the same radius of convergence as the original power series $\sum_{n=0}^{\infty} a_n(z-z_0)^n$. Consequently, the power series $\sum_{n=0}^{\infty} a_{n+1}(z-z_0)^n$ converges absolutely for $|z-z_0| < R$. Let $r \in \mathbb{R}$ such that 0 < r < R. Then

for all $z \in D(z_0, r)$, we have from (1) the estimate:

$$|f(z) - f(z_0)| = |z - z_0| \cdot \left| \sum_{n=0}^{\infty} a_{n+1} (z - z_0)^n \right|$$

$$\leq |z - z_0| \sum_{n=0}^{\infty} |a_{n+1}| |z - z_0|^n$$

$$\leq |z - z_0| \sum_{n=0}^{\infty} |a_{n+1}| \cdot r^n.$$

$$< +\infty \text{ since } r < R$$

Taking the limit as $z \to z_0$, we conclude that $\lim_{z \to z_0} f(z) = f(z_0)$, so f is continuous at z_0 .

Corollary 1.2.4 (Immediate): Let Ω be a non empty open subset of \mathbb{C} and $f:\Omega\longrightarrow\mathbb{C}$. If f is analytic on Ω , then f is continuous on Ω .

Proposition 1.2.5 (Composition of Analytic functions): Let Ω_1 and Ω_2 be two nonempty open subsets of $\mathbb C$ and let $f:\Omega_1\longrightarrow\Omega_2$ and $g:\Omega_2\longrightarrow\mathbb C$ be two maps. Let also $z_0\in\Omega_1$. If f is analytic at z_0 and g is analytic at $f(z_0)$, then $(g\circ f)$ is analytic at z_0 .

Proof. Exercise

Corollary 1.2.6 (Immediate): Let Ω_1 and Ω_2 be two nonempty open subsets of $\mathbb C$ and let $f:\Omega_1\longrightarrow\Omega_2$ and $g:\Omega_2\longrightarrow\mathbb C$ be two maps. If f is analytic on Ω_1 and g is analytic on Ω_2 then $(g\circ f)$ is analytic on Ω_1 .

Proposition 1.2.7 (Quotient of Analytic Functions) : Let Ω be a nonempty open subsets of \mathbb{C} and let $z_0 \in \Omega$. Let also $f,g:\Omega \longrightarrow \mathbb{C}$ be two functions which are both analytic at z_0 and such that $g(z_0) \neq 0$. Then the function $\frac{f}{g}$ is analytic at z_0 .

Proof. Since $g(z_0) \neq 0$ then the function $h: w \longrightarrow \frac{1}{w}$ is analytic at $g(z_0)$ (as seen in previous examples). Therefore, by *Proposition* 1.2.5, the function $\frac{1}{g} = h \circ g$ is analytic at z_0 . It then follows from *Proposition* 1.2.1 that the product $f \cdot \left(\frac{1}{g}\right)$ is analytic at z_0 .

Corollary 1.2.8 (Immediate): Let Ω be a non empty open subset of $\mathbb C$ and let $f,g:\Omega\longrightarrow\mathbb C$ be two analytic functions on Ω such that $g(z)\neq 0$ for every $z\in\Omega$. Then the function $\frac{f}{g}$ is analytic on Ω .

Example: Every rational function is analytic on its domain of definition. This is because a rational function is a quotient of two polynomials, and polynomials are analytic on \mathbb{C} .

1.3 Power series define Analytic functions

Theorem 1.3.1: A power series with a positive radius of converges defines an analytic function on its disk of convergence.

Proof. Let $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ be a power series $(z_0 \in \mathbb{C}, (a_n)_{n \in \mathbb{N}}) \subset \mathbb{C}$ with radius of convergence R > 0. Define the function f on the disk $D(z_0, R)$ by:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n.$$

We must show that f is analytic on $D(z_0, R)$. Let $z_1 \in D(z_0, R)$ arbitrary. We will show that f is analytic at z_1 . For $z \in D(z_1, R - |z_1 - z_0|)$, we have

$$|z - z_0| \stackrel{T.I}{\leq} \underbrace{|z - z_1|}_{< R - |z_1 - z_0|} + |z_1 - z_0| < R$$

Thus $D(z_1, R - |z_1 - z_0|) \subset D(z_0, R)$, so the power series $\sum_{n=0}^{\infty} a_n (z - z_0)^n$ converges absolutely. so:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

$$= \sum_{n=0}^{\infty} a_n ((z - z_1) + (z_1 - z_0))^n$$

$$= \sum_{n=0}^{\infty} a_n \sum_{k=0}^n \binom{n}{k} (z - z_1)^k (z_1 - z_0)^{n-k}$$

$$= \sum_{k=0}^{\infty} \left(\sum_{n=k}^{\infty} a_k \binom{n}{k} (z_1 - z_0)^{n-k}\right) (z - z_1)^k$$

The interchange of summation is justified by the absolute convergence of the double series for $z \in D(z_1, R - |z_1 - z_0|)$. This express f(z) as a power series in $(z - z_1)$ in the disk $D(z_1, R - |z_1 - z_0|)$, proving that f is analytic at z_1 . Since z_1 was arbitrary in $D(z_0, R)$, then f is analytic on $D(z_0, R)$.

Lecture 3

08:14 AM Mon, Oct 13 2025

Example: The power series $\sum_{n=0}^{\infty} \frac{z^n}{n!}$ has radius of convergence $R = +\infty$. Therefore (by the previous Theorem), it defines an analytic function on the whole complex plane \mathbb{C} .

Definition 1.3.1: The analytic function on C defined by:

$$\exp(z) = e^z := \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

is called the exponential function.

Definition 1.3.2 (Entire function) : A complex function $f : \mathbb{C} \longrightarrow \mathbb{C}$ which is analytic on the whole complex plane \mathbb{C} is called an <u>entire function</u>.

Example:

- ① Every complex polynomial is an entire function.
- ② The exponential function $\exp(z)$ is an entire function.

1.3.1 Properties of the exponential function

Proposition 1.3.2: The exponential function defines the following properties:

① $\forall z_1, z_2 \in \mathbb{C}$, we have:

$$e^{z_1+z_2}=e^{z_1}\cdot e^{z_2}$$
 and $e^{z_1-z_2}=\frac{e^{z_1}}{e^{z_2}}$.

- ② for all $z \in \mathbb{C}$, we have $e^z \neq 0$.
- ③ (EULER'S FORMULA): $\forall \theta \in \mathbb{R}$, we have:


$$e^{i\theta} = \cos\theta + i\sin\theta.$$


 $\textcircled{4} \ \forall z \in \mathbb{C}$, we have:

$$e^z = 1 \iff z \in 2\pi i \mathbb{Z}.$$

More generally, for all $z, z' \in \mathbb{C}$, we have:

$$e^z = e^{z'} \iff z - z' \in 2\pi \mathbb{Z}.$$





1.3.2 Trigonometric and hyperbolic functions

Definition 1.3.3 (Complex Trigonometric functions) : We define the trigonometric functions cosine and sine by:

$$\cos z := \sum_{n=0}^{+\infty} (-1)^n \frac{z^{2n}}{(2n)!},$$

$$\sin z := \sum_{n=0}^{+\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!} \qquad (\forall z \in \mathbb{C}).$$

Clearly, these functions extend the real functions cos and sin. The power series defining cos and sin have infinite radius of convergence, thus (By a previous theorem) cos and sin are analytic on C; that is, cos and sin are <u>entire functions.</u>

Remark 🐿

We easily verify the extended Euler's formula:

$$e^{iz} = \cos z + i \sin z$$
 $(\forall z \in \mathbb{C}).$

From this formula, we derive:

$$\cos z = \frac{e^{iz} + e^{-iz}}{2},$$

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \qquad (\forall z \in \mathbb{C}).$$

Exercise

Using property @ of *Proposition 1.3.2* and Euler's formula, show the following properties:

- ① The functions cos and sin are both 2π -periodic.
- ② The set of zeros of $z \mapsto \cos z$ is $(\frac{\pi}{2} + \pi \mathbb{Z})$, while the set of zeros of $z \mapsto \sin z$ is $\pi \mathbb{Z}$.
- ③ For all $z \in \mathbb{C}$, we have

$$\cos^2 z + \sin^2 z = 1.$$

These functions are not bounded in C, when you replace $x \leftarrow ix$, you get $\cos ix = \cosh x$.

FOR EXAMPLE, FOR ③: By the Euler formula, we have for all $z \in \mathbb{C}$:

$$\cos^{2} z + \sin^{2} z = \left(\frac{e^{iz} + e^{-iz}}{2}\right)^{2} + \left(\frac{e^{iz} - e^{-iz}}{2i}\right)^{2}$$
$$= \frac{4}{4} = 1$$

Definition 1.3.4 (Complex hyperbolic functions): We define the hyperbolic functions cosh and sinh by:

$$\cosh z := \sum_{n=0}^{+\infty} \frac{z^{2n}}{(2n)!} = \frac{e^z + e^{-z}}{2} = \cos(iz),
\sinh z := \sum_{n=0}^{+\infty} \frac{z^{2n+1}}{(2n+1)!} = \frac{e^z - e^{-z}}{2} = -i\sin(iz) \qquad (\forall z \in \mathbb{C}).$$

Clearly, these definitions extend the real functions cosh and sinh. Like the trigonometric functions cos and sin, the hyperbolic functions cosh and sinh are also entire functions.

Exercise

Using the expressions of cosh and sinh in terms of cos and sin, verify the following properties:

- ① The functions cosh and sinh are both 2π -periodic.
- ② The set of zeros of cosh is $(\frac{\pi}{2}i + \pi i \mathbb{Z})$, while the set of zeros of sinh is $\pi i \mathbb{Z}$.

$$\cosh^2 z - \sinh^2 z = 1.$$

Definition 1.3.5 (Further trigonometric and hyperbolic functions) : We define the following functions:

$$\tan z := \frac{\sin z}{\cos z} \qquad \left(\forall z \in \mathbb{C} \setminus \left(\frac{\pi}{2} + \pi \mathbb{Z} \right) \right),$$

$$\cot z := \frac{\cos z}{\sin z} \qquad \left(\forall z \in \mathbb{C} \setminus \pi \mathbb{Z} \right),$$

$$\tanh z := \frac{\sinh z}{\cosh z} \qquad \left(\forall z \in \mathbb{C} \setminus \left(\frac{\pi}{2} i + \pi i \mathbb{Z} \right) \right),$$

$$\coth z := \frac{\cosh z}{\sinh z} \qquad \left(\forall z \in \mathbb{C} \setminus \pi i \mathbb{Z} \right).$$

This clearly extends the well-known real functions tan, cot, tanh, and coth. Note that each of these four functions is analytic in its domain of definition (according to the previous

results on analytic functions).

1.4 Holomorph functions

Definition 1.4.1: Let Ω be a nonempty open subset of \mathbb{C} and z_0 be a point in Ω . Let also $f:\Omega\longrightarrow\mathbb{C}$ be a map.

• We say that f is holomorphic at z_0 if the limit

$$\lim_{z \to z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists and belong to C. In this case, the limit is called the <u>derivative</u> of f at the point z_0 and denoted by $f'(z_0)$.

• We say that f is holomorphic on Ω if it is holomorphic at every point in Ω . In this case, the function

$$f': \Omega \longrightarrow \mathbb{C}$$
 $z \longmapsto f'(z)$

is called the derivative of f.

Proposition 1.4.1 (Holomorphy of power series) : Let $z_0 \in \mathbb{C}$, $(a_n)_{n \in \mathbb{N}_0} \subset \mathbb{C}$, and S be the power series

$$S(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n.$$

Suppose that *S* has a positive radius of convergence *R*. Then *S'* is holomorphic on $D(z_0, R)$ and we have for all $z \in D(z_0, R)$:

$$S'(z) = \sum_{n=0}^{+\infty} n a_n (z - z_0)^{n-1}$$
$$= \sum_{n=0}^{+\infty} (n+1) a_{n+1} (z - z_0)^n.$$

Proof. For simplicity, suppose without loss of generality that $z_0 = 0$. First, remark that by using the Hadamard formula, the power series

$$\sum_{n=1}^{+\infty} n a_n z^{n-1} = \sum_{n=0}^{+\infty} (n+1) a_{n+1} z^n$$

has the same radius of convergence R as S. It follows that $\sum_{n=1}^{+\infty} na_nz^{n-1}$ is absolutely convergent on D(0,R); That is, for all 0 < r < R, the series $\sum_{n=1}^{+\infty} n |a_n| r^{n-1}$ converges. Now, let $z_1 \in D(0,R)$ be arbitrary and show that S is holomorphic at z_1 . Choose $r \in \mathbb{R}$ such that $|z_1| < r < R$. For all $z \in D(0,r) \setminus \{z_1\}$, we have

$$\frac{S(z) - S(z_1)}{z - z_1} = \frac{\sum_{n=0}^{+\infty} a_n z^n - \sum_{n=0}^{+\infty} a_n z_1^n}{z - z_1}$$

$$= \sum_{n=0}^{+\infty} a_n \frac{z^n - z_1^n}{z - z_1}$$

$$= \sum_{n=1}^{+\infty} a_n \sum_{k=0}^{n-1} z^k z_1^{n-1-k}$$

$$= \sum_{n=1}^{+\infty} a_n \sum_{k=0}^{n-1} z^k z_1^{n-1-k} \qquad (*).$$

Next, we show that this last series of functions converges normally on $D(0,r)\setminus\{z_1\}$. For $z\in D(0,r)\setminus\{z_1\}$, we have:

$$\begin{vmatrix} a_n \sum_{k=0}^{n-1} z^k z_1^{n-1-k} \\ \le |a_n| \sum_{k=0}^{n-1} \underbrace{|z|^k}_{< r} \underbrace{|z_1|^{n-1-k}}_{< r}$$

$$\le |a_n| \sum_{k=0}^{n-1} r^{n-1}$$

$$= n |a_n| r^{n-1}$$
 (independent on z).

Since the series $\sum_{n=1}^{+\infty} n |a_n| r^{n-1}$ converges (as explained at the beginning of this of this proof) then the series of function $\sum_{n=1}^{+\infty} a_n \sum_{k=0}^{n-1} z^k z_1^{n-1-k}$ converges normally (no uniformally) on $D(0,r) \setminus \{z_1\}$. Therefore, we can interchange the limit as $z \to z_1$ and the summation for computing

 $\lim_{z\to z_1}\sum_{n=1}^{+\infty}\sum_{k=0}^{n-1}z^kz_1^{n-1-k}$. Doing so, we get according to (*);

$$\lim_{z \to z_1} \frac{S(z) - S(z_1)}{z - z_1} = \sum_{n=1}^{+\infty} \lim_{z \to z_1} a_n \sum_{k=0}^{n-1} z^k z_1^{n-1-k}$$

$$= \sum_{n=1}^{+\infty} a_n \sum_{k=0}^{n-1} z_1^k z_1^{n-1-k}$$

$$= \sum_{n=1}^{+\infty} n a_n z_1^{n-1} \in \mathbb{C}.$$

Hence S is holomorphic at z_1 and we have

$$S'(z_1) = \sum_{n=1}^{+\infty} n a_n z_1^{n-1}$$
$$= \sum_{n=0}^{+\infty} (n+1) a_{n+1} z_1^n.$$

Since z_1 is arbitrary in D(0, R) then S is holomorphic on D(0, R) and we have for all $z \in D(0, R)$:

$$S'(z) = \sum_{n>1} n a_n z^{n-1} = \sum_{n=0}^{+\infty} (n+1) a_{n+1} z^n.$$

Lecture 4

08:04 AM Mon, Oct 20 2025

Corollary 1.4.2 (Infinite differentiability of power series) : Let $z_0 \in \mathbb{C}$, $(a_n)_{n \in \mathbb{N}_0} \subset \mathbb{C}$, and S be the power series

$$S(z) := \sum_{n=0}^{+\infty} a_n (z - z_0)^n.$$

Suppose that *S* has a positive radius of convergence *R*. Then *S* is infinitely \mathbb{C} —differentiable on $D(z_0, R)$ and we have for all $k \in \mathbb{N}_0$ and all $z \in D(z_0, R)$:

$$S^{(k)}(z) = \sum_{n=k}^{+\infty} n(n-1)\dots(n-k+1)a_n(z-z_0)^{n-k}$$

$$= \sum_{n=0}^{+\infty} (n+k)(n+k-1)\dots(n+1)a_{n+k}(z-z_0)^n$$

$$= \sum_{n=0}^{+\infty} \frac{(n+k)!}{n!} a_{n+k}(z-z_0)^n.$$

In particular, we have for all $k \in \mathbb{N}_0$:

$$S^{(k)}(z_0) = k!a_k.$$

Corollary 1.4.3 (Analytic functions are \mathbb{C} -infinitely differentiable): Let Ω be a nonempty open subset of \mathbb{C} and $z_0 \in \Omega$. Let also $f : \Omega \longrightarrow \mathbb{C}$ be a map.

① If f is analytic at z_0 then f is infinitely \mathbb{C} -differentiable (no holomorphic) on some neighborhood of z_0 and we have in that neighborhood:

$$f(z) = \sum_{n=0}^{+\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n$$

TAYLOR'S FORMULA

② If f is analytic on Ω then f is infinitely \mathbb{C} -differentiable (so holomorphic) on Ω .

Proof. Represent f by a power series in S in a neighborhood of z_0 and apply Corollary 3.

Remark 🐿

Analytic \implies holomorphic

② CAUCHY (1825):

 f_n holomorphic + f' is continuous $\implies f$ is analytic.

3 GOURSAT (1900):

f is holomorphic $\implies f$ is analytic.

Definition 1.4.2: Let Ω be a nonempty open subset of $\mathbb C$ and $f:\Omega\longrightarrow\mathbb C$ be a map. An antiderivative of f is a holomorphic function $F:\Omega\longrightarrow\mathbb C$ such that F'=f.

Proposition 1.4.4 (Existence of Local antiderivatives): Let Ω be a nonempty open subset of $\mathbb C$ and $z_0 \in \Omega$. Let also $f: \Omega \longrightarrow \mathbb C$ be a map. If f is analytic at z_0 then f admits an antiderivative in a neighborhood of z_0 . Precisely, $\exists r > 0$ and $F: D(z_0, r) \longrightarrow \mathbb C$ analytic such that F'(z) = f(z) for all $z \in D(z_0, r)$.

Proof. Suppose that f is analytic at z_0 . then $\exists r > 0$, $\exists (a_n)_{n \in \mathbb{N}_0} \subset \mathbb{C}$ such that for all $z \in D(z_0, r)$:

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n.$$

Define $F: D(z_0, r) \longrightarrow \mathbb{C}$ by

$$F(z) = \sum_{n=0}^{+\infty} \frac{a_n}{n+1} (z-z_0)^{n+1} = \sum_{n=1}^{+\infty} \frac{a_{n-1}}{n} (z-z_0)^n \qquad (\forall z \in D(z_0, r)).$$

The Hadamard formula shows that this last power series has the name radius of convergence as the original power series $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ representing f (which is $\geq r$). Consequently, F is well-defined on $D(z_0,r)$, and by the previous results, F is even analytic on $D(z_0,r)$ so holomorphic on

 $D(z_0, r)$ and for all $z \in D(z_0, r)$:

$$F'(z) = \sum_{n=1}^{+\infty} \frac{a_{n-1}}{n} n(z - z_0)^{n-1}$$
$$= \sum_{n=1}^{+\infty} a_{n-1} (z - z_0)^{n-1}$$
$$= \sum_{n=0}^{+\infty} a_n (z - z_0)^n = f(z).$$

Thus, *F* is an antiderivative of *f* on $D(z_0, r)$, completing the proof.

Remark 🐿

The rules of differentiation for analytic/holomorphic functions are the same as those of real-valued functions. For example:

$$(fg)' = f'g + fg'$$
$$(f \circ g) = g' \cdot (f' \circ g).$$

On the other hand, the derivatives of known elementary functions, such that $z \to e^z$, $z \to \cos z$, $z \to \sin z$, etc are the same as in the real case. For example:

$$(e^z)' = e^z \qquad (\forall z \in \mathbb{C})$$

 $(\sin z)' = \cos z \qquad (\forall z \in \mathbb{C})$

Proof.

$$e^z = \sum_{n=0}^{+\infty} \frac{z^n}{n!} \qquad R = +\infty.$$

$$(e^{z})' = \sum_{n=1}^{+\infty} \frac{n}{n!} z^{n-1}$$
$$= \sum_{n=1}^{+\infty} \frac{z^{n-1}}{(n-1)!}$$
$$= \sum_{n=0}^{+\infty} \frac{z^{n}}{n!} = e^{z}.$$

1.5 The Cauchy-Riemann equations

Theorem 1.5.1 (Cauchy-Riemann equations): Let Ω be a nonempty open subset of \mathbb{C} , $z_0 = x_0 + iy_0$ with $(x_0, y_0 \in \mathbb{C})$ a point in Ω , and $f : \Omega \longrightarrow \mathbb{C}$ be a map. Let $P : \text{Re} f : \Omega \longrightarrow \mathbb{R}$ and $Q : \text{Im} f : \Omega \longrightarrow \mathbb{R}$ so that

$$f(z) = P(x,y) + iQ(x,y).$$

for all $z = x + iy \in \Omega$, with $x, y \in \mathbb{R}$ then f is holomorphic at z_0 if and only if P and Q are differentiable at (x_0, y_0) and satisfy the following Cauchy-Riemann equations at (x_0, y_0) :

$$\begin{cases} \frac{\partial P}{\partial x}(x_0, y_0) = \frac{\partial Q}{\partial y}(x_0, y_0) \\ \frac{\partial P}{\partial y}(x_0, y_0) = -\frac{\partial Q}{\partial x}(x_0, y_0) \end{cases}$$

Proof.

$$(\Longrightarrow)$$

Suppose that f in holomorphic at z_0 . Then for $h = u + iv \quad (u, v \in \mathbb{R})$, sufficiently small, we have:

$$f(z_0 + h) = f(z_0) + \cosh + o(h),$$

with $c = c_1 + ic_2 \in \mathbb{C}$ $(c_1, c_2 \in \mathbb{R})$. expanding this, we find:

$$P(x_0 + u, y_0 + v) + iQ(x_0 + u, y_0 + v) = P(x_0, y_0) + iQ(x_0, y_0) + (c_1 + ic_2)(u + iv) + o(u, v).$$

Identifying real and imaginary parts gives:

$$P(x_0 + u, y_0 + v) = P(x_0, y_0) + c_1 u - c_2 v + o(u, v),$$

$$Q(x_0 + u, y_0 + r) = Q(x_0, y_0) + c_2 u + c_1 v + o(u, v).$$

$$\frac{\partial P}{\partial x}(x_0, y_0) = c_1, \quad \frac{\partial P}{\partial y}(x_0, y_0) = -c_2, \quad \frac{\partial Q}{\partial x}(x_0, y_0) = c_2, \quad \frac{\partial Q}{\partial y}(x_0, y_0) = c_1.$$

Thus, *P* and *Q* indeed satisfying the Cauchy-Riemann condition at (x_0, y_0) .

$$(\Leftarrow)$$

Conversly, suppose that P and Q are differentiable at (x_0, y_0) and satisfy the Cauchy-Riemann conditions at this point. Set

$$c_1 := \frac{\partial P}{\partial x}(x_0, y_0) = \frac{\partial Q}{\partial y}(x_0, y_0) \in \mathbb{R}$$

$$c_2 := \frac{\partial Q}{\partial x}(x_0, y_0) = -\frac{\partial f}{\partial y}(x_0, y_0) \in \mathbb{R}$$

By hypothesis, for $(u, v) \in \mathbb{R}^2$ sufficiently small, we have:

$$P(x_0 + u, y_0 + v) = P(x_0 + y_0) + c_1 u - c_2 v + o(u, v)$$

$$Q(x_0 + u, y_0 + v) = Q(x_0, y_0) + c_2 u + c_1 v + o(u, v).$$

Then, setting h = u + iv:

$$f(z_0 + h) = P(x_0 + u, y_0 + v) + iQ(x_0 + u, y_0 + v)$$

$$= P(x_0, y_0) + iQ(x_0, y_0) + \underbrace{(c_1 + ic_2)}_{c}(u + iv) + o(u, v)$$

$$= f(z_0) + ch + o(h),$$

with $c = c_1 + ic_2$. This shows that f is holomorphic at z_0 . The theorem is proved.

Lecture 5

14:41 PM Tue, Oct 21 2025

Corollary 1.5.2 (Cauchy-Riemann equations on an open set): Let Ω be an open subset of \mathbb{C} and $f:\Omega\longrightarrow\mathbb{C}$ be a map. Let $P:=\mathrm{Re} f:\Omega\longrightarrow\mathbb{R}$ and $Q:=\mathrm{Im} f:\Omega\longrightarrow\mathbb{R}$, so that

$$f(z) = P(x, y) + iQ(x, y)$$

for all $z = x + iy \in \Omega$, with $x, y \in \mathbb{R}$. Then f is holomorphic on Ω if and only if P and Q are differentiable on Ω and satisfy the following Cauchy-Riemann equations:

$$\frac{\partial P}{\partial x} = \frac{\partial Q}{\partial y},$$
$$\frac{\partial P}{\partial y} = -\frac{\partial Q}{\partial x},$$

on Ω .

1.5.1 The isolated zeros thereom

Some toplogical remainders:

Definition 1.5.1 (Limit points) : Let E be a toplogical space, $A \subset E$, $x \in E$. we say that x is a limit point of A if every neighborhood of x intersect A in a point different from x; That is,

$$\forall V \in \mathcal{V}(x): V \cap (A \setminus \{x\}) \neq \emptyset.$$

Note that this is equivalent to $x \in \overline{A \setminus \{x\}}$. The set of all limit points of A is denoted by A' and is called the derived set of A.

→ In metric spaces, we have the following equivallent definition:

Definition 1.5.2 : Let *E* be a metric space, $A \subset E$, and $x \in E$. We say that *x* is a <u>limit point</u> of *A* if there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in $A \setminus \{x\}$ that converges to *x* .

Definition 1.5.3: Let *E* be a toplogical space and $A \subset E$.

- ① We say that an element $a \in A$ is <u>isolated</u> if its not a limit point of A.
- ② We say that *A* is a discrete set if all its points are isolated.

Example: The set \mathbb{N} and \mathbb{Z} are discete in \mathbb{R} , whereas the set \mathbb{Q} is not (even though its countable).

Proposition 1.5.3 : In $\mathbb{R}^n (n \in \mathbb{N})$, every discrete set is at most countable.

Proof. ⇔ Exercise!

Theorem 1.5.4 (The isolated zeros theorem) : Let Ω be a nonempty connected open set in \mathbb{C} and let f be a an analytic function on Ω that is not identically zero. Then the zero of f in Ω are all isolated; In other words, the set of zeros of f in Ω is discrete.

Proof. We proceed by contradiction. Suppose that there exists a zero $z_0 \in \Omega$ of f that is not isolated; i.e., z_0 is a limit point of the set of all zeros of f in Ω . Therefore, there exists a sequence $(z_k)_{k\geq 1}$ of zeros of f in Ω , with all terms distinct from z_0 , that converges to z_0 . Since f is analytic at z_0 , $\exists r>0$ and a power series representation:

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$$
 $(\forall z \in D(z_0, r))$

with $a_n \in \mathbb{C}$ for all $n \in \mathbb{N}_0$.

•• ① Let us first show that the coefficients $a_n (n \in \mathbb{N})$ must necessarily all be zero. We proceed by contradictiom, assuming the contrary, and consider

$$P := \min \left\{ n \in \mathbb{N}_0 : a_n \neq 0 \right\}.$$

We then have for all $z \in D(z_0, r)$:

$$f(z) = \sum_{n=p}^{+\infty} a_n (z - z_0)^n$$

$$= (z - z_0)^p \left[a_p + a_{p+1} (z - z_0) + a_{p+2} (z - z_0)^2 \right]$$

$$= (z - z_0)^p \sum_{n=0}^{+\infty} a_{n+p} (z - z_0)^n.$$

By specializing to $z = z_k$ (for k sufficiently large so that $z_k \in D(z_0, r)$) we find that (for $k \ge 1$ sufficiently large):

$$\underbrace{f(z_k)}_{=0} = \underbrace{(z_k - z_0)^p}_{\neq 0} \sum_{n=0}^{+\infty} a_{n+p} (z_k - z_0)^n.$$

Hence

$$\sum_{n=0}^{+\infty} a_{n+p} (z_k - z_0)^n = 0,$$

taking the limit as $k \to +\infty$ and noting the normal convergence of the series on the left, we obtain: $a_p = 0$. This contradicts the definition of p and shows that $a_n = 0$ for all $n \in \mathbb{N}_0$. In follows from this that:

$$f(z) = 0$$
 $(\forall z \in D(z_0, r)).$

Now, we will show that f is identically zero on all Ω , which will yield the desired contradiction. Let $\omega \in \Omega$ be arbitrary and show that $f(\omega) = 0$. Since Ω is connected (no path-connected, as its an open subset of \mathbb{C}), there exists a continuous path $\gamma : [0,1] \longrightarrow \Omega$ such that $\gamma(0) = z_0$ and $\gamma(1) = \omega$. Consider

$$t_0 := \sup \{ t \in [0,1] : (f \circ \gamma)(t) = 0 \}$$

The supremum exists because the set is non empty, as it contains 0, and it is bounded from above by 1. Since f and γ are continuous, the function $f \circ \gamma$ are continuous on [0,1]. Consequently, the set

$$\{t \in [0,1] : (f \circ \gamma)(t) = 0\} = (f \circ \gamma)^{-1}(\{0\})$$

which is closed in [0,1]. Thereforem t_0 belongs to this set; in other words, we have

$$(f \circ \gamma)(t_0) = 0 \tag{1}$$

Let us show that $t_0 = 1$. Suppose for contradiction, that $t_0 < 1$. By the reasoning from the previous part of this proof (replacing z_0 by $\gamma(t_0)$, which is a limit point of the zero of f), there exists r' > 0 such that

$$f(z) = 0$$
 $(\forall z \in D(\gamma(t_0), r')).$

FOr $\varepsilon > 0$, sufficiently small, we have:

$$\gamma(t_0 + \varepsilon) \in D(\gamma(t_0), r'),$$

(by the continuity of γ). Therefore:

$$g(\gamma(t_0 + \varepsilon)) = 0,$$

i.e. $(f \circ \gamma)(t_0 + \varepsilon) = 0$. This contradicts the very definition of t_0 as the supremum. Hence, necessarily $t_0 = 1$. This gives, from (1), $f(\omega) = 0$. Since ω was arbitrary in Ω , we have $f \equiv 0$ on Ω . Contradiction. This final contradiction ensures that the zeros of f in Ω are all isolated. The theorem is proved. there is a tiny error in this proof, will be fixed next time.

Corollary 1.5.5 (Principle of analytic continuation): Let f and g be two analytic functions on a nonempty <u>connected</u> open subset Ω of $\mathbb C$ that coincide on a subset $A \subset \Omega$ pocessing a limit point in Ω . Then f and g are identical on Ω .

Proof. Let $\varphi := f - g$. Then φ is analytic on Ω and vanishes on the set $A \subset \Omega$, which has a limit point $a \in \Omega$. So $a \in \overline{A \setminus \{a\}} \subset \overline{A}$. Since φ vanishes on A and is continuous on Ω , then it vanishes on $\overline{A} \cap \Omega$. In particular, φ vanishes at a. Therefore, a is a non-isolated zero of φ . By the isolated zero theorem, this implies that $\varphi \equiv 0$ on Ω ; That is, $f \equiv g$ on Ω .

Example: Let us show that (without using the extended Euler formulas) that for all $z \in \mathbb{C}$, we have:

$$\cos^2 z + \sin^2 z = 1.$$

consider $f(z) := \cos^2 z + \sin^2 z$ and g(z) := 1. f and g are analytic on \mathbb{C} (which is an connected open subset of \mathbb{C}) and coincide on \mathbb{R} , which posesses a limit point in \mathbb{C} . Thus,

by the principle of analytic continuation $f \equiv g$ on \mathbb{C} ; i.e.

$$\cos^2 z + \sin^2 z = 1 \qquad (\forall z \in \mathbb{C}).$$

1.5.2 Multiplicity of a zero of an analytic function

Theorem 1.5.6: Let Ω be a nonempty connected open subset of \mathbb{C} , and let f be an analytic function on Ω , not identically zero. Let $z_0 \in \Omega$ be a zero of f. Then there exists a unique positive integer p and a unique analytic function g on Ω does not vanish at z_0 , such that

$$f(z) = (z - z_0)^p g(z) \qquad (\forall z \in \Omega)$$

Proof. ightharpoonup Existence of p and g:

Since f is analytic at z_0 , $\exists r > 0$ and a complex sequence $(a_n)_{n \in \mathbb{N}_0}$ such that $D(z_0, r) \subset \Omega$ and

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n \qquad (\forall z \in D(z_0, r)).$$

Since f is not identically zero on Ω , it is certainly not identically zero on $D(z_0, r)$ (By the isolated zeros theorem). Thus the coefficients a_n are not all zero. We can therefore define

$$p:=\min\left\{n\in\mathbb{N}_0:a_n\neq 0\right\}.$$

since $a_0 = f(z_0)$, we have $p \ge 1$. Then, for all $z \in D(z_0, r)$:

$$f(z) = \sum_{n=p}^{+\infty} a_n (z - z_0)^n$$
$$= (z - z_0)^p \sum_{n=0}^{+\infty} a_{n+p} (z - z_0)^n$$

Now, define $g: \Omega \longrightarrow \mathbb{C}$ by:

$$g(z) := \begin{cases} \frac{f(z)}{(z-z_0)^p} & \text{if } z \neq z_0, \\ a_p & \text{if } z = z_0. \end{cases}$$

We observe that:

- g is analytic on $\Omega \setminus \{z_0\}$ (as a quotitent of two analytic functions on $\Omega \setminus \{z_0\}$).
- For $z \in D(z_0, r)$, $g(z) = \sum_{n=0}^{+\infty} a_{n+p} (z z_0)^n$ which shows that g is analytic at z_0 . Hence, g is analytic on Ω . Moreover, we have

$$f(z) = (z - z_0)^p g(z) \qquad (\forall z \in \Omega)$$

and
$$g(z_0) = a_p \neq 0$$
.

$\bullet \bullet$ Uniqueness of p and g:

Suppose there exists $p_1, p_2 \in \mathbb{N}$ and analytic functions g_1, g_2 on Ω that do not vanish at z_0 , such that

$$f(z) = (z - z_0)^{p_1} g_1(z) = (z - z_0)^{p_2} g_2(z) \qquad (\forall z \in \Omega).$$

Then, for all $z \in \Omega \setminus \{z_0\}$, we have:

$$g_1(z) = (z - z_0)^{p_2 - p_1} g_2(z).$$

If $p_1 < p_2$ then taking the limit as $z \to z_0$ and using the continuity of g_1 and g_2 at z_0 , we obtain $g_1(z_0) = 0$, which contradicts the hypothesis $g_1(z_0) \neq 0$. Therefore, we must have $p_1 \geq p_2$. By symmetry, we also have $p_2 \geq p_1$, so $p_1 = p_2$. Then, from above, we get

$$\forall z \in \Omega \setminus \{z_0\} : g_1(z) = (z - z_0)^{p_2 - p_1} g_2(z),$$

since $p_2 - p_1 = 0$, we get

$$g_1(z) = g_2(z).$$

Since g_1 and g_2 are continuous at z_0 , taking the limit as $z \to z_0$ gives $g_1(z_0) = g_2(z_0)$. Hence $g_1 \equiv g_2$ on Ω , which completes the proof of uniqueness.

Definition 1.5.4: In the context of the above theorem, the positive integer p is called the multiplicity of the zero z_0 of f. if z_1 is not a zero of f, its multiplicity is conventionally taken to be 0.

Chapter 2

The Cauchy integral formula on a circle and applications

Lecture 6

08:09 AM Mon, Oct 27 2025

Theorem 2.0.1 : Let $z_0 \in \mathbb{C}$ and $f : \mathbb{C} \longrightarrow \mathbb{C}$ be a function, analytic at z_0 . Let also R be the radius of convergence of the power series representing f in a neighborhood of z_0 . Then for every $n \in \mathbb{N}_0$ and every 0 < r < R, we have:

$$f^{(n)}(z_0) = \frac{n!}{2\pi r^n} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-ni\theta} d\theta$$

In particular, for n = 0 and 0 < r < R.

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) d\theta$$

Proof. By hypothesis, \exists ! complex sequence $(a_n)_{n \in \mathbb{N}_0}$ such that for all $z \in D(z_0, R)$

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n \qquad (\forall z \in D(z_0, R))$$

Fix $r \in (0, R)$. For every $\theta \in [0, 2\pi]$, taking $z = z_0 + re^{i\theta} \in C(z_0, r) \subset D(z_0, R)$ in the formula

above, we obtain

$$f(z_0 + re^{i\theta}) = \sum_{n=0}^{+\infty} a_n r^n e^{in\theta}.$$

Since the trigonometric series on the right converges normally (so uniformally) with respect to $\theta \in [0,2\pi]$ (Because for all $n \in \mathbb{N}_0$, we have $|a_n r^n e^{ni\theta}| = |a_n r^n|$, and the series $\sum_{n=0}^{+\infty} |a_n| r^n$ converges by properties of power series) then that is the Fourier series of the function $\theta \mapsto f(z_0 + re^{i\theta})$. Consequently, for every $n \in \mathbb{N}_0$, the Fourier coefficients are given by:

$$a_n r^n = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-ni\theta} d\theta.$$

Thus

$$a_n = \frac{1}{2\pi r^n} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-ni\theta} d\theta.$$

Comparing this with Taylor's formula

$$a_n = \frac{f^{(n)}(z_0)}{n!}$$

, we obtain for every $n \in \mathbb{N}_0$:

$$f^{(n)}(z_0) = \frac{n!}{2\pi r^n} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-in\theta} d\theta,$$

as required.

Remark ®

In the context of *Theorem* 1, the right-hand side of the second formula (corresponding to n = 0):

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) d\theta.$$

is precisely the average value of f on the circle $C(z_0, r)$. Therefore, according to *Theorem 1*, the average value of f on any circle centered at z_0 and contained in $D(z_0, R)$ is equal to the value of f at z_0 .

2.0.1 Analytic continuation of power series

Theorem 2.0.2: Let Ω be a nonempty open subset of \mathbb{C} , and let $f:\Omega \longrightarrow \mathbb{C}$ be an analytic function on Ω . Let $z_0 \in \Omega$, and let $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ (where $a_n \in \mathbb{C}$, for all $n \in \mathbb{N}_0$) be the unique power series representing f in a small neighborhood $D(z_0,r) \subset \Omega$ of z_0 with (with r > 0). Then, the power series $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ converges on every disk centered at z_0 and contained in Ω , and it sum remains f(z).

Proof. We have

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n \qquad (\forall z \in D(z_0, r)).$$

Let R > r such that $D(z_0, R) \subset \Omega$ and show that the above formula remains valid on $D(z_0, R)$. Given $n \in \mathbb{N}_0$, by Cauchy's integral formula on a circle, we have for all $0 < \sigma < r$:

$$a_n = \frac{1}{2\pi\sigma^n} \int_0^{2\pi} f(z_0 + \sigma e^{i\theta}) e^{-ni\theta} d\theta.$$

hence $\sigma \mapsto \frac{1}{2\pi\sigma^n} \int_0^{2\pi} f(z_0 + \sigma e^{i\theta}) e^{-ni\theta} d\theta$ is constant (with respect to σ) in (0, r). This suggests us to define

$$\varphi: (0,R) \longrightarrow \mathbb{C}$$

$$\sigma \longmapsto \varphi(\sigma) = \frac{1}{2\pi\sigma^n} \int_0^{2\pi} f(z_0 + \sigma e^{i\theta}) e^{-ni\theta} d\theta.$$

 $(\varphi \text{ is constant in } (0,r).$ with value a_n). Since f is analytic (so continuous) on $D(z_0,R)\subset\Omega$ then φ is well-defined and differentiable on (0,R). We now show that φ remains constant on (0,R) on (0,R). By applying results from measure theory and integration (which allow differentiation under the integral sign), we have for all $\sigma\in(0,R)$:

$$\varphi'(\sigma) = -\frac{n\sigma^{-n-1}}{2\pi} \int_{0}^{2\pi} f(z_{0} + \sigma e^{i\theta}) e^{-ni\theta} d\theta + \frac{\sigma^{-n}}{2\pi} \int_{0}^{2\pi} f'(z_{0} + \sigma e^{i\theta}) e^{(-n+1)i\theta} d\theta.$$

$$= \frac{\sigma^{-n-1}}{2\pi} \int_{0}^{2\pi} \underbrace{\left[-nf(z_{0} + \sigma e^{i\theta}) e^{-ni\theta} + \sigma f'(z_{0} + \sigma e^{i\theta}) e^{(-n+1)i\theta} \right]}_{\frac{d}{d\theta}(-if(z_{0} + \sigma e^{i\theta}) e^{-ni\theta})} d\theta$$

$$= \frac{\sigma^{-n-1}}{2\pi} \underbrace{\left[-if(z_{0} + \sigma e^{i\theta}) e^{-ni\theta} \right]_{0}^{2\pi}}_{0} = 0.$$

This shows that φ is constant on (0, R), and its value is a_n (since $\varphi \equiv a_n$ on (0, r)). Thus, we have shown that:

$$a_n = \frac{1}{2\pi\sigma^n} \int_0^{2\pi} f(z_0 + \sigma e^{i\theta}) e^{-ni\theta} d\theta \qquad (\forall n \in \mathbb{N}_0, \forall \sigma \in (0, R))$$

This formula allows us to estimate $|a_n|$ in terms of n. Giving $n \in \mathbb{N}_0$, $\sigma \in (0, R)$, and setting

$$M(f) := \sup_{z \in C(z_0, \sigma)} |f(z)|$$
, we have:

$$a_n = \left| \frac{1}{2\pi\sigma^n} \int_0^{2\pi} f(z_0 + \sigma e^{i\theta}) e^{-ni\theta} d\theta \right|$$

$$\leq \frac{1}{2\pi\sigma^n} \int_0^{2\pi} \left| \underbrace{f(z_0 + \sigma e^{i\theta})}_{\leq M(f)} \right| d\theta$$

$$|a_n| \le \frac{M(f)}{\sigma^n} \quad \forall n \in \mathbb{N}_0, \forall \sigma \in (0, R)$$

So $\forall z \in D(z_0, \sigma)$:

$$|a_n(z-z_0)^n| \le M(f) \left| \frac{z-z_0}{\sigma} \right|^n$$

implying that the power series $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ converges absolutely on $D(z_0,\sigma)$, $\forall \sigma \in (0,R)$. Consequently, $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ converges (absolutely) on $D(z_0,R)$. Further, define

$$g(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n \quad (\forall z \in D(z_0, R)).$$

Since g is analytic on $D(z_0, R)$ and coincides with f on $D(z_0, r)$, by the principle of analytic continuation, g coincides with f on $D(z_0, R)$. Hence

$$f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$$
 $(\forall z \in D(z_0, R)),$

completing the proof.

Corollary 2.0.3 : Let Ω be a nonempty open subset of \mathbb{C} and z_0 be a point in Ω . Let also $f:\Omega \longrightarrow \mathbb{C}$ be an analytic function on Ω and $\sum_{n=0}^{+\infty} a_n(z-z_0)^n$ (where $(a_n)_{n\in\mathbb{N}_0}\subset\mathbb{C}$) be a power series representing f in a neighborhood of z_0 . Then the radius of convergence R of that power series satisfies;

$$R \geq \operatorname{dist}(z_0, \mathbb{C} \setminus \Omega).$$

Proof. Set $\sigma := \operatorname{dist}(z_0, \mathbb{C} \setminus \Omega)$. By definition of S, we have $D(z_0, \sigma) \subset \Omega$. (Indeed, if $z \in \mathbb{C} \setminus \Omega$ then $d(z, z_0) \geq \inf_{w \in \mathbb{C} \setminus \Omega} d(w, z_0) = d(z_0, \mathbb{C} \setminus \Omega) = \sigma$, thus $z \in \mathbb{C} \setminus D(z_0, \sigma)$.) Thus, by *Theorem 2*, the power series $\sum_{n=0}^{+\infty} a_n (z-z_0)^n$ converges on $D(z_0, \sigma)$. Hence $R \geq \sigma$, as required.

Example: ♥ VERY IMPORTANT.

Consider the function $f(z) = \frac{z}{e^z - 1}$ with the convention f(0) = 1. For all z in a small

neighborhood of θ , with $z \neq 0$, we have:

$$f(z) = \frac{z}{z + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots}$$
$$= \frac{1}{1 + \frac{z}{2!} + \frac{z^2}{3!} + \dots},$$

Which also holds for z=0, showing that f is analytic at 0 (as a quotient of two analytic functions at 0, with the denominator nonzero at 0). Moreover, the zero of $z\mapsto e^z-1$ are the complex numbers $2k\pi i$ with $k\in\mathbb{Z}$, implying that f is analytic on $\mathbb{C}\setminus 2\pi i\mathbb{Z}$ (as a quotient of two analytic functions on this domain, with the denominator nonzero at any point of the domain). In conclusion, f is analytic on the domain

$$\Omega = (\mathbb{C} \setminus 2\pi \mathbb{Z}) \cup \{0\}$$
$$= \mathbb{C} \setminus 2\pi i \mathbb{Z}^*.$$

Consider the power series expansion of f at 0:

$$\frac{z}{e^z - 1} = \sum_{n=0}^{+\infty} B_n \frac{z^n}{n!}, \quad (*)$$

where $B_n(n \in \mathbb{N}_0)$ are the famous Bernoulli numbers, which appear in nearly all branches of mathematics (number theory, analysis, combinatorics, algebraic geometry, etc...).:w

$$B_0 = 1, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}, B_3 = 0, B_4 = \frac{1}{42}, \dots$$

 $(B+1)^n = B^n \quad (\forall n > \mathbb{N}_2).$

for n = 2:

$$B^2 + 2B + 1 = B_2 \implies B_1 = -\frac{1}{2}$$

for n = 3:

$$(B+1)^3 = B^3 \implies 3B_2 = \frac{1}{2} \implies B_2 = \frac{1}{6}.$$

Theorem 2 shows that formula (*) is valid on the largest disk centered at 0 and contained in Ω , which is $D(0,2\pi)$. Hence

$$\frac{z}{e^z - 1} = \sum_{n=0}^{+\infty} B_n \frac{z^n}{n!} \qquad (\forall z \in \mathbb{C}, |z| < 2\pi).$$

Remark 🐿 It can be shown that $B_{2n+1}=0 \quad \forall n\geq 1,$ and $B_{2n} \sim_{+\infty} (-1)^{n-1} \cdot \frac{2(2n)!}{(2\pi)^{2n}}.$ This allows us to directly verify (using Hadamard formula) that the radius of convergence of the power series $\sum_{n=0}^{+\infty} B_n \frac{z^n}{n!}$ is exactly 2π .

CHAPTER 2. THE CAUCHY INTEGRAL FORMULA ON A CIRCLE AND APPLICATIONS 35