

# Complex Analysis Lecture Notes

*Hand written summary from lectures*

---

## Acknowledgment

Special thanks to my professor **MR.BAKIR FARHI**, who gave the lectures and explanations, this work wouldn't exist without his teaching, here is the link to his website:

<http://farhi.bakir.free.fr/home/index-fr.html>

---

## Disclaimer

These notes were written in real-time during the lectures, this is not the final version, yet. so they may contain :

- Incomplete or incorrect information
- Typos, transcription mistakes, or missing content
- Interpretations or notations that reflect my own understanding at the moment

*Please double check anything important with official material or trusted sources.*

if you spot an error feel free to open an issue or submit a pull request, or contact me via gmail :

kara.abderahmane@nhsm.edu.dz

**Notes on Contribution :**

This document is a collaborative effort. students who contribute by reporting errors or helping to complete the content will be credited in the next page as contributors in future versions, your help is appreciated and helps improve this document for everyone.

## My Github Page

Last Update : 2025-09-28

### CONTRIBUTORS

*Main Writer*

KARA ABDERAHMANE

Main Drawer

Haddar Noureddine

ARMWRESTLING4EVER

# Contents

|          |  |          |
|----------|--|----------|
| <b>1</b> | <b>Power Series</b>  | <b>2</b> |
| 1.1      | Formulas for Calculating the Radius of Convergence . . . . . | 5        |
| 1.2      | Analytic Functions . . . . .                                 | 8        |
| 1.2.1    | Properties of Analytic Functions . . . . .                   | 9        |
| 1.3      | Power series define Analytic functions . . . . .             | 11       |

# Chapter 1

## Power Series

### Lecture 1

08:06 AM Mon, Sep 29 2025

**Definition 1.0.1 (Power Series)** : A power series is a formal series of the form  $\sum_{n=1}^{\infty} a_n z^n$ , where  $a_n \in \mathbb{C}$  for all  $n \in \mathbb{N}_0$ .

More generally, given  $z_0 \in \mathbb{C}$ , a power series centered at  $z_0$  is a formal series of the form:

$$\sum_{n=1}^{\infty} a_n (z - z_0)^n$$

where  $a_n \in \mathbb{C} \quad (\forall n \in \mathbb{N})$

#### Remark:

The set of all complex power series (centered at 0) is denoted by  $\mathbb{C}[[z]]$ . More generally, given  $z_0 \in \mathbb{C}$ , the set of all complex power series at  $z_0$  is denoted by  $\mathbb{C}[[z - z_0]]$

#### Operations on Formal Power Series:

Given  $z_0 \in \mathbb{C}$ , we equip  $\mathbb{C}[[z - z_0]]$ , with the following operations:

1. **Additions:** For all  $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \in \mathbb{C}$ :

$$\sum_{n=1}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} b_n (z - z_0)^n = \sum_{n=1}^{\infty} (a_n + b_n) (z - z_0)^n$$

2. **Multiplication:**

$$\sum_{n=1}^{\infty} a_n (z - z_0)^n \times \sum_{n=1}^{\infty} b_n (z - z_0)^n = \sum_{n=1}^{\infty} c_n (z - z_0)^n$$

where,  $c_n = \sum_{k=1}^n a_k b_{n-k}$  for all  $n \in \mathbb{N}_0$ . Also  $(c_n)_{n \in \mathbb{N}}$  is called the convolution of the two sequences  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$ .

3. **Scalar Multiplication:** For all  $\lambda \in \mathbb{C}$ , and all  $(a_n)_{n \in \mathbb{N}} \subset \mathbb{C}$ :

$$\lambda \sum_{n=1}^{\infty} a_n (z - z_0)^n = \sum_{n=1}^{\infty} (\lambda a_n) (z - z_0)^n$$

It's straight forward to verify that  $\mathbb{C}[[z - z_0]]$  equipped with these operations forms a commutative algebra over  $\mathbb{C}$ . The Multiplicative identity is the constant power series:

$$1 = 1 + 0 \cdot (z - z_0) + 0 \cdot (z - z_0)^2 + \dots$$

**Definition 1.0.2 (Domain of Convergence) :** The domain of convergence of a power series  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$  is the set of all points  $z \in \mathbb{C}$  for which the series converge the structure of this domain is very specific. Its a disk (possibly with some points in its boundary) centered at  $z_0$ .

**Proposition 1.0.1 (Abel's Lemma) :** Let  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$  be a power series and let  $z_1 \in \mathbb{C} \setminus \{z_0\}$ . Suppose that the sequence  $\{a_n (z_1 - z_0)^n\}_{n \in \mathbb{N}}$  is bounded. Then, the power series in question converges absolutely (so converges) for every  $z \in \mathbb{C}$ , such that:

$$|z - z_0| < |z_1 - z_0|$$

*Proof.* By hypothesis,  $\exists M > 0$  such that  $\forall n \in \mathbb{N}_0$ :

$$|a_n (z_1 - z_0)^n| \leq M$$

Then, for all  $z \in \mathbb{C}$  such that  $|z - z_0| < |z_1 - z_0|$  we have:

$$\begin{aligned} |a_n (z - z_0)^n| &= \underbrace{|a_n (z_1 - z_0)^n|}_{\leq M} \cdot \underbrace{\left| \frac{z - z_0}{z_1 - z_0} \right|^n}_{< 1} \\ &\leq M \underbrace{\left| \frac{z - z_0}{z_1 - z_0} \right|^n}_{< 1} \end{aligned}$$


since  $\left| \frac{z - z_0}{z_1 - z_0} \right| < 1$  then the geometric series

$$\sum_{n=1}^{\infty} M \left| \frac{z - z_0}{z_1 - z_0} \right|^n \text{ Converges}$$

Thus, the series  $\sum_{n=1}^{\infty} |a_n (z - z_0)^n|$  also converges, that is  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$  is absolutely convergent.  $\square$

**Corollary 1.0.2 :** Let  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  be a power series which converges at some  $z = z_1 \in \mathbb{C} \setminus \{z_0\}$ , then the power series in question converges absolutely (so converges), for every  $z \in \mathbb{C}$  such that:

$$|z - z_0| < |z - z_1|$$



*Proof.*  $\sum_{n=1}^{\infty} a_n(z_1 - z_0)^n$  converges implies that  $a_n(z - z_0)^n \rightarrow 0$  as  $n \rightarrow \infty$ , which implies that the sequence  $\{a_n(z_1 - z_0)^n\}_{n \geq 0}$  is bounded. Lemma 1 ( proposition 1 ) permits us to conclude the required result.  $\square$

**Theorem 1.0.3 (Radius of Convergence) :** Let  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  be a power series then there exists a unique  $R \in [0, \infty]$ , called the radius of convergence with the following properties:

1. The power series converges absolutely for every  $z \in \mathbb{C}$  with  $|z - z_0| < R$ .
2. The power series diverges for every  $z \in \mathbb{C}$  with  $|z - z_0| > R$ , the disk  $D(z_0, R) = \{z \in \mathbb{C} : |z - z_0| < R\}$  is called the disk of convergence.

*Proof.* Define the set  $A \subset \mathbb{R}_{\geq 0}$  of nonnegative real numbers for which the sequence  $\{|a_n| r^n\}_{n \in \mathbb{N}_0}$  is bounded.

$$A := \left\{ r \geq 0 : \sup_{n \in \mathbb{N}} |a_n| r^n < \infty \right\}$$

we have  $A \neq \emptyset$  because  $0 \in A$ .

Define  $R := \sup A \in [0, \infty]$ , we now show that  $R$  has the stated properties.

1. Let  $z \in D(z_0, R)$ . By definition of the supremum, there exists  $r \in A$ , (i.e.,  $|a_n| r^n$  is bounded) such that  $|z - z_0| < r \leq R$ , since  $|z - z_0| < r$  and  $\{|a_n| r^n\}_{n \geq 0}$  is bounded. then by Abel's lemma, we deduce that the series  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  converges absolutely.
2. Let  $z \in \mathbb{C}$  such that  $|z - z_0| > R$ , suppose for contradictions that the power series converges at  $z$ . Then by the corollary 2, it would converge absolutely for any  $\omega$  with  $|\omega - z_0| < |z - z_0|$ , In particular, for any  $r$  such that:

$$R < r < |z - z_0|$$

The series would converge at points on the circle  $C(z_0, r)$  implying  $r \in A$ . This contradicts the fact that  $R = \sup A$ , Therefore. the power series diverges.

*The Uniqueness of  $R$ :*

If another  $R' \in [0, \infty]$  satisfies the same properties, a point  $z$  such that  $|z - z_0|$  lies between  $R$  and  $R'$  would lead to a contradiction regarding the convergence or divergence of the power series.  $\square$

## 1.1 Formulas for Calculating the Radius of Convergence

**Proposition 1.1.1 (Hadamard's Formula) :** Let  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  be a power series centered at  $z_0 \in \mathbb{C}$ . Denote by  $R$  its radius of convergence. Then:

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$$

with the convention  $\frac{1}{0} = \infty$  and  $\frac{1}{\infty} = 0$

*Proof.* Let  $L := \lim_{n \rightarrow \infty} \sup |a_n|^{\frac{1}{n}} \in [0, \infty]$ , we must show that  $R = \frac{1}{L}$ . Let  $z \in \mathbb{C} \setminus \{z_0\}$ , we distinguish three cases:

1. if  $(L = 0)$ . In this case, we have:

$$0 \leq \liminf_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} \leq \limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}} = 0$$

Thus,  $\lim_{n \rightarrow \infty} \inf |a_n|^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \sup |a_n|^{\frac{1}{n}} = 0$ . This implies that  $\lim_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}$  exists and equals to 0, so for all  $n$  sufficiently large, we have:

$$|a_n|^{\frac{1}{n}} < \frac{1}{z |z - z_0|}$$

That is

$$|a_n(z - z_0)^n| < \frac{1}{z^n}$$

for all  $n$  sufficiently large, since the geometric series  $\sum_{n=1}^{\infty} \frac{1}{z^n}$  converges then the series  $\sum_{n=1}^{\infty} |a_n(z - z_0)^n|$  converges  $\forall z \in \mathbb{C}$ , thus  $R = +\infty = \frac{1}{L}$

2.  $(L = +\infty)$ , we have  $L = \lim_{n \rightarrow \infty} \sup |a_n|^{\frac{1}{n}} = +\infty$  is equivalent to the fact the sequence  $\left\{ |a_n|^{\frac{1}{n}} \right\}_{n \in \mathbb{N}}$  is bounded. Therefore, the sequence:

$$|a_n(z - z_0)^n|^{\frac{1}{n}} = |a_n|^{\frac{1}{n}} |z - z_0|$$

is also unbounded. This implies that  $|a_n(z - z_0)^n|$  is unbounded, thus  $|a_n(z - z_0)^n|$  does not converge to 0 as  $n \rightarrow \infty$ . Hence  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  diverges. Hence  $R = 0$



3. ( $L \in (0, \infty)$ ), Let  $z \in \mathbb{C}$ . We consider two subcases:

(a) if  $|z - z_0| < r < \frac{1}{L}$ , choose  $r$  such that  $|z - z_0| < r < \frac{1}{L}$ , thus  $L < \frac{1}{r}$ . By definition of a  $\lim_{n \rightarrow \infty} \sup$ , for all  $n$  sufficiently large we have:

$$|a_n|^{\frac{1}{n}} < \frac{1}{r}$$

which implies that:

$$|a_n(z - z_0)^n| < \underbrace{\left(\frac{|z - z_0|}{r}\right)^n}_{<1}$$

since  $\left|\frac{z - z_0}{r}\right| < 1$ , the geometric series  $\sum_{n=1}^{\infty} \left|\frac{z - z_0}{r}\right|^n$  converges, by comparison, the power series  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  converges absolutely

(b) if ( $|z - z_0| > \frac{1}{L}$ ), in this case, we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup |a_n(z - z_0)^n|^{\frac{1}{n}} &= \lim_{n \rightarrow \infty} \sup \left( |a_n|^{\frac{1}{n}} |z - z_0| \right) \\ &= L |z - z_0| > 1 \end{aligned}$$

Thus,  $\{a_n(z - z_0)^n\}_{n \in \mathbb{N}}$  is unbounded, hence  $|a_n(z - z_0)^n|$  does not converge to zero as  $n \rightarrow \infty$ , implying that  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  diverges therefore:

$$R = \frac{1}{L}$$

□

## Lecture 2

08:00 AM Mon, Oct 06 2025

**Proposition 1.1.2 (Ratio Test Formula) :** Let  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  be a power series. suppose that the limit

$$\alpha = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists (i.e.,  $\in [0, \infty]$ ). Then the radius of convergence  $R$  of the power series in question is  $R = \alpha$ .

*Proof.* We use the d'Allembert rule for the series

$$\sum_{n=1}^{\infty} a_n(z - z_0)^n \quad (z \in \mathbb{C} \setminus \{z_0\})$$

let  $z \in \mathbb{C} \setminus \{z_0\}$ . we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}(z - z_0)^{n+1}}{a_n(z - z_0)^n} \right| &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \cdot |z - z_0| \\ &= |z - z_0| \cdot \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \\ &= \frac{|z - z_0|}{\alpha} \end{aligned}$$

By the d'Allembert rule, we have:

- The series  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  converges if

$$\frac{|z - z_0|}{\alpha} < 1 \quad \text{i.e.} \quad |z - z_0| < \alpha$$

- The series  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  diverges if

$$\frac{|z - z_0|}{\alpha} > 1 \quad \text{i.e.} \quad |z - z_0| > \alpha$$

Hence  $R = \alpha$ .

□

**Example:** Determine the radius of convergence of the power series  $\sum_{n=1}^{\infty} \frac{z^n}{n!}$  where  $z_0 = 0$ .

1<sup>st</sup> method: (By Hadamard formula)

We must compute  $\lim_{n \rightarrow \infty} \sup \left( \frac{1}{n!} \right)^{\frac{1}{n}}$ , by the stirling formula, we have that:

$$n! \stackrel{+\infty}{\sim} n^n e^{-n} \sqrt{2\pi n}$$

Thus we get:

$$(n!)^{\frac{1}{n}} \stackrel{+\infty}{\sim} ne^{-1} (2\pi n)^{\frac{1}{2n}}$$

Thus

$$\left( \frac{1}{n!} \right)^{\frac{1}{n}} \stackrel{+\infty}{\sim} \frac{e}{n} (2\pi n)^{-\frac{1}{2n}} \rightarrow 0 \text{ as } n \rightarrow \infty$$

Thus  $R = \frac{1}{0} = +\infty$ .

This means that the power series  $\sum_{n=1}^{\infty} \frac{z^n}{n!}$  converges for all  $z \in \mathbb{C}$ .

2<sup>nd</sup> method:

We use proposition 2. we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{\frac{1}{n!}}{\frac{1}{(n+1)!}} \right| &= \lim_{n \rightarrow \infty} \frac{(n+1)!}{n!} \\ &= \lim_{n \rightarrow \infty} (n+1) = +\infty \end{aligned}$$

Thus  $R = +\infty$

## 1.2 Analytic Functions


**Definition 1.2.1 :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and let  $z_0 \in \Omega$ .

Let  $f : \Omega \rightarrow \mathbb{C}$  be a map. then:

1.  $f$  is said to be analytic at  $z_0$  if there exists  $r > 0$  and a complex sequence  $(a_n)_{n \in \mathbb{N}_0}$  such that  $D(z_0, r) \subset \Omega$  and:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n \quad * \forall z \in D(z_0, r)$$

2.  $f$  is said to be analytic on  $\Omega$  if its analytic at every point of  $\Omega$



**Example:**

1. Every complex polynomial is analytic on  $\mathbb{C}$ . Indeed, let  $P \in \mathbb{C}[\mathbb{Z}]$ , and  $z_0 \in \mathbb{C}$ . since  $P(z + z_0) \in \mathbb{C}[\mathbb{Z}]$ , we can write:

$$P(z + z_0) = \sum_{n=0}^d a_n z^n \quad (d \in \mathbb{N}_0)$$

substituting  $z$  by  $(z - z_0)$ . we get:

$$P(z) = \sum_{n=0}^d a_n (z - z_0)^n$$

which is a power series centered at  $z_0$  with infinite radius of convergence. Thus,  $P$  is analytic at  $z_0$ , since  $z_0$  was arbitrary  $P$  is analytic on  $\mathbb{C}$ .

2. The function  $z \rightarrow \frac{1}{z}$  is analytic on  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ . Indeed, let  $z_0 \in \mathbb{C}^*$  arbitrary.

For  $z \in D(z_0, |z_0|)$ , we have:

$$\left| \frac{z - z_0}{z_0} \right| < 1$$

we can write

$$\begin{aligned} \frac{1}{z} &= \frac{1}{z_0 + (z - z_0)} \\ &= \frac{1}{z_0} \cdot \frac{1}{1 + \frac{z - z_0}{z_0}} \\ &= \frac{1}{z_0} \cdot \sum_{n=1}^{\infty} (-1)^n \left( \frac{z - z_0}{z_0} \right)^n \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n}{z_0^{n+1}} (z - z_0)^n \end{aligned}$$

which is a power series centered at  $z_0$ , valid on  $D(z_0, |z_0|)$ . Hence  $z \rightarrow \frac{1}{z}$  is analytic at  $z_0$ . since  $z_0 \in \mathbb{C}^*$  was arbitrary, then  $z \rightarrow \frac{1}{z}$  is analytic on  $\mathbb{C}^*$

### 1.2.1 Properties of Analytic Functions

**Proposition 1.2.1 :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and let  $z_0 \in \Omega$  if,  $f, g : \Omega \rightarrow \mathbb{C}$  are analytic at  $z_0$ , then the same for  $(f + g)$  and  $(f \cdot g)$ . Moreover, if  $f$  and  $g$  are represented by power series with radii of convergence  $R_f$  and  $R_g$  respectively then  $(f + g)$  and  $(f \cdot g)$  represented by power series with radii of convergence  $\geq \min(R_f, R_g)$

Proof. Exercise □

**Corollary 1.2.2 :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and let  $f, g : \Omega \rightarrow \mathbb{C}$ . If  $f$  and  $g$  are both analytic on  $\Omega$ , then the same is for  $(f + g)$  and  $(f \cdot g)$ .

**Proposition 1.2.3 (Analyticity  $\implies$  Continuity) :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and let  $z_0 \in \Omega$ , Let also  $f : \Omega \longrightarrow \mathbb{C}$  be a map.  
if  $f$  is analytic at  $z_0$  then  $f$  is continuous at  $z_0$

*Proof.* Suppose that  $f$  is analytic at  $z_0$  then there exists  $R > 0$  and a complex sequence  $(a_n)_{n \in \mathbb{N}_0}$  such that  $D(z_0, R) \subset \Omega$  and:

$$f(z) = \sum_{n=1}^{\infty} a_n (z - z_0)^n \quad (\forall z \in D(z_0, R))$$

in particular,  $f(z_0) = a_0$ . Thus for all  $z \in D(z_0, R)$  we have:

$$\begin{aligned} f(z) - f(z_0) &= \sum_{n=1}^{\infty} a_n (z - z_0)^n \\ &= (z - z_0) \sum_{n=1}^{\infty} a_n (z - z_0)^{n-1} \\ &= (z - z_0) \underbrace{\sum_{n=1}^{\infty} a_{n+1} (z - z_0)^n}_{(1)} \end{aligned}$$

By the Hadamard formula, we see that the power series  $\sum_{n=1}^{\infty} a_{n+1} (z - z_0)^n$  has the same radius of convergence as the original power series  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$

$$\left( R' = \frac{1}{\lim_{n \rightarrow \infty} \sup |a_{n+1}|^{\frac{1}{n}}} = \frac{1}{\lim_{n \rightarrow \infty} \sup |a_n|^{\frac{1}{n-1}}} \right)$$

Consequently, the power series  $\sum_{n=1}^{\infty} a_{n+1} (z - z_0)^n$  converges absolutely for  $|z - z_0| < R$ , let  $r \in \mathbb{R}$  such that  $0 < r < R$ . then for all  $z \in D(z_0, r)$ , we have from (1) the estimate:

$$\begin{aligned} |f(z) - f(z_0)| &= |z - z_0| \cdot \left| \sum_{n=1}^{\infty} a_{n+1} (z - z_0)^n \right| \\ &\leq |z - z_0| \sum_{n=1}^{\infty} |a_{n+1}| |z - z_0|^n \\ &\leq |z - z_0| \underbrace{\sum_{n=1}^{\infty} |a_{n+1}| \cdot r^n}_{(\text{since } r < R)} \end{aligned}$$

Taking the limit as  $z \rightarrow z_0$ , we conclude that  $\lim_{z \rightarrow z_0} f(z) = f(z_0)$ , so  $f$  is continuous at  $z_0$ .  $\square$

**Corollary 1.2.4 (Immediate) :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and  $f : \Omega \longrightarrow \mathbb{C}$ . If  $f$  is analytic on  $\Omega$ , then  $f$  is continuous on  $\Omega$ .

**Proposition 1.2.5 (Composition of Analytic functions) :** Let  $\Omega_1$  and  $\Omega_2$  be two non empty open subsets of  $\mathbb{C}$  and let  $f : \Omega_1 \rightarrow \Omega_2$  and  $g : \Omega_2 \rightarrow \mathbb{C}$  be two maps. Let also  $z_0 \in \Omega_1$ . If  $f$  is analytic at  $z_0$  and  $g$  is analytic at  $f(z_0)$ , then  $(g \circ f)$  is analytic at  $z_0$ .

*Proof.* Exercise □

**Corollary 1.2.6 (Immediate) :** Let  $\Omega_1$  and  $\Omega_2$  be two non empty open subsets of  $\mathbb{C}$  and let  $f : \Omega_1 \rightarrow \Omega_2$  and  $g : \Omega_2 \rightarrow \mathbb{C}$  be two maps. If  $f$  is analytic on  $\Omega_1$  and  $g$  is analytic on  $\Omega_2$  then  $(g \circ f)$  is analytic on  $\Omega_1$ .

**Proposition 1.2.7 (Quotient of Analytic Functions) :** Let  $\Omega$  be a non empty open subsets of  $\mathbb{C}$  and let  $z_0 \in \Omega$ . let also  $f, g : \Omega \rightarrow \mathbb{C}$  be two functions which are both analytic at  $z_0$  and such that  $g(z_0) \neq 0$ . Then the function  $\frac{f}{g}$  is analytic at  $z_0$

*Proof.* since  $g(z_0) \neq 0$  then the function  $h : w \rightarrow \frac{1}{w}$  is analytic at  $g(z_0)$  (as seen in previous examples), therefore, by proposition 5, the function  $\frac{1}{g} = h \circ g$  is analytic at  $z_0$ .

It then follows from proposition 1 that the product  $f \cdot \left(\frac{1}{g}\right)$  is analytic at  $z_0$  □

**Corollary 1.2.8 (Immediate) :** Let  $\Omega$  be a non empty open subset of  $\mathbb{C}$  and let  $f, g : \Omega \rightarrow \mathbb{C}$  be two analytic functions on  $\Omega$  such that  $g(z) \neq 0$ . for every  $z \in \Omega$ . Then the function  $\frac{f}{g}$  is analytic on  $\Omega$ .

**Example:** Every rational function is analytic on its domain of definition. this is because a rational function is a quotient of two polynomials, and polynomials are analytic on  $\mathbb{C}$

## 1.3 Power series define Analytic functions

**Theorem 1.3.1 :** A power series with a positive radius of converges defines an analytic function on its disk of convergence

*Proof.* Let  $\sum_{n=1}^{\infty} a_n(z - z_0)^n$  be a power series ( $z_0 \in \mathbb{C}, (a_n)_{n \in \mathbb{N}} \subset \mathbb{C}$ ) with radius of convergence

$R > 0$ . define the function  $f$  on the disk  $D(z_0, R)$  by:

$$f(z) = \sum_{n=1}^{\infty} a_n (z - z_0)^n$$

we must show that  $f$  is analytic on  $D(z_0, R)$ . arbitrary we will show that  $f$  is analytic at  $z_1$  for  $z \in D(z_1, R - |z_1 - z_0|)$ , we have

$$|z - z_0| \stackrel{I.I}{\leq} \underbrace{|z - z_1|}_{< R - |z_1 - z_0|} + |z_1 - z_0|$$

Thus  $D(z_1, R - |z_1 - z_0|) \subset D(z_0, R)$ , so the power series  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$  converges absolutely. so:

$$\begin{aligned} f(z) &= \sum_{n=1}^{\infty} a_n (z - z_0)^n \\ &= \sum_{n=1}^{\infty} a_n ((z - z_1) + (z_1 - z_0))^n \\ &= \sum_{n=1}^{\infty} a_n \sum_{k=0}^n \binom{n}{k} (z - z_1)^k (z_1 - z_0)^{n-k} \\ &= \sum_{k=1}^{\infty} \left( \sum_{n=k}^{\infty} a_n \binom{n}{k} (z_1 - z_0)^{n-k} \right) (z - z_1)^k \end{aligned}$$

The interchange of summation is justified by the absolute convergence of the double series for  $z \in D(z_1, R - |z_1 - z_0|)$ .

This express  $f(z)$  as a power series in  $(z - z_1)$  in the disk  $D(z_1, R - |z_1 - z_0|)$  proving that  $f$  is analytic at  $z_1$ . since  $z_1$  was arbitrary in  $D(z_0, R)$  then  $f$  is analytic on  $D(z_0, R)$ .  $\square$