

## Differentiability

A multivariable function  $\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be differentiable at  $\mathbf{a} \in \mathbb{R}^n$  if all its partial derivatives  $\frac{\partial \mathbf{f}}{\partial x_i}$  exist *and* the local tangent plane  $\mathbf{h}(\mathbf{x}) = \mathbf{f}(\mathbf{a}) + \nabla \mathbf{f}(\mathbf{a}) \cdot (\mathbf{x} - \mathbf{a})$  is a “good” approximation of the function:

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \frac{\mathbf{f}(\mathbf{x}) - \mathbf{h}(\mathbf{x})}{|\mathbf{x} - \mathbf{a}|} = 0.$$

A nice little shortcut, because we’re college math students and not hardcore real analysts, is that a function is differentiable if its partial derivatives are all continuous.

## Chain Rule

The chain rule deals with the *composition* of two functions. In one-variable calculus, the chain rule looks something like, for some function  $f = g \circ h$  (that is,  $f(x) = g(h(x))$ ),

$$\frac{df}{dx} = \frac{df}{dh} \frac{dh}{dx} = \frac{dg}{dh} \frac{dh}{dx} = g'(h(x))h'(x).$$

For multi-variable functions ( $D$ , the chain rule takes on a matrix form:

$$D(\mathbf{g} \circ \mathbf{h}) = D(\mathbf{g})D(\mathbf{h}).$$

**Example** Consider  $\mathbf{f}: \mathbb{R}^3 \rightarrow \mathbb{R}^2$  given by

$$\mathbf{f}(x_1, x_2, x_3) = (x_1 - x_2, x_1 x_2 x_3),$$

and  $\mathbf{x}: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  given by

$$\mathbf{x}(t_1, t_2) = (t_1 t_2, t_1^2, t_2^2).$$

**Context** Here’s how the mapping looks:

$$\mathbb{R}^2 \xrightarrow{\mathbf{x}} \mathbb{R}^3 \xrightarrow{\mathbf{f}} \mathbb{R}^2,$$

or

$$(t_1, t_2) \xrightarrow{\mathbf{x}} (\underbrace{x_1(t_1, t_2)}_{t_1 t_2}, \underbrace{x_2(t_1, t_2)}_{t_1^2}, \underbrace{x_3(t_1, t_2)}_{t_2^2}) \xrightarrow{\mathbf{f}} (\underbrace{f_1(x_1, x_2, x_3)}_{x_1 - x_2}, \underbrace{f_2(x_1, x_2, x_3)}_{x_1 x_2 x_3}).$$

Notice the overall (composed) mapping is then

$$\mathbb{R}^2 \xrightarrow{\mathbf{f} \circ \mathbf{x}} \mathbb{R}^2.$$

**Solution (method 1): tedious substitution** Find the derivative of the composition of  $f$  and  $x$ ,

$$D(\mathbf{f} \circ \mathbf{x})(\mathbf{t}).$$

We can simply plug in the composed function

$$(\mathbf{f} \circ \mathbf{x})(\mathbf{t}) = (x_1 - x_2, x_1 x_2 x_3) = (t_1 t_2 - t_1^2, (t_1 t_2)(t_1^2)(t_2^2)) = (t_1 t_2 - t_1^2, t_1^3 t_2^3)$$


and take the derivative:

$$D(\mathbf{f} \circ \mathbf{x})(\mathbf{t}) = \begin{pmatrix} \partial_{t_1} (t_1 t_2 - t_1^2) & \partial_{t_2} (t_1 t_2 - t_1^2) \\ \partial_{t_1} (t_1^3 t_2^3) & \partial_{t_2} (t_1^3 t_2^3) \end{pmatrix} = \begin{pmatrix} t_2 - 2t_1 & t_1 \\ 3t_1^2 t_2^3 & 3t_1^3 t_2^2 \end{pmatrix}.$$


**Solution (method 2): chain rule (*better!!!*)** Alternatively, apply the chain rule:

$$\begin{aligned}
 D(\mathbf{f} \circ \mathbf{x})(\mathbf{t}) &= D\mathbf{f}(\mathbf{x})D\mathbf{x}(\mathbf{t}) \\
 &= \begin{pmatrix} \partial_{x_1}(x_1 - x_2) & \partial_{x_2}(x_1 - x_2) & \partial_{x_3}(x_1 - x_2) \\ \partial_{x_1}(x_1 x_2 x_3) & \partial_{x_2}(x_1 x_2 x_3) & \partial_{x_3}(x_1 x_2 x_3) \end{pmatrix} \begin{pmatrix} \partial_{t_1}(t_1 t_2) & \partial_{t_2}(t_1 t_2) \\ \partial_{t_1}(t_1^2) & \partial_{t_2}(t_1^2) \\ \partial_{t_1}(t_2^2) & \partial_{t_2}(t_2^2) \end{pmatrix} \\
 &= \begin{pmatrix} 1 & -1 & 0 \\ x_2 x_3 & x_1 x_3 & x_1 x_2 \end{pmatrix} \begin{pmatrix} t_2 & t_1 \\ 2t_1 & 0 \\ 0 & 2t_2 \end{pmatrix} \\
 &= \begin{pmatrix} t_2 - 2t_1 & t_1 \\ x_2 x_3 t_2 + 2x_1 x_3 t_1 & x_2 x_3 t_1 + 2x_1 x_2 t_2 \end{pmatrix} \\
 &= \begin{pmatrix} t_2 - 2t_1 & t_1 \\ (t_1^2)(t_2^2)t_2 + 2(t_1 t_2)(t_2^2)t_1 & (t_1^2)(t_2^2)t_1 + 2(t_1 t_2)(t_1^2)t_2 \end{pmatrix} \\
 &= \begin{pmatrix} t_2 - 2t_1 & t_1 \\ 3t_1^2 t_2^3 & 3t_1^3 t_2^2 \end{pmatrix}.
 \end{aligned}$$

**Solution (method 3?): cool diagrams**



The partial derivative of the function's  $i$ -th component  $f_i$  with respect to some input  $t_j$  can be found by tracing all “paths” on the diagram from  $t_j$  to  $f_i$ . For example, the partial derivative of  $f_2$  with respect to  $t_1$  is found by



$$\partial_{t_1} f_2 = \partial_{x_1} f_2 \cdot \partial_{t_1} x_1 + \partial_{x_2} f_2 \cdot \partial_{t_1} x_2 + \partial_{x_3} f_2 \cdot \partial_{t_1} x_3$$

Why does it work? Turns out, this is just another way to do, or *visualize*, derivative matrix multiplication. Tracing the “paths” of each component is the same thing as component-wise matrix multiplication:

$$\begin{aligned} D_{\mathbf{t}} \mathbf{f} &= D_{\mathbf{x}} \mathbf{f} D_{\mathbf{t}} \mathbf{x} \\ &= \begin{pmatrix} \partial_{x_1} f_1 & \partial_{x_2} f_1 & \partial_{x_3} f_1 \\ \partial_{x_1} f_2 & \partial_{x_2} f_2 & \partial_{x_3} f_2 \end{pmatrix} \begin{pmatrix} \partial_{t_1} x_1 & \partial_{t_2} x_1 \\ \partial_{t_1} x_2 & \partial_{t_2} x_2 \\ \partial_{t_1} x_3 & \partial_{t_2} x_3 \end{pmatrix} \\ &= \begin{pmatrix} \partial_{x_1} f_1 \partial_{t_1} x_1 + \partial_{x_2} f_1 \partial_{t_1} x_2 + \partial_{x_3} f_1 \partial_{t_1} x_3 & \partial_{x_1} f_1 \partial_{t_2} x_1 + \partial_{x_2} f_1 \partial_{t_2} x_2 + \partial_{x_3} f_1 \partial_{t_2} x_3 \\ \partial_{x_1} f_2 \partial_{t_1} x_1 + \partial_{x_2} f_2 \partial_{t_1} x_2 + \partial_{x_3} f_2 \partial_{t_1} x_3 & \partial_{x_1} f_2 \partial_{t_2} x_1 + \partial_{x_2} f_2 \partial_{t_2} x_2 + \partial_{x_3} f_2 \partial_{t_2} x_3 \end{pmatrix} \end{aligned}$$

## Implicit Derivatives

Consider some surface defined by the equation  $F(x, y, z) = 0$ . Implicitly take partial derivatives with respect to each variable, considering that  $z$  to be a function of  $x$  and  $y$ :

$$\partial_x F + \partial_z F \partial_x z = 0,$$

$$\partial_y F + \partial_z F \partial_y z = 0.$$

Consequently,

$$\begin{aligned} \partial_x z &= -\frac{\partial_x F}{\partial_z F}, \\ \partial_y z &= -\frac{\partial_y F}{\partial_z F}. \end{aligned}$$

**Example** Consider the surface given by


$$x^2 + y^2 + z^2 - 1 = 0,$$

a sphere with radius 1. The partial derivatives of  $z$  with respect to  $x$  and  $y$  are then

$$\begin{aligned}\partial_x z &= -\frac{\partial_x(x^2 + y^2 + z^2 - 1)}{\partial_z(x^2 + y^2 + z^2 - 1)} = -\frac{2x}{2z} = -\frac{x}{z} \\ \partial_y z &= -\frac{y}{z}.\end{aligned}$$

## Directional Derivatives

Consider the derivative of a function  $\mathbf{f}$  along some direction  $\hat{\mathbf{v}}$ . The *directional derivative* can be thought of as a derivative of a function inside a “cross section”:



The line along the direction of differentiation is parametrized by some

$$\mathbf{l}(t) = \mathbf{p} + t\hat{\mathbf{v}},$$

so that the function along the line

$$\mathbf{f}|_{\mathbf{l}(t)} = \mathbf{f}(\mathbf{p} + t\hat{\mathbf{v}}),$$

so that the directional derivative can be found by the chain rule

$$D_{\hat{\mathbf{v}}} \mathbf{f} = d_t \mathbf{f}(\mathbf{p} + t\hat{\mathbf{v}}) = D\mathbf{f} \cdot \hat{\mathbf{v}}.$$