

1

Results

1.1 Abstract

Cite more

We present finite-size scaling results using the corner transfer matrix renormalization group method on two-dimensional classical square lattices [11]. We compare the results of conventional finite-size scaling in the system size N with scaling in the number of states kept during the renormalization step of the algorithm, denoted by m . Such a comparison was first done in [10]. We highlight the areas in which method excels over the other.

Calculate critical temperature and exponents using information that is directly extractable from the corner transfer matrix.

1.2 Introduction

The first direct comparison of finite-size scaling in the system size N with scaling in the bond dimension of the corner transfer matrix renormalization group method m was done in [10]. In explaining the basic concepts, we largely follow this paper.

The error in the approximation of the partition function (and thus all thermodynamic quantities) in the thermodynamic limit with the corner transfer matrix method depends on two characteristic length scales. The first is the size of the system N . After n steps of the infinite-system algorithm, we have

$$N = 2n + 1. \tag{1.1}$$

The second length scale is related to the finite bond dimension m . Baxter [2], and later Östlund and Rommer [13] (in the context of one-dimensional quantum systems) showed that in the thermodynamic limit, CTMRG and




Figure 1.1: Upper panel: expectation value of the central spin $\langle \sigma_0 \rangle$ after n CTMRG steps. m is chosen such that the truncation error is smaller than 10^{-6} . Lower panel: $\langle \sigma_0 \rangle$ for systems with bond dimension m .

DMRG are variational optimizations in the space of matrix product states.

Can extend this idea a bit.

It is known that an MPS-ansatz with finite bond dimension inherently limits the correlation length of the system to a finite value [18]. Hence, thermodynamic quantities obtained from the CTMRG algorithm with finite m , in the limit $N \rightarrow \infty$, cannot diverge and must show finite-size effects similar to those of some effective finite system of size $N_{\text{eff}}(m)$ depending on the bond dimension m .

Figure 1.1 shows the behaviour of the order parameter of the two-dimensional Ising model for systems of finite-size, where m has been chosen such that the truncation error is smaller than 10^{-6} , and for systems of finite m , where the result is converged in the system size N . The results look very similar and support the above claim.

Order parameter is not the same as magnetization central spin. Where to explain this?

1.2.1 Definition of N_{eff} in terms of the correlation length at T_c

In the thermodynamic limit (corresponding to infinite m and N), we have the following expression for the correlation length of a classical system [3]

$$\xi(T) = \frac{1}{\log\left(\frac{T_0}{T_1}\right)}. \quad (1.2)$$

Here, T_0 and T_1 are the largest and second-largest eigenvalues of the row-to-row transfer matrix T , respectively. With N tending towards infinity and finite m , near the critical point $\xi(T)$ should obey a scaling law of the form

$$\xi(T, m) = N_{\text{eff}}(m) \mathcal{F}(N_{\text{eff}}(m)/\xi(T)) \quad (1.3)$$

with

$$\mathcal{F}(x) = \begin{cases} \text{const} & \text{if } x \rightarrow 0, \\ x^{-1} & \text{if } x \rightarrow \infty. \end{cases} \quad (1.4)$$

Hence, the effective length scale corresponding to the finite bond dimension m is proportional to the correlation length of the system at the critical point $t = 0$.

$$N_{\text{eff}}(m) \propto \xi(T = T_c, m). \quad (1.5)$$

Under this assumption, the order parameter should obey the following scaling relation at the critical temperature

$$M(T = T_c, m) \propto \xi(T = T_c, m)^{-\beta/\nu}. \quad (1.6)$$

The left panel of Figure 1.2 shows that this scaling relation holds. The fit yields $\frac{\beta}{\nu} \approx 0.125(5)$, close to the true value of $\frac{1}{8}$.

The right panel shows the conventional finite-size scaling relation

$$M(T = T_c, N) \propto N^{-\beta/\nu}, \quad (1.7)$$

yielding $\beta/\nu \approx 0.1249(1)$, which can be systematically improved by fitting to larger system sizes, obtained with a fixed truncation error.




Figure 1.2: Left panel: fit to the relation in Equation 1.6, yielding $\frac{\beta}{\nu} \approx 0.125(5)$. The data points are obtained from simulations with $m = 2, 4, \dots, 64$. The smallest 10 values of m have not been used for fitting, to diminish correction terms to the basic scaling law. Right panel: fit to conventional finite-size scaling law given in Equation 1.7.

In the case of scaling in correlation length $\xi(m)$, the exponent does not improve when taking bigger values of m , while keeping the termination criterion (relative change of singular values) fixed. This points to a flaw in the termination criterion of the algorithm.

Furthermore, the correlation length $\xi(m)$ shows characteristic half-moon patterns on a log-log scale, stemming from the degeneracies in the corner transfer matrix spectrum. This makes the data harder to interpret, since the effect of increasing m depends on how much of the spectrum is currently retained.

Talk about how to alleviate this partially by using entropy S as length scale.

To further test the hypothesis that N and $\xi(m)$ are the only relevant length scales, the authors of [10] propose a scaling relation for the order parameter M at the critical temperature of the form

$$M(N, m) = N^{-\beta/\nu} \mathcal{G}(\xi(m)/N) \quad (1.8)$$

with

$$\mathcal{G}(x) = \begin{cases} \text{const} & \text{if } x \rightarrow \infty, \\ x^{-\beta/\nu} & \text{if } x \rightarrow 0, \end{cases} \quad (1.9)$$




Figure 1.3: Scaling function $\mathcal{G}(\xi(m)/N)$ given in Equation 1.8.

meaning that Equation 1.8 reduces to Equation 1.7 in the limit $\xi(m) \gg N$ and to Equation 1.6 in the limit $N \gg \xi(m)$. Figure 1.3 shows that the scaling relation of Equation 1.8 is justified.

Figure 1.4 shows the cross-over behaviour from the N -limiting regime, where $M(N, m) \propto N^{-\beta/\nu}$ to the $\xi(m)$ -limiting regime, where $M(N, m)$ does not depend on N .

1.2.2 Scaling relations away from the critical point

In general, the position of the critical point is not known. In that situation, the scaling relation in Equation 1.8 is no longer valid. Instead, in the limit $N \rightarrow \infty$, we should have

$$M(t, m) \propto \xi(m)^{-\beta/\nu} \mathcal{P}(t\xi(m)^{1/\nu}), \quad (1.10)$$

which is confirmed in Figure 1.5.




Figure 1.4: Behaviour of the order parameter at fixed m as function of the number of renormalization steps n . For small n , all curves coincide, since the system size is the only limiting length scale. For large enough n , the order parameter is only limited by the length scale $\xi(m)$. In between, there is a cross-over described by $\mathcal{G}(\xi(m)/N)$, given in Equation 1.8.

1.3 Finite-entropy scaling

Articles to cite:

- [16]: postulates exponent κ , provides numerical support and extracts basic results for 1D quantum systems with periodic boundary conditions. Also compares to nishino 2D results. Replicate his experiments? Because CTM seems to give different value for κ .
- [1]: establishes $\xi \propto m^{1.3}$ for correlation length of converged DMRG ground states of a gapless systems of free fermions.
- [14]: assumes existence of κ and compares finite-size scaling with finite- m scaling for 1D quantum systems with periodic boundary conditions.




Figure 1.5: Scaling function $\mathcal{P}(t\xi(m)^{1/\nu})$ in Equation 1.10

- [15]: proves expression for κ in asymptotically large m limit that depends only on central charge c , but with corrections of order $1/\log(\chi)$.
- [17]: Also finds $S_A \propto \log(L)$ for 1D quantum systems.
- [4]: proves relation $S_A \propto \mathcal{A}(c/6) \log(\xi)$, with \mathcal{A} the number of boundary points (verified for integrable 2d classical lattice systems).
- [5]: proves relation for classical eight vertex model

Check for validity of $\xi(m) \propto m^\kappa$

- entropy scaling at T_c (using exact value of c).
- correlation length scaling at T_c .
- data collapse of order parameter (using exact value of T_c, β, ν)

1.4 Finding the critical point by maximizing entropy

Articles to cite:

- [6]: theory and numerical support for directly extracting information from ctm spectrum, equivalence to 1D quantum systems.
- [8]: characterisation of phase transition by ctm spectrum for $q = 6$ clock model.
- [7]: characterisation of phase transition by ctm spectrum for truncated tetrahedral model.
- [9]: characterisation of phase transition by ctm spectrum for Widom-Rowlinson models.
- [12]: for XY and Ising model, proves that next-to-nearest neighbor entanglement peaks at critical point (though not nearest-neighbor entanglement.)

Bibliography

- [1] Martin Andersson, Magnus Boman, and Stellan Östlund. “Density-matrix renormalization group for a gapless system of free fermions”. In: *Physical Review B* 59.16 (1999), p. 10493.
- [2] RJ Baxter. “Variational approximations for square lattice models in statistical mechanics”. In: *Journal of Statistical Physics* 19.5 (1978), pp. 461–478.
- [3] Rodney J Baxter. *Exactly solved models in statistical mechanics*. Elsevier, 1982. Chap. 7.
- [4] Pasquale Calabrese and John Cardy. “Entanglement entropy and quantum field theory”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06 (2004), P06002.
- [5] Elisa Ercolessi, Stefano Evangelisti, and Francesco Ravanini. “Exact entanglement entropy of the XYZ model and its sine-Gordon limit”. In: *Physics Letters A* 374.21 (2010), pp. 2101–2105.
- [6] Ching-Yu Huang, Tzu-Chieh Wei, and Roman Orus. “Holographic encoding of universality in corner spectra”. In: *arXiv preprint arXiv:1702.01598* (2017).
- [7] Roman Krčmar, Andrej Gendiar, and Tomotoshi Nishino. “Phase diagram of a truncated tetrahedral model”. In: *Phys. Rev. E* 94 (2 Aug. 2016), p. 022134. DOI: 10.1103/PhysRevE.94.022134. URL: <https://link.aps.org/doi/10.1103/PhysRevE.94.022134>.
- [8] Roman Krčmár, Andrej Gendiar, and Tomotoshi Nishino. “Phase transition of the six-state clock model observed from the entanglement entropy”. In: *arXiv preprint arXiv:1612.07611* (2016).
- [9] Roman Krčmár and Ladislav Šamaj. “Reentrant disorder-disorder transitions in generalized multicomponent Widom-Rowlinson models”. In: *Physical Review E* 92.5 (2015), p. 052103.

- [10] T Nishino, K Okunishi, and M Kikuchi. “Numerical renormalization group at criticality”. In: *Physics Letters A* 213.1-2 (1996), pp. 69–72.
- [11] Tomotoshi Nishino and Kouichi Okunishi. “Corner transfer matrix renormalization group method”. In: *Journal of the Physical Society of Japan* 65.4 (1996), pp. 891–894.
- [12] Tobias J Osborne and Michael A Nielsen. “Entanglement in a simple quantum phase transition”. In: *Physical Review A* 66.3 (2002), p. 032110.
- [13] Stellan Östlund and Stefan Rommer. “Thermodynamic limit of density matrix renormalization”. In: *Physical review letters* 75.19 (1995), p. 3537.
- [14] B Pirvu et al. “Matrix product states for critical spin chains: Finite-size versus finite-entanglement scaling”. In: *Physical review b* 86.7 (2012), p. 075117.
- [15] Frank Pollmann et al. “Theory of finite-entanglement scaling at one-dimensional quantum critical points”. In: *Physical review letters* 102.25 (2009), p. 255701.
- [16] L Tagliacozzo et al. “Scaling of entanglement support for matrix product states”. In: *Physical review b* 78.2 (2008), p. 024410.
- [17] Guifre Vidal et al. “Entanglement in quantum critical phenomena”. In: *Physical review letters* 90.22 (2003), p. 227902.
- [18] Michael M Wolf et al. “Quantum phase transitions in matrix product systems”. In: *Physical review letters* 97.11 (2006), p. 110403.