

Contents

1	Results clock model	1
1.1	Abstract	1
1.2	Introduction	1
1.3	Previous work	2
1.4	Results	2

1

Results clock model

1.1 Abstract

We present results of scaling in bond dimension and system size with the CTMRG algorithm for the five- and six-state clock model.

1.2 Introduction

In the field of phase transitions and critical phenomena, the two-dimensional topological phase transition discovered by Kosterlitz and Thouless [1, 2] received much attention. This phase transition is characterized not by an order parameter which indicates a breaking of symmetry, but by the proliferation of topological defects.

The XY model consists of planar rotors on the square lattice. It exhibits the Kosterlitz-Thouless (BK) phase transition and by the Mermin-Wagner-Hohenberg theorem the symmetry of the ground state is broken for all temperatures, due to the $O(2)$ (planar rotational) symmetry of the potential [3, 4].

The q -state clock model possesses the discrete \mathbb{Z}_q symmetry and is an interpolation between the Ising model, which corresponds to $q = 2$, and the XY model, which corresponds to $q \rightarrow \infty$. Its energy function is given by

$$H_q = - \sum_{\langle ij \rangle} \cos(\theta_i - \theta_j). \quad (1.1)$$

It has been proven that for high enough q , this model indeed exhibits a Kosterlitz-Thouless transition [5].

In the Villain formulation of the potential [6], it has been proven that the model has two phase transitions for $q \geq 5$: a symmetry broken phase for

$T < T_1$, an intermediate phase with power law decay of the correlation function, and a high-temperature phase with exponential decay of the correlation function for $T > T_2$. The transition at T_2 is a BK-transition, and for a broad range of temperatures, the thermodynamic behaviour becomes identical to the XY model for high enough q [7].

1.3 Previous work

1.4 Results

Bibliography

- [1] John Michael Kosterlitz and David James Thouless. “Ordering, metastability and phase transitions in two-dimensional systems”. In: *Journal of Physics C: Solid State Physics* 6.7 (1973), p. 1181.
- [2] JM Kosterlitz. “The critical properties of the two-dimensional xy model”. In: *Journal of Physics C: Solid State Physics* 7.6 (1974), p. 1046.
- [3] N David Mermin and Herbert Wagner. “Absence of ferromagnetism or antiferromagnetism in one-or two-dimensional isotropic Heisenberg models”. In: *Physical Review Letters* 17.22 (1966), p. 1133.
- [4] PC Hohenberg. “Existence of long-range order in one and two dimensions”. In: *Physical Review* 158.2 (1967), p. 383.
- [5] Jürg Fröhlich and Thomas Spencer. “The Kosterlitz-Thouless transition in two-dimensional abelian spin systems and the Coulomb gas”. In: *Communications in Mathematical Physics* 81.4 (1981), pp. 527–602.
- [6] J Villain. “Theory of one-and two-dimensional magnets with an easy magnetization plane. II. The planar, classical, two-dimensional magnet”. In: *Journal de Physique* 36.6 (1975), pp. 581–590.
- [7] Cintia M Lapilli, Peter Pfeifer, and Carlos Wexler. “Universality away from critical points in two-dimensional phase transitions”. In: *Physical review letters* 96.14 (2006), p. 140603.