HKUSTx: ELEC1200.1x A System View of Communications: From Signals to Packets (Part 1)

KarenWest (/dashboard)

Courseware (/courses/HKUSTx/ELEC1200.1x/3T2014/courseware) Course Info (/courses/HKUSTx/ELEC1200.1x/3T2014/info)

 $Course\ Outline\ (/courses/HKUSTx/ELEC1200.1x/3T2014/05fb01b36df14eb99ab54545dabc47f6/)$

Grading Scheme (/courses/HKUSTx/ELEC1200.1x/3T2014/6e2be4dac3e44b4d9f812e7b5a5d5a29/)

으 인 Instructors (/courses/HKUSTx/ELEC1200.1x/3T2014/674fdd6887fe4f4bb73b984df4a5675b/)

Resources (/courses/HKUSTx/ELEC1200.1x/3T2014/a6a8267fef364cccbccd0128d091f11c/)

Discussion (/courses/HKUSTx/ELEC1200.1x/3T2014/discussion/forum)

Progress (/courses/HKUSTx/ELEC1200.1x/3T2014/progress)

6.3 QUIZ QUESTION 1 (1/1 point)

Which of the following is an equivalent recursive specification for the sequence $x(n) = 0.5 \cdot n + 1$?

Select the correct answer.

$$x(0) = 0$$
, $x(n) = x(n-1) + 0.5$

$$x(0)=0$$
, $x(n)=0.5\cdot x(n-1)$

$$x(n) = 0.5 \cdot (n+2)$$

$$\bullet$$
 $x(0) = 1$, $x(n) = x(n-1) + 0.5$

$$x(0)=1$$
, $x(n)=0.5\cdot x(n-1)$

EXPLANATION

Base cases:

By inspection, the explicit formula and the recursive specification both give x(0) = 1.

For n = 1, using the explicit formula, $x(1) = 0.5 \cdot 1 + 1 = 1.5$

For n = 1, using the recursive formulation, x(1)=x(0)+0.5=1+0.5=1.5

Thus, the explicit formula and the recursive formulation are the same for n = 1.

Inductive step:

Assume that for
$$n \geq 1$$
, $x(n) = 0.5 \cdot n + 1$ and $x(n) = x(n-1) + 0.5$.

By the recursive formulation, x(n+1)=x(n)+0.5

Substituting $x(n) = 0.5 \cdot n + 1$, we obtain

$$x(n+1) = (0.5 \cdot n + 1) + 0.5 = 0.5 \cdot (n+1) + 1$$


Hide Answer

You have used 2 of 2 submissions

Help

6.3 QUIZ QUESTION 2 (1/1 point)

The response of a channel, y(n), to an input, x(n), can be modelled by the following recursive system:



Assume x(0:4)=[01110], and y(0)=0. Determine the value of the output y(2).

Please key in the numerical value of your answer to two significant digits in the box provided below.

0.72

0.72

Answer: 0.72

EXPLANATION

This channel response can be determined iteratively using $y(n) = 0.2 \cdot y(n-1) + 0.6 \cdot x(n)$

$$y(1) = 0.2 \cdot y(0) + 0.6 \cdot x(1) = 0.2 \cdot 0 + 0.6 \cdot 1 = 0.6$$

$$y(2) = 0.2 \cdot y(1) + 0.6 \cdot x(2) = 0.2 \cdot 0.6 + 0.6 \cdot 1 = 0.72$$

Check

Save

Hide Answer

You have used 1 of 3 submissions



Follow Us

UTx and many other universities. Topics include biology, business, che and the contract of the co engineering, food and nutrition, history, humanities, law, literature, math, medicine, music, philosophy, physics, science, statistics and more. EdX is a non-profit online initiative created by founding partners Harvard and MIT.

© 2014 edX, some rights reserved.

erms of Service and Honor Code (https://www.edx.org/edx-terms-service)

Privacy Policy (Revised 4/16/2014) (https://www.edx.org/edx-privacypolicy)

News (https://www.edx.org

Contact (https://www.edx.org /contact)

FAQ (https://www.edx.org /student-faq)

edX Blog (https://www.edx.org /edx-blog)

Donate to edX (https://www.edx.org/donate)

Jobs at edX (https://www.edx.org/jobs)

Facebook /news) https://courses.edx.org/@urses/HKUSTx/Eh...

/EdxOnline)

Meetup

(http://www.meetup.com /edX-Global-Community)

in LinkedIn

(http://www.linkedin.com /company/edx)

Google+

(https://plus.google.com /+edXOnline)

3 of 3 10/12/2014 12:27 PM