

# SOLVING PARTIAL DIFFERENTIAL EQUATIONS WITH NEURAL NETWORKS

FYS-STK4155: PROJECT 3

Morten Ledum, Håkon Kristiansen & Kari Eriksen

 [github.com/mortele/FYS-STK4155](https://github.com/mortele/FYS-STK4155)

December 7, 2018

## Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

# Contents

<b>I</b>	<b>Introduction</b>	<b>3</b>
<b>II</b>	<b>Theory</b>	<b>3</b>
A	The heat equation . . . . .	3
B	Closed form solution . . . . .	3
	Applying the boundary conditons . . . . .	4
	Applying the initial condition . . . . .	4
C	Explicit scheme . . . . .	5
D	Finite difference method . . . . .	5
E	Implicit schemes . . . . .	6
F	The Crank-Nicolson scheme . . . . .	7
G	Stability analysis . . . . .	7
H	Finite element methods . . . . .	7
	Functional spaces and weak forms . . . . .	7
I	Solving differential equations with neural networks . . . . .	7
<b>III</b>	<b>Results and discussion</b>	<b>8</b>
<b>IV</b>	<b>Conclusion</b>	<b>8</b>

## I. INTRODUCTION

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## II. THEORY

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

### A. The heat equation

The heat equation is a partial differential equation (in space  $x$  and time  $t$ ) which describes the evolution of temperature differences in a region of space over a time interval. It is based on *Fourier's law*; the rate of heat flow through a surface is proportional to the temperature gradient across the surface, i.e.

$$\mathbf{q} = -k\nabla T = -k\frac{\partial T}{\partial x}, \quad (1)$$

where  $\mathbf{q}$  denotes the heat flux density,  $k$  is the thermal conductivity of the surface material, and  $T$  represents the temperature. Changes in temperature are proportional to changes in internal energy, with the proportionality constant

being the specific heat capacity  $c_p$ . With the arbitrary energy zero point placed at absolute zero, this can be written as

$$Q = c_p \rho T, \quad (2)$$

with  $Q$  being the internal energy and  $\rho$  denoting the mass density. This is essentially just a restatement of (a shifted) *first law of thermodynamics*, in the absence of applied work. The total heat energy contained in a region  $[a, b]$  is given by the integral

$$\int_a^b dx c_p \rho T(x, t). \quad (3)$$

Integrating over a small region of space and considering the change in internal energy over a short time interval (assuming  $c_p$  and  $\rho$  are both time-independent and spatially homogeneous) gives

$$\begin{aligned} \Delta Q &= c_p \rho \int_x^{x+\Delta x} d\chi \left[ T(\chi, t + \Delta t) - T(\chi, t) \right] \\ &= c_p \rho \int_x^{x+\Delta x} d\chi \int_t^{t+\Delta t} d\tau \frac{\partial T}{\partial \tau}. \end{aligned} \quad (4)$$

Over a short time period  $\Delta t$ , the change in internal energy of a short segment of length  $\Delta x$  must be entirely due to the heat flux in/out of the boundaries,

$$\begin{aligned} \Delta Q &= k \int_t^{t+\Delta t} d\tau \left[ \frac{\partial T(x + \Delta x, \tau)}{\partial x} - \frac{\partial T(x, \tau)}{\partial x} \right] \\ &= k \int_t^{t+\Delta t} d\tau \int_x^{x+\Delta x} d\chi \frac{\partial^2 T}{\partial \chi^2}. \end{aligned} \quad (5)$$

By conservation of energy, the difference between Eq. (4) and Eq. (5) must obviously vanish. Since we are integrating over the same spatial and temporal regions in both equations, this means that the integrand must vanish identically:

$$\frac{k}{c_p \rho} \frac{\partial^2 T}{\partial x^2} = \frac{\partial T}{\partial t}. \quad (6)$$

This is known as the *heat equation* and is a special case of the more general diffusion equation.

### B. Closed form solution

The 1D heat equation may be solved by applying a separation of variables ansatz, i.e. we assume the solution  $u(x, t)$  takes the form

$$u(x, t) = X(x)T(t), \quad (7)$$

with  $X(x)$  carrying all the  $x$ -dependence, and  $T(t)$  carrying the corresponding time dependence. Introducing the compact notation

$$u_x \equiv \partial_x u = \frac{\partial u}{\partial x}, \quad \text{and} \quad u_t \equiv \partial_t u = \frac{\partial u}{\partial t}, \quad (8)$$

we find by insertion of the ansatz into Eq. (6):

$$\begin{aligned} \alpha^2 u_{xx} &= u_t \\ \alpha^2 \partial_x \left[ \partial_x X(x) T(t) \right] &= \partial_t X(x) T(t) \\ \alpha^2 \partial_x \left[ X_x T + X T_x \right] &= X_t T + X T_t \\ \alpha^2 \left[ X_{xx} T + 2 X_x T_x + X T_{xx} \right] &= X T_t \\ \frac{1}{X(x)} \frac{\partial^2 X(x)}{\partial x^2} &= \frac{1}{\alpha^2 T(t)} \frac{\partial T(t)}{\partial t}, \end{aligned} \quad (9)$$

where we defined  $\alpha^2 \equiv k/c_p \rho$  and used the fact that  $X_t = T_x = 0$ . As the left hand side is independent of  $t$  and the right hand side is independent of  $x$ , the equality can only be achieved if both sides are constant. This reduces the original partial differential equation into a set of two ordinary differential equations

$$\frac{1}{X(x)} \frac{\partial^2 X(x)}{\partial x^2} = k \quad (10)$$

$$\frac{1}{\alpha^2 T(t)} \frac{\partial T(t)}{\partial t} = k, \quad (11)$$

with  $k$  an undetermined constant.

Depending on the value of  $k$ , the spatial part has solutions  $X(x) = Ax + B$  (if  $k = 0$ ),  $X(x) = Ae^{\mu x} + Be^{-\mu x}$  (if  $k = \mu^2 > 0$ ), or  $X(x) = Ae^{i\mu x} + Be^{-i\mu x}$  (if  $k = -\mu^2 < 0$ ). The temporal equation has solutions  $T(t) = C$  (if  $k = 0$ ), or  $T(t) = Ce^{\alpha^2 \mu^2 x}$  (otherwise).

### Applying the boundary conditions

In order to make progress, we need to apply the specific boundary and initial conditions. In our case, the boundaries at  $x = 0$  and  $x = L = 1$  vanish, and the initial spatial solution takes the form  $u(x, t = 0) = \sin \pi x$ . If the  $k$  constant of Eq. (10) vanishes, then both constants  $A$  and  $B$  vanish due to the boundary conditions. The same is true if  $k = \mu^2 > 0$ . It follows that the only non-trivial solutions arise when  $k = -\mu^2 < 0$ , in which case we find (left boundary)

$$X(0) = A \cos \mu x + B \sin \mu x = 0 \Rightarrow A = 0$$

and (right boundary)

$$X(1) = B \sin \mu x = 0 \Rightarrow \mu = \pi n.$$

This gives rise to an infinite set of equations—one for each  $n$ —which determine the Fourier coefficients of the initial condition  $u(x, t = 0)$ :

$$u(x, t = 0) = \sum_{n=1}^{\infty} B_n \sin(n\pi x), \quad (12)$$

with

$$B_n = 2 \int_0^1 dx u(x, t = 0) \sin(n\pi x). \quad (13)$$

The temporal equation, Eq. (11), can now be solved only applying the boundary conditions. With the value of  $\mu$  fixed at  $\mu = n\pi$ , we obtain

$$T(t) = e^{-n^2 \pi^2 \alpha^2 t}. \quad (14)$$

### Applying the initial condition

Combining the  $X(x)$  and  $T(t)$  solutions we obtain a series representation of the solution in terms of the Fourier coefficients of the initial condition  $u(x, t = 0)$ , in which the higher frequency modes of the initial solution decays more rapidly than the corresponding lower frequency modes,

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin(n\pi x) e^{-n^2 \pi^2 \alpha^2 t}. \quad (15)$$

It is easy to see that the steady-state solution is achieved when  $u(x, t) = 0$ , since both boundaries (left and right) act as sinks for the initial heat energy contained in the system, and no heat is ever *added*. In our case, the initial condition makes the general solution Eq. (15) take on a very simple form. Applying  $u(x, t = 0) = \sin \pi x$ , it is trivial to evaluate

$$B_n = 2 \int_0^1 dx \sin(\pi x) \sin(n\pi x) = \delta_{1n}, \quad (16)$$

due to the orthogonality of  $\sin(n\pi x)$  and  $\sin(m\pi x)$ . This means  $B_1 = 1$  and  $B_2, B_3, \dots = 0$ , and the full solution to Eq. (6) is given by

$$u(x, t) = \sin(\pi x) e^{-\pi^2 \alpha^2 t}. \quad (17)$$

### C. Explicit scheme

One of the more simple methods in the world of solving ordinary and partial differential equation numerically is the explicit scheme. This is a method that bases itself upon using the previous value of the derivatives in order to find the new ones. We will here derive the explicit scheme for the diffusion equation, Eq. (18).

$$\frac{\partial^2 u(x, t)}{\partial x^2} = \frac{\partial u(x, t)}{\partial t} \quad (18)$$

We begin with Taylor expansion of  $u(x, t)$  in  $t$ .

$$u(x, t + \Delta t) = u(x, t) + \Delta t u'(x, t) + \frac{1}{2} \Delta t^2 u''(x, t) + \dots$$

Truncating the series after the first term we get an expression of the first derivative of  $u$  wrt.  $t$  with the truncation error  $O(\Delta t)$ .

$$u'(x, t) = \frac{u(x, t + \Delta t) - u(x, t)}{\Delta t} + O(\Delta t)$$

This can be simplified if we rewrite the derivative of  $u$  wrt.  $t$  as  $u_t$  and discretize.

$$\begin{aligned} u_t &\approx \frac{u(x_i, t_j + \Delta t) - u(x_i, t_j)}{\Delta t} \\ &\approx \frac{u_{i,j+1} - u_{i,j}}{\Delta t} \end{aligned}$$

This is called the forward Euler method, we move one step forward in time in the integration process using the previous step. This is the simplest version of the explicit scheme. Now we carry out the same procedure for the spatial part.

$$\begin{aligned} u(x + \Delta x, t) &= u(x, t) + \Delta x u'(x, t) + \frac{1}{2} \Delta x^2 u''(x, t) + \frac{1}{3!} \Delta x^3 u'''(x, t) + \dots \\ u(x - \Delta x, t) &= u(x, t) - \Delta x u'(x, t) + \frac{1}{2} \Delta x^2 u''(x, t) - \frac{1}{3!} \Delta x^3 u'''(x, t) + \dots \end{aligned}$$

We truncate at second term in order to get the second order derivative and rewrite to simplify as before.

$$\begin{aligned} u_{xx} &\approx \frac{u(x_i + \Delta x, t_j) - 2u(x_i, t_j) + u(x_i - \Delta x, t_j)}{\Delta x^2} \\ &\approx \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{\Delta x^2} \end{aligned}$$

This is a centered difference method as we are using both previous and latter values in space to find the new second derivative of the spatial

coordinate. Adding the two solutions into the original differential equation we get the following.

$$\begin{aligned} u_t &= u_{xx} \\ u_{i,j+1} - u_{i,j} &= u_{i+1,j} - 2u_{i,j} + u_{i-1,j} \frac{\Delta t}{\Delta x^2} \end{aligned}$$

The full diffusion equation on discretized form can further be rewritten by putting  $\beta = \Delta t / \Delta x^2$ .

$$\begin{aligned} u_{i,j+1} &= \beta(u_{i+1,j} - 2u_{i,j} + u_{i-1,j}) + u_{i,j} \\ &= \beta(u_{i+1,j} + u_{i-1,j}) + (1 - 2\beta)u_{i,j} \end{aligned}$$

### D. Finite difference method

The most straightforward way to solve Eq. (6) numerically is through *finite difference methods*, i.e. discretizing time and space on a grid and Taylor expanding the solution to obtain algebraic equation sets. A multitude of different strategies and schemes exists, but we will consider the *explicit forward Euler* scheme.

The spatial region  $[0, L]$  is discretized by splitting it into  $N$  segments, and considering only the functional values on the points  $x_i = i\Delta x$  with  $i \in [0, N - 1]$ . We denote a function  $f(x, t)$  evaluated at  $x_i$  (and at time  $t$ ) by  $f_i^t = f(x_i, t)$ . Considering  $N$  spatial points gives a step size  $\Delta x$  between each point of

$$\Delta x = \frac{L}{N - 1}. \quad (19)$$

In addition, we introduce a discretization in the temporal dimension with step size  $\Delta t$ .

Let us consider the Taylor expansion of a function  $f(x, t)$  considered at fixed  $t$ ,  $f(x, t)$ , around a spatial point  $x$ . We use the shorthand  $h \equiv (x - a)$ , and consider the expansion at a point  $a \neq x$ . The expansions of  $f(x + h; t)$  and  $f(x - h; t)$  are given by,

$$f(x + h) \approx f(x) + hf'(x) + h^2 f''(x), \quad (20)$$

$$f(x - h) \approx f(x) - hf'(x) + h^2 f''(x), \quad (21)$$

respectively. As  $t$  is considered a fixed parameter for the moment, we suppressed the second functional argument for notational brevity. The shorthand  $f'(x)$  is here used to denote differentiation w.r.t.  $x$ . Note that both equations hold with equality if an overall error term proportional to  $h^3$  is added on the right hand side, i.e.  $O(h^3)$ . Adding Eqs. (20) and (21) and dividing through by  $h^2$  yields

$$\begin{aligned} f(x + h) + f(x - h) &= 2f(x) + h^2 f''(x) + O(h^4) \\ f''(x) &= \frac{f(x + h) - 2f(x) + f(x - h)}{h^2} + O(h^2), \end{aligned} \quad (22)$$

which is the three-point central difference approximation to the second derivative. Note that the resulting error is proportional to  $h^2$  because the third order contributions from Eqs. (20) and (21)— $h^3 f'''(x+h)$  and  $-h^3 f'''(x-h)$ —cancel exactly. A corresponding temporal first order derivative approximation may be found by simply considering Eq. (20), and considering  $h$  to be a small temporal step,  $x$  to be a fixed parameter, and varying  $t$ . This gives

$$f'(t) = \frac{f(t+h) - f(t)}{h} + \mathcal{O}(h), \quad (23)$$

where the error term is proportional to  $h^2$  when disregarding the last term on the right hand side of Eq. (20), which gives  $\mathcal{O}(h)$  after dividing through by  $h$ .

Let us now consider Eq. (22) on the previously defined grid, and take  $h = \Delta x$ . In the Eq. (23) case, we define  $h = \Delta t$ , and equate  $\alpha^2 f''(x)$  with  $f'(t)$  as in the heat equation Eq. (6). The result can be solved for  $f(x, t + \Delta t)$ , i.e. the *next* time step given that the previous step is known:

$$\begin{aligned} \frac{f_i^{j+1} - f_i^j}{\Delta t} &= \alpha^2 \frac{f_{i+1}^j - 2f_i^j + f_{i-1}^j}{\Delta x^2} \\ f_i^{j+1} &= f_i^j + \beta [f_{i+1}^j - 2f_i^j + f_{i-1}^j], \end{aligned} \quad (24)$$

where  $f_i^j$  denotes the discretized  $f(x_i, t_j)$  and

$$\beta \equiv \alpha^2 \left( \frac{\Delta t}{\Delta x^2} \right). \quad (25)$$

Equation (24) is known as the *explicit Euler* scheme, and can be solved directly for  $f(x, t + \Delta t)$  since the right hand side is all known at time step  $t$ .

### E. Implicit schemes

As noted in section D, applying the forward difference approximation for the first derivative in time—derived from the Taylor expansion of a function around the point  $x+h$ —yields the explicit Euler scheme. However, as we will see in section G, the forward method has a rather weak stability criterion which necessitates the use of very small time steps to ensure the solution remains well-behaved.

If we instead employ a backward difference approximation in time (solving Eq. (21) for an  $\mathcal{O}(h)$  approximation of the first derivative) we obtain equations sets which are stable for any combination of  $\Delta x$  and  $\Delta t$ . This allows us to

use more reasonable time steps and ultimately speeds up and improves the computed solution.

However, using this approximation,

$$f'(t) = \frac{f(t) - f(t-h)}{h} + \mathcal{O}(h), \quad (26)$$

introduces additional complexity in that we are no longer able to simply solve for the unknown  $f(x, t + \Delta t)$  in the resulting heat equation approximation. Equating Eqs. (22) and (26) yields the *implicit Euler* scheme:

$$\begin{aligned} \frac{f_i^j - f_i^{j-1}}{\Delta t} &= \alpha^2 \frac{f_{i+1}^j - 2f_i^j + f_{i-1}^j}{\Delta x^2} \\ f_i^{j-1} &= -\beta [f_{i+1}^j + f_{i-1}^j] + (1 + 2\beta) f_i^j. \end{aligned} \quad (27)$$

We note that the only known quantity is the left hand side  $f_i^{j-1} = f(x, t - \Delta t)$ , which means we must solve all the equations for every spatial point  $i$  simultaneously in order to ensure they are all satisfied. This gives rise to a matrix-vector equation in the place of the single algebraic equation needed for the explicit scheme. Denoting  $\gamma \equiv 1 + 2\beta$  we may write the equation set corresponding to the integration from  $t - \Delta t$  to  $t$  as

$$A \mathbf{f}^j = \mathbf{f}^{j-1}, \quad (28)$$

with (zeros omitted)

$$A = \begin{pmatrix} \gamma & -1 & & & \\ -1 & \gamma & -1 & & \\ & -1 & \gamma & -1 & \\ & & & \ddots & \\ & & & -1 & \gamma & -1 \\ & & & & -1 & \gamma \end{pmatrix} \quad (29)$$

and

$$\mathbf{f}^j = \begin{pmatrix} f_0^j \\ f_1^j \\ f_2^j \\ \vdots \\ f_N^j \end{pmatrix}, \quad \mathbf{f}^{j-1} = \begin{pmatrix} f_0^{j-1} \\ f_1^{j-1} \\ f_2^{j-1} \\ \vdots \\ f_N^{j-1} \end{pmatrix} \quad (30)$$

As the matrix  $A$  is tri-diagonal, efficient linear scaling solution algorithms exist which circumvent the traditional solution schemes, e.g. LU decomposition at  $\mathcal{O}(N^3)$ .

## F. The Crank-Nicolson scheme

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

## G. Stability analysis

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

## H. Finite element methods

In the case of the finite difference approximation, the derivatives in the original partial differential equation is transformed into a set of algebraic equations by application of *finite difference* approximations arising from Taylor expansions of the solution function. A fundamentally different, and somewhat more sophisticated, solution strategy entails writing the original differential equation in its *weak form*. This is done by integrating the original equation multiplied by a test function  $v(x)$  over the domain in question. Such a *finite element* approach is normally accompanied by a simple finite element scheme in time when partial differential equations are considered.

## Functional spaces and weak forms

Let us consider the vector space of *all* functions  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ , which we will call  $A(\mathbb{R}^2)$ . Obvi-

ously, the solution to the heat equation is contained in this space, being a real-valued function of two parameters,  $x$  and  $t$ . However, we may consider subspaces with more useful structure than the all-encompassing  $A(\mathbb{R}^2)$ . Assuming the initial and boundary conditions are specified in a sufficiently non-pathological way, the solution at each instant in time will be continuous across the spatial integration domain,  $\Omega$ . In other words, the solution (at some  $\tau \geq 0$ )  $u(x; \tau) \in C(\Omega)$ : The vector space of *continuous* functions  $f : \Omega \rightarrow \mathbb{R}$ . As the heat equation involves a second spatial derivative, we might assume that the solution be twice continuously differentiable, i.e.  $u(x; \tau) \in C^2(\Omega)$ .

It turns out that demanding  $u \in C^2(\Omega)$  is too strict a bound on the behaviour of the solution. For example, a valid solution may be *not* differentiable on a subset  $S \subset \Omega$  with measure zero<sup>1</sup>, known as differentiable almost everywhere (a.e.). It is more useful to consider *weakly differentiable* functions, which encompass the ordinary (strongly) differentiable functions and includes other possible solutions of partial differential equations.

Equipping our functional space with the  $L^1$  integral inner product (and the associated norm), consider a family of functions  $\phi \in C^\infty(\Omega)$ , with  $\phi(\partial\Omega) = 0$ , where we denote by  $\partial\Omega$  the boundary of the integration domain (in one dimension, this is just the end two points). The weak derivative of  $u$  is defined as a function  $v$  for which the following holds

$$\int_{\Omega} dx \quad (31)$$

## I. Solving differential equations with neural networks

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut au-

<sup>1</sup>A set  $S \subset \mathbb{R}$  of measure zeros is a set with vanishing "size," e.g. a finite set of *points*.

gue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

### III. RESULTS AND DISCUSSION

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

### IV. CONCLUSION

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.