

SOLVING PARTIAL DIFFERENTIAL EQUATIONS WITH NEURAL NETWORKS

FYS-STK4155: PROJECT 3

Morten Ledum, Håkon Kristiansen & Kari Eriksen

 github.com/mortele/FYS-STK4155

November 22, 2018

Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Contents

I	Introduction	3
II	Theory	3
A	The heat equation	3
B	Closed form solution	3
	Applying the boundary conditons	4
	Applying the initial condition	4
C	Solving differential equations with neural networks	5
III	Results and discussion	5
IV	Conclusion	5

I. INTRODUCTION

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

II. THEORY

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

A. The heat equation

The heat equation is a partial differential equation (in space x and time t) which describes the evolution of temperature differences in a region of space over a time interval. It is based on *Fourier's law*; the rate of heat flow through a surface is proportional to the temperature gradient across the surface, i.e.

$$\mathbf{q} = -k\nabla T = -k\frac{\partial T}{\partial x}, \quad (1)$$

where \mathbf{q} denotes the heat flux density, k is the thermal conductivity of the surface material, and T represents the temperature. Changes in temperature are proportional to changes in internal energy, with the proportionality constant

being the specific heat capacity c_p . With the arbitrary energy zero point placed at absolute zero, this can be written as

$$Q = c_p \rho T, \quad (2)$$

with Q being the internal energy and ρ denoting the mass density. This is essentially just a restatement of (a shifted) *first law of thermodynamics*, in the absence of applied work. The total heat energy contained in a region $[a, b]$ is given by the integral

$$\int_a^b dx c_p \rho T(x, t). \quad (3)$$

Integrating over a small region of space and considering the change in internal energy over a short time interval (assuming c_p and ρ are both time-independent and spatially homogeneous) gives

$$\begin{aligned} \Delta Q &= c_p \rho \int_x^{x+\Delta x} d\chi \left[T(\chi, t + \Delta t) - T(\chi, t) \right] \\ &= c_p \rho \int_x^{x+\Delta x} d\chi \int_t^{t+\Delta t} d\tau \frac{\partial T}{\partial \tau}. \end{aligned} \quad (4)$$

Over a short time period Δt , the change in internal energy of a short segment of length Δx must be entirely due to the heat flux in/out of the boundaries,

$$\begin{aligned} \Delta Q &= k \int_t^{t+\Delta t} d\tau \left[\frac{\partial T(x + \Delta x, \tau)}{\partial x} - \frac{\partial T(x, \tau)}{\partial x} \right] \\ &= k \int_t^{t+\Delta t} d\tau \int_x^{x+\Delta x} d\chi \frac{\partial^2 T}{\partial \chi^2}. \end{aligned} \quad (5)$$

By conservation of energy, the difference between Eq. (4) and Eq. (5) must obviously vanish. Since we are integrating over the same spatial and temporal regions in both equations, this means that the integrand must vanish identically:

$$\frac{k}{c_p \rho} \frac{\partial^2 T}{\partial x^2} = \frac{\partial T}{\partial t}. \quad (6)$$

This is known as the *heat equation* and is a special case of the more general diffusion equation.

B. Closed form solution

The 1D heat equation may be solved by applying a separation of variables ansatz, i.e. we assume the solution $u(x, t)$ takes the form

$$u(x, t) = X(x)T(t), \quad (7)$$

with $X(x)$ carrying all the x -dependence, and $T(t)$ carrying the corresponding time dependence. Introducing the compact notation

$$u_x \equiv \partial_x u = \frac{\partial u}{\partial x}, \quad \text{and} \quad u_t \equiv \partial_t u = \frac{\partial u}{\partial t}, \quad (8)$$

we find by insertion of the ansatz into Eq. (6):

$$\begin{aligned} \alpha^2 u_{xx} &= u_t \\ \alpha^2 \partial_x \left[\partial_x X(x) T(t) \right] &= \partial_t X(x) T(t) \\ \alpha^2 \partial_x \left[X_x T + X T_x \right] &= X_t T + X T_t \\ \alpha^2 \left[X_{xx} T + 2 X_x T_x + X T_{xx} \right] &= X T_t \\ \frac{1}{X(x)} \frac{\partial^2 X(x)}{\partial x^2} &= \frac{1}{\alpha^2 T(t)} \frac{\partial T(t)}{\partial t}, \end{aligned} \quad (9)$$

where we defined $\alpha^2 \equiv k/c_p \rho$ and used the fact that $X_t = T_x = 0$. As the left hand side is independent of t and the right hand side is independent of x , the equality can only be achieved if both sides are constant. This reduces the original partial differential equation into a set of two ordinary differential equations

$$\frac{1}{X(x)} \frac{\partial^2 X(x)}{\partial x^2} = k \quad (10)$$

$$\frac{1}{\alpha^2 T(t)} \frac{\partial T(t)}{\partial t} = k, \quad (11)$$

with k an undetermined constant.

Depending on the value of k , the spatial part has solutions $X(x) = Ax + B$ (if $k = 0$), $X(x) = Ae^{\mu x} + Be^{-\mu x}$ (if $k = \mu^2 > 0$), or $X(x) = Ae^{i\mu x} + Be^{-i\mu x}$ (if $k = -\mu^2 < 0$). The temporal equation has solutions $T(t) = C$ (if $k = 0$), or $T(t) = Ce^{\alpha^2 \mu^2 x}$ (otherwise).

Applying the boundary conditions

In order to make progress, we need to apply the specific boundary and initial conditions. In our case, the boundaries at $x = 0$ and $x = L = 1$ vanish, and the initial spatial solution takes the form $u(x, t = 0) = \sin \pi x$. If the k constant of Eq. (10) vanishes, then both constants A and B vanish due to the boundary conditions. The same is true if $k = \mu^2 > 0$. It follows that the only non-trivial solutions arise when $k = -\mu^2 < 0$, in which case we find (left boundary)

$$X(0) = A \cos \mu x + B \sin \mu x = 0 \Rightarrow A = 0$$

and (right boundary)

$$X(1) = B \sin \mu x = 0 \Rightarrow \mu = \pi n.$$

This gives rise to an infinite set of equations—one for each n —which determine the Fourier coefficients of the initial condition $u(x, t = 0)$:

$$u(x, t = 0) = \sum_{n=1}^{\infty} B_n \sin(n\pi x), \quad (12)$$

with

$$B_n = 2 \int_0^1 dx u(x, t = 0) \sin(n\pi x). \quad (13)$$

The temporal equation, Eq. (11), can now be solved only applying the boundary conditions. With the value of μ fixed at $\mu = n\pi$, we obtain

$$T(t) = e^{-n^2 \pi^2 \alpha^2 t}. \quad (14)$$

Applying the initial condition

Combining the $X(x)$ and $T(t)$ solutions we obtain a series representation of the solution in terms of the Fourier coefficients of the initial condition $u(x, t = 0)$, in which the higher frequency modes of the initial solution decays more rapidly than the corresponding lower frequency modes,

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin(n\pi x) e^{-n^2 \pi^2 \alpha^2 t}. \quad (15)$$

It is easy to see that the steady-state solution is achieved when $u(x, t) = 0$, since both boundaries (left and right) act as sinks for the initial heat energy contained in the system, and no heat is ever *added*. In our case, the initial condition makes the general solution Eq. (15) take on a very simple form. Applying $u(x, t = 0) = \sin \pi x$, it is trivial to evaluate

$$B_n = 2 \int_0^1 dx \sin(\pi x) \sin(n\pi x) = \delta_{1n}, \quad (16)$$

due to the orthogonality of $\sin(n\pi x)$ and $\sin(m\pi x)$. This means $B_1 = 1$ and $B_2, B_3, \dots = 0$, and the full solution to Eq. (6) is given by

$$u(x, t) = \sin(\pi x) e^{-\pi^2 \alpha^2 t}. \quad (17)$$

C. Solving differential equations with neural networks

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

III. RESULTS AND DISCUSSION

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

IV. CONCLUSION

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.