

	100			200			400			600		
Item	FitResid	χ^2	ANOVA	FitResid	χ^2	ANOVA	FitResid	χ^2	ANOVA	FitResid	χ^2	A
1	-0,18	0,38	0,36	2,10	0,26	0,35	1,84	0,23	0,32	1,99	0,04	
2	-0,85	0,82	0,79	-1,26	0,27	0,19	-3,10	0,02	0,02	-3,85	0,00	
3	0,12	0,36	0,30	-0,95	0,56	0,47	0,78	0,47	0,49	1,10	0,56	
4	-0,62	0,37	0,25	-1,25	0,11	0,03	-0,74	0,41	0,33	-0,83	0,42	
5	0,60	0,47	0,46	0,84	0,28	0,28	0,93	0,22	0,26	1,09	0,35	
6	-1,30	0,49	0,29	-1,83	0,10	0,05	-2,24	0,04	0,02	-3,26	0,01	
7	-0,73	0,78	0,79	0,39	0,52	0,55	-1,59	0,18	0,20	-1,31	0,33	
8	-0,91	0,49	0,34	-0,02	0,60	0,65	-1,02	0,23	0,27	-1,09	0,14	
9	-0,82	0,97	0,94	-1,18	0,42	0,09	-0,62	0,46	0,33	-1,36	0,39	

Karl Bang Christensen, kach@sund.ku.dk

3. oktober 2017

1. Little interest or pleasure in doing things 2. Feeling down, depressed, or hopeless 3. Trouble falling or staying asleep, or sleeping too much 4. Feeling tired or having little energy 5. Poor appetite or overeating 6. Feeling bad about yourself - or that you are a failure or have let yourself or your family down 7. Trouble concentrating on things, such as reading the newspaper or watching television 8. Moving or speaking so slowly that other people could have noticed. Or the opposite being fidgety or restless and moving around a lot more than usual 9. Thoughts that you would be better off dead or of hurting yourself in some way