

1 Relative Frequency

- $f_k(n) = \frac{N_k(n)}{n} \leftarrow$ **Relative Frequency**
 - k is the outcome
 - $N_k(n)$ is the number of times outcome k
- $\lim_{n \rightarrow \infty} f_k(n) = p_k \leftarrow$ **Statistical Regularity**
 - p_k is the probability of event k occurring

1.1 Properties of Relative Frequencies

1. $f_k(n) = \frac{N_k(n)}{n}$
2. $0 \leq N_k(n) \leq n$
3. $0 \leq f_k(n) \leq 1 = \frac{0}{n} \leq \frac{N_k(n)}{n} \leq \frac{n}{n}$
4. $\sum_{k=1}^k f_k(n) = \sum_{k=1}^k \frac{N_k(n)}{n} = \frac{\sum_{k=1}^k N_k(n)}{n} = \frac{n}{n} = 1$
5. $\sum_{k=1}^k f_k(n) = 1$
6. If events A and B are disjoint and event C is " A or B ", then $F_C = F_A(n) + F_B(n)$

2 Set Theory

- A *set* is a collection of objects, denoted by capital letters
- Denote the *universal set*, U ; consisting of all possible objects of interest in a given setting/application
- For any set A , we say that " x is an element of A ", denoted $x \in A$ if object x of the universal set U is contained in A
- We say that " x is not an element of A ", denoted $x \notin A$ if object x of the universal set U is not contained in A
- We say that " A is a subset of B ", denoted $A \subset B$ if every element in A also belongs to B , $x \in A \rightarrow x \in B$
- The *empty set*, \emptyset is defined as the set with no elements
 - The empty set is a subset of every set
- Sets A and B are equal if they contain the same elements. To show this:
 1. Enumerate the elements of each set
 2. Thm: $A = B \iff A \subset B \text{ AND } B \subset A$
- The *union of 2 sets* A, B , denoted $A \cup B$ is defined as the set of outcomes that are either in A , or in B , or both
- The *intersection of 2 sets*, A, B , denoted $A \cap B$ is defined as the set of outcomes in A and B
- The 2 sets A, B are said to be *disjoint or mutually exclusive* if $A \cap B = \emptyset$
- The *complement of a set* A , denoted A^C is defined as the set of elements of U not in A
 - $A^C = \{x \in U | x \notin A\}$
- *Relative complement or difference*, denoted $A - B$, is the set of elements in A that are not in B
 - $A - B = A \cap B^C$
 - $A^C = U - A$

2.1 Properties of Set Operations

Set Operators are:

1. Commutative, Equation (1)

$$\begin{aligned} A \cup B &= B \cup A \\ A \cap B &= B \cap A \end{aligned} \tag{1}$$

2. Associative, Equation (2)

$$\begin{aligned} A \cup (B \cup C) &= (A \cup B) \cup C \\ A \cap (B \cap C) &= (A \cap B) \cap C \end{aligned} \tag{2}$$

3. Distributive, Equation (3)

$$\begin{aligned} A \cup (B \cap C) &= (A \cup B) \cap (A \cup C) \\ A \cap (B \cup C) &= (A \cap B) \cup (A \cap C) \end{aligned} \tag{3}$$

4. Set Operations obey De Morgan's Laws, Equation (4)

$$\begin{aligned} (A \cup B)^C &= A^C \cap B^C \\ (A \cap B)^C &= A^C \cup B^C \end{aligned} \tag{4}$$

Additionally,

Defn 1 (Union of n Sets). The *union of n sets* $\bigcup_{k=1}^n A_k = A_1 \cup A_2 \cup A_3 \cup \dots \cup A_n$ is the set consisting of all elements such that $x \in A_k$ for some $1 \leq k \leq n$.

- All sets need to be empty to make $\bigcup_{k=1}^n A_k = \emptyset$

3 Probability Theory

There are 3 main components to Probability Theory.

1. Set Theory
2. Axioms of Probability
3. Conditional Probability and Independence

3.1 Random Experiments

Defn 2 (Random Experiment). A *random experiment* is an experiment whose outcome varies in an unpredictable fashion when performed under the same conditions.

Defn 3 (Sample Space). A *sample space*, S of a random experiment is the set of all possible experiments.

Defn 4 (Outcome/Sample Point). An *outcome*, or *sample point* of a random experiment is a result that cannot be decomposed into other results.

Defn 5 (Event). An *event* corresponds to a subset of the sample space. We say an event occurs if and only if (iff) the outcome of the experiment is in the subset representing the event.

Defn 6 (Event Classes). An event class \mathcal{F} .