

---

COMMENTARY

---

## Prescription opioids, abuse and public health in Canada: is fentanyl the new centre of the opioid crisis?

Benedikt Fischer<sup>1,2,3\*</sup>, Cayley Russell<sup>1</sup>, Yoko Murphy<sup>1</sup> and Paul Kurdyak<sup>1,2,4</sup>

<sup>1</sup>*Social and Epidemiological Research Department, Centre for Addiction and Mental Health, Toronto, ON, Canada*

<sup>2</sup>*Department of Psychiatry and Institute of Medical Science, University of Toronto, Toronto, ON, Canada*

<sup>3</sup>*Centre for Applied Research in Mental Health and Addiction, Faculty of Health Sciences, Simon Fraser University, Vancouver, BC, Canada*

<sup>4</sup>*Institute for Clinical Evaluative Sciences, Toronto, ON, Canada*

For most of the present century, prescription opioid (PO) misuse and related morbidity (e.g. emergency room and substance use treatment admissions), as well as mortality (e.g. fatal poisonings), have formed a major and unique public health crisis in North America.<sup>1,2</sup> In both the USA and Canada, a large proportion of these PO-related harms had been associated with extended-release formulations of oxycodone (e.g. mainly Oxycontin).<sup>2–4</sup> In Canada, specifically, Oxycontin's inclusion in the province of Ontario's public drug benefit formulary was found to be associated with a significant increase in opioid-related deaths in the period 1991–2004.<sup>5</sup> Approximately one-third of POs dispensed (in defined daily doses (DDD))—and up to half or more of opioid-related poisoning deaths—in Canada were Oxycontin-related until 2012, when the majority of Canadian provinces de-listed the drug from its formularies in order to reduce the PO-related public health toll; subsequent to this intervention, both oxycodone formulation dispensing levels and related poisoning mortality—for example in Ontario—substantially declined.<sup>6,7</sup>

While these oxycodone-focused intervention measures have led to reductions in oxycodone-specific use and harm levels, recent data suggest that improvements on this specific front may have been replaced by increases in harms related to other PO formulations—specifically including fentanyl. Fentanyl is a distinct PO formulation product in that it is a highly potent synthetic

opioid designed for chronic pain treatment; in recent years—for supposed benefits of increased tamper resistance—it has been marketed largely in formulations for transdermal, intranasal and transmucosal administration.<sup>8,9</sup> Consumption levels of fentanyl have more than doubled in Canada in the past decade (from 4428 DDDs in 2003–2005 to 11 007 DDDs in 2011–2013) and currently make up >35% of overall PO consumption (in DDDs); Canadian fentanyl consumption rates are the highest in the world except for a few European countries, yet surpassing the USA where fentanyl prescriptions have also increased by >500% in the period 1997–2007.<sup>1,10</sup>

Notably, fentanyl is one of two strong PO formulations (besides hydromorphone) registering with persistently increasing dispensing levels in Canada following the delisting of Oxycontin in 2012, and which hence—to an overall substantial extent—have substituted for the reductions in consumption of oxycodone formulations.<sup>6,11</sup> Specifically, while oxycodone dispensing levels (in DDDs) in Canada in 2013—after peaking in 2010/11—have returned to 2007/08 levels, fentanyl dispensing levels have continuously increased by about 50% during that period.<sup>11</sup> Yet markedly more important from a public health perspective is that fentanyl-related poisoning deaths have rapidly increased in jurisdictions across Canada. In Ontario, the annual number of fentanyl-related poisoning deaths more than doubled—from 45 in 2008 to 110 in 2013—and thus counteracted about half the reductions in oxycodone-related deaths observed in 2010–2013.<sup>12</sup> In British Columbia, fentanyl-related deaths more than tripled from 15 in 2012, to 51 in 2013 and 84 in 2014<sup>13</sup>; in Alberta, fentanyl-related poisoning deaths increased from 12 in

---

\*Correspondence to: B. Fischer, Department of Social and Epidemiological Research, Centre for Addiction and Mental Health (CAMH), University of Toronto, 33 Russell St., Toronto, Ontario M5S 2S1, Canada. Email: benedikt.fischer@utoronto.ca

2011 to 150 in 2014.<sup>14</sup> Other Canadian provinces—for example New Brunswick and Saskatchewan—have also reported marked increases in fentanyl-related fatalities; a total number of 655 fentanyl-related poisoning deaths have been estimated for Canada for the period 2009–2014 (but may include underestimations).<sup>15</sup> Moreover, other national jurisdictions (including the USA, Nordic countries, Estonia and Australia) on several continents have reported substantial increases in fentanyl-related mortality over the past decade, largely in the context of parallel increases in the medical availability of fentanyl.<sup>16–19</sup>

The circumstances of fentanyl-related overdose deaths have varied by substance (with either fentanyl alone or combined with other opioid/non-opioid drugs) and by route of administration (including intravenous injection).<sup>8,20,21</sup> Fentanyl mortality rates arose unexpectedly despite the drug's supposed 'tamper-resistant' design (including the transdermal matrix formulation) introduced several years ago. It became quickly evident, however, that the supposed abuse barriers could easily be circumvented and that fentanyl formulations—related to a combination of factors involving their distinct physical design, pharmacological characteristics and high psychoactive utility appeal among abusers—carried an amplified high-risk profile for morbidity and/or mortality outcomes.<sup>9,22,23</sup> Specifically, as fentanyl is a highly potent opioid, transdermal patches provide a high-dose load sufficient for several days of therapeutic use, and even previously used or discarded patches still contain sufficient opioid drug amounts to produce substantial euphorogenic effects, or to contribute to opioid poisoning, and the opioid quantity extracted from patches by tampering is likely to be highly variable and unpredictable.<sup>24</sup> An added distinct feature of recent fentanyl-related mortality incidents in Canada is that they have involved both legitimate fentanyl pharmaceutical products as well as illicitly produced fentanyl products from (domestic and international, mainly suspected in China) clandestine sources; the latter have commonly been mixed into and sold as Oxycontin tablets or heroin by street dealers, presumably trying to capitalize on the limited availability but continued high market appeal of these specific opioid types.<sup>25–27</sup> This constitutes a watershed development in the sourcing pathways for PO misuse in Canada, which previously was dominated by diversion from medical sources but now includes a substantial illicit production and distribution element.

While both public health and law enforcement officials in different Canadian jurisdictions are—in ways closely resembling efforts to the then unfolding

'Oxycontin epidemic' just a few years prior—in the process of developing ad hoc counter-strategies for the emerging fentanyl crisis, this case study reinforces several essential lessons and experiences for public health and policy making in the area of PO misuse and harms. The first is that a series of studies have consistently shown that levels of PO-related morbidity and mortality are strongly correlated with levels of PO-drug availability in the population. The described recent increases in fentanyl dispensing in Canada, combined with the high-risk characteristics of the drug, and subsequent rising mortality rates provide another real-life confirmation of these dynamics.<sup>28–30</sup> Second, there is solid evidence in reference to both legal and illegal psycho-pharmaceutical drug markets that reductions of availability in specific opioid drug commodities (e.g. Oxycontin) without corresponding reductions in demand will likely lead to existing similar drugs (e.g. fentanyl) to increasingly being available and (mis-)used as substitutes for the restricted drug.<sup>7,31</sup> Comparable 'substitution effect' phenomena have been observed in, for example the USA (where recent restrictions on POs have been associated with increases in heroin abuse and harms) and Australia (where a 'heroin drought' has been associated with marked shifts to other, e.g. stimulant, drugs in the early 2000s) in the context of reduced availability of or 'shortages' for specific opioid classes.<sup>32–34</sup>

With persistently high PO availability and demand in Canada, PO-related problems like abuse and/or mortality are unlikely to be effectively addressed by only limited focus or interventions targeting one specific drug type (e.g. whether Oxycontin yesterday or fentanyl today) or ad hoc 'emergency response'-type measures. Existing evidence indicates that these kinds of specific interventions mainly result in displacements or shifts of PO-related problems from one drug type to another. These outcomes are undesirable and ambivalent in their public health benefits and mainly constitute a sort of 'Pyrrhic victory' in efforts of improved PO control. Instead, effective and sustained interventions to reduce PO-related harms are required to comprehensively address the known fundamental determinants and drivers of PO abuse and mortality, which primarily ought to include—but not to be limited to—sensible reductions of the extremely high PO dispensing levels in the Canadian population.<sup>30,35</sup>

## CONFLICT OF INTEREST

The authors declared no conflict of interest.

## KEY POINTS

- Extensive spikes in fentanyl abuse and related fatal poisonings have occurred in Canada.
- These increases are following recent reductions in oxycodone-related availability and harms.
- Substitution effects between opioid classes may be at work; a more comprehensive intervention and policy approach is needed.

## ACKNOWLEDGEMENT

Dr Fischer acknowledges funding from Canadian Institutes of Health Research Team Grant SAF-94814 and from a CIHR/PHAC Applied Public Health Research Chair supporting the present work.

## REFERENCES

- Manchikanti L, Helm S, Fellows B, *et al.* Opioid epidemic in the United States. *Pain Physician* 2012; **15**(3): ES9–ES38.
- Fischer B, Argento E. Prescription opioid related misuse, harms, diversion and interventions in Canada: a review. *Pain Physician* 2012; **15**(3): ES191–ES203.
- Paulozzi LJ, Ryan GW. Opioid analgesics and rates of fatal drug poisoning in the United States. *Am J Prev Med* 2006; **31**(6): 506–511.
- Okie S. A flood of opioids, a rising tide of deaths. *N Engl J Med* 2010; **363**(21): 1981–1985.
- Dhalla IA, Mamdani MM, Sivilotti ML, *et al.* Prescribing of opioid analgesics and related mortality before and after the introduction of long-acting oxycodone. *CMAJ* 2009; **181**(12): 891–896.
- Fischer B, Jones W, Rehm J. Trends and changes in prescription opioid analgesic dispensing in Canada 2005–2012: an update with a focus on recent interventions. *BMC Health Serv Res* 2014; **14**(1): 90.
- Fischer B, Keates A. 'Opioid drought', Canadian-style? Potential implications of the 'natural experiment' of delisting Oxycontin in Canada. *Int J Drug Policy* 2012; **23**(6): 495–497.
- Nelson L, Schwaner R. Transdermal fentanyl: pharmacology and toxicology. *J Med Toxicol* 2009 Dec; **5**(4): 230–241.
- Mastropietro DJ, Omidian H. Current approaches in tamper-resistant and abuse-deterrent formulations. *Drug Dev Ind Pharm* 2013 May; **39**(5): 611–624.
- International Narcotics Control Board (INCB). *Narcotic Drugs Estimated World Requirements for 2014—Statistics for 2012*. T.14.XI.2: United Nations, New York, NY., 2014.
- Fischer B, Murphy Y, Jones W, *et al.* Recent developments in prescription opioid-related dispensing and harm indicators in Ontario, Canada. *Pain Physician* 2015; **18**(4): E659–E662.
- Chief Coroner of Ontario. Number of opioid toxicity deaths by drug in Ontario. 2015 [Email]. [05 May 2015].
- Talmazan Y. VDP warn about a spike in fentanyl induced overdose deaths. Global News. 2015. Available at: <http://globalnews.ca/news/1858902/vdp-warn-about-a-spike-in-fentanyl-induced-overdose-deaths/> [accessed 05 May 2015].
- Jones G. *Fentanyl-related Deaths in Alberta*. Chief Medical Examiner: Alberta [Personal Communication]. [05 May 2015], 2011–14.
- Canadian Centre on Substance Abuse (CCSA). *CCENDU Bulletin: Deaths Involving Fentanyl in Canada, 2009–2014*, Aug; Ottawa (ed.): ON, Canadian Centre on Substance Abuse, 2015.
- Roxburgh A, Burns L, Drummer OH, *et al.* Trends in fentanyl prescriptions and fentanyl-related mortality in Australia. *Drug Alcohol Rev* 2013 May; **32**(3): 269–275.
- Simonsen KW, Edvardsen HM, Thelander G, *et al.* Fatal poisoning in drug addicts in the Nordic countries in 2012. *Forensic Sci Int* 2015 Mar; **248**: 172–180.
- Calcaterra S, Glanz J, Binswanger IA. National trends in pharmaceutical opioid related overdose deaths compared to other substance related overdose deaths: 1999–2009. *Drug Alcohol Depend* 2013 Aug 1; **131**(3): 263–270.
- Ojanpera I, Gergov M, Liiv M, *et al.* An epidemic of fatal 3-methylfentanyl poisoning in Estonia. *Int J Legal Med* 2008 Sep; **122**(5): 395–400.
- Martin TL, Woodall KL, McLellan BA. Fentanyl-related deaths in Ontario, Canada: toxicological findings and circumstances of death in 112 cases (2002–2004). *J Anal Toxicol* 2006; **30**: 603–610.
- Hull MJ, Juhascik M, Mazur F, *et al.* Fatalities associated with fentanyl and co-administered cocaine or opiates. *J Forensic Sci* 2007 Nov; **52**(6): 1383–1388.
- Jumbelic MI. Deaths with transdermal fentanyl patches. *Am J Forensic Med Pathol* 2010 Mar; **31**(1): 18–21.
- Firestone M, Goldman B, Fischer B. Fentanyl use among street drug users in Toronto, Canada: behavioural dynamics and public health implications. *Int J Drug Policy* 2009; **20**(1): 90–92.
- Department of Health and Human Services. Fentanyl patch misuse: serious injury, overdose and death. [Internet]. 2012. Available at: <http://docs2.health.vic.gov.au/docs/doc/Fentanyl-patch-misuse:-serious-injury-overdose-and-death> [accessed 25 August 2015].
- Cotter J. Illegal fentanyl linked to more than 100 deaths in Alberta. CTV News. [Internet]. 2015. Available at: <http://www.ctvnews.ca/canada/illegal-fentanyl-linked-to-more-than-100-deaths-in-alberta-last-year-rcmp-1.2286229> [accessed 05 May 2015].
- Carter A. Two more deaths probed by coroner linked to toxic heroin. CBC News. [Internet]. 2014. Available at: <http://www.cbc.ca/news/canada/hamilton/news/two-more-deaths-probed-by-coroner-for-links-to-toxic-heroin-1.2769543> [accessed 05 May 2015].
- Burgmann T. Lethal fentanyl profiting gangs in Western Canada while deaths climb. The Globe and Mail. [Internet]. 2015. Available at: <http://www.theglobeandmail.com/news/british-columbia/lethal-fentanyl-profiting-gangs-in-western-canada-while-deaths-climb/article25866895/> (accessed 25 August 2015).
- Fischer B, Jones W, Urbanoski K, *et al.* Correlations between prescription opioid analgesic dispensing levels and related mortality and morbidity in Ontario, Canada, 2005–2011. *Drug Alcohol Rev* 2013; **33**(1): 19–26.
- King NB, Fraser V, Boikos C, *et al.* Determinants of increased opioid-related mortality in the United States and Canada, 1990–2013: a systematic review. *Am J Public Health* 2014 Aug; **104**(8): 32–42.
- Webster LR, Cochella S, Dasgupta N, *et al.* An analysis of the root causes for opioid-related overdose deaths in the United States. *Pain Med* 2011; **12**(2 Suppl): S26–S35.
- Fishman SM. Commentary in response to Paulozzi *et al.*: prescription drug abuse and safe pain management. *Pharmacoepidemiol Drug Saf* 2006; **15**(9): 628–631.
- Dart RC, Surratt HL, Cicero TJ, *et al.* Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015 Jan 15; **372**(3): 241–248.
- Degenhardt L, Day C, Dietze P, *et al.* Effects of a sustained heroin shortage in three Australian states. *Addiction* 2005 Jul; **100**(7): 908–920.
- Unick GJ, Rosenblum D, Mars S, *et al.* Intertwined epidemics: national demographic trends in hospitalizations for heroin- and opioid-related overdoses, 1993–2009. *PLoS One* 2013; **8**(2): e54496.
- Fischer B, Burnett C, Rehm J. Considerations towards a population health approach to reduce prescription opioid-related harms (with a primary focus on Canada). *Drugs-Educ Prev Policy* 2015; **22**(1): 60–65.