

Machine Learning with Python


MTH786U/P 2020/21

Estimating the height of a person

Mihail Poplavskyi, Queen Mary University of London (QMUL)

What is regression?


Examples:




From “Machine Learning for Hackers” by
Conway & White

What is regression?

Examples:



From “Machine Learning for Hackers” by Conway & White



From Avi Feller et al. 2013

What is regression?

Mathematical formulation:

Given input/output pairs $\{(x_i, y_i)\}_{i=1}^s$ find function f with

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\}$$

Example: linear regression

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\}$$



Example: linear regression

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\} \quad \text{How do we parametrise } f ?$$



Example: linear regression

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\} \quad \text{How do we parametrise } f ?$$

Example:

$$f(x) = w_0 + \sum_{j=1}^d w_j x_j$$

Example: linear regression

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\} \quad \text{How do we parametrise } f ?$$

Example:

$$f(x) = w_0 + \sum_{j=1}^d w_j x_j$$

Affine linear transformation of vector $x = (x_1, \dots, x_d)$ with weights $w \in \mathbb{R}^{d+1}$

Example: linear regression

$$y_i \approx f(x_i) \quad \forall i \in \{1, \dots, s\} \quad \text{How do we parametrise } f ?$$

Example:

$$f(x) = w_0 + \sum_{j=1}^d w_j x_j$$

Affine linear transformation of vector $x = (x_1, \dots, x_d)$ with weights $w \in \mathbb{R}^{d+1}$

Note that $i \neq j$:

$$f(x_i) = w_0 + \sum_{j=1}^d w_j x_{ij}$$

Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle$

with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$



Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle$

with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

(slight abuse of x -notation)



Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle = w^\top x = x^\top w$ with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

(slight abuse of x -notation)



Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle = w^\top x = x^\top w$ with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

How do we choose w such that $y_i \approx f(x_i)$?

(slight abuse of x -notation)



Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle = w^\top x = x^\top w$ with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

How do we choose w such that $y_i \approx f(x_i)$?

(slight abuse of x -notation)

Imagine $s = 3$ and $d = 2$:

$$w_0 + x_{11}w_1 + x_{12}w_2 = y_1$$

$$w_0 + x_{21}w_1 + x_{22}w_2 = y_2$$

$$w_0 + x_{31}w_1 + x_{32}w_2 = y_3$$

Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle = w^\top x = x^\top w$ with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

How do we choose w such that $y_i \approx f(x_i)$? (slight abuse of x -notation)

Imagine $s = 3$ and $d = 2$:

$$\begin{pmatrix} 1 & x_{11} & x_{12} \\ 1 & x_{21} & x_{22} \\ 1 & x_{31} & x_{32} \end{pmatrix} \begin{pmatrix} w_0 \\ w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

Cost function

Notation: $f(x) = w_0 + \sum_{j=1}^d w_j x_j = \langle w, x \rangle = w^\top x = x^\top w$ with $x := \begin{pmatrix} 1 \\ x_1 \\ x_2 \\ \vdots \\ x_d \end{pmatrix} \in \mathbb{R}^{d+1}$

How do we choose w such that $y_i \approx f(x_i)$? (slight abuse of x -notation)

Imagine $s = 3$ and $d = 2$:


$$\begin{pmatrix} 1 & x_{11} & x_{12} \\ 1 & x_{21} & x_{22} \\ 1 & x_{31} & x_{32} \end{pmatrix} \begin{pmatrix} w_0 \\ w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

This system of linear equations has a unique solution if...?


But is it realistic to assume $s = d + 1$?



But is it realistic to assume $s = d + 1$?



But is it realistic to assume $s = d + 1$?



$$s \gg d + 1 = 2$$

Instead we need to find an approximation that is optimal in some sense

Example: Mean-Square Error (MSE)

$$\text{MSE}(w) := \frac{1}{2s} \sum_{i=1}^s |f(x_i) - y_i|^2$$



Instead we need to find an approximation that is optimal in some sense

Example: Mean-Square Error (MSE)

$$\text{MSE}(w) := \frac{1}{2s} \sum_{i=1}^s |f(x_i) - y_i|^2$$

Obtain ‘optimal’ parameters \hat{w} by minimising MSE:

$$\hat{w} = \arg \min_w \text{MSE}(w)$$



Instead we need to find an approximation that is optimal in some sense

Example: Mean-Square Error (MSE)

$$\text{MSE}(w) := \frac{1}{2s} \sum_{i=1}^s |f(x_i) - y_i|^2$$

Obtain ‘optimal’ parameters \hat{w} by minimising MSE:

$$\hat{w} = \arg \min_w \text{MSE}(w)$$

How can we do this?

How do we compute \hat{w} ?

Example: $f(x_i) = w_0 \quad \forall i \in \{1, \dots, s\}$



How do we compute \hat{w} ?

Example: $f(x_i) = w_0 \quad \forall i \in \{1, \dots, s\}$

MSE cost function:
$$\text{MSE}(w_0) := \frac{1}{2s} \sum_{i=1}^s |w_0 - y_i|^2$$



How do we compute \hat{w} ?

Example: $f(x_i) = w_0 \quad \forall i \in \{1, \dots, s\}$

MSE cost function:
$$\text{MSE}(w_0) := \frac{1}{2s} \sum_{i=1}^s |w_0 - y_i|^2$$

We do what we did in school: we compute the derivative and set it to zero:

$$\nabla \text{MSE}(\hat{w}_0) = \text{MSE}'(\hat{w}_0) = \frac{1}{s} \sum_{i=1}^s (\hat{w}_0 - y_i) \stackrel{!}{=} 0$$

How do we compute \hat{w} ?

Example: $f(x_i) = w_0 \quad \forall i \in \{1, \dots, s\}$


MSE cost function:
$$\text{MSE}(w_0) := \frac{1}{2s} \sum_{i=1}^s |w_0 - y_i|^2$$

We do what we did in school: we compute the derivative and set it to zero:


$$\nabla \text{MSE}(\hat{w}_0) = \text{MSE}'(\hat{w}_0) = \frac{1}{s} \sum_{i=1}^s (\hat{w}_0 - y_i) \stackrel{!}{=} 0$$

$$\Rightarrow \hat{w}_0 = \frac{1}{s} \sum_{i=1}^s y_i$$

Example:




Example:




$$\hat{w}_0 \approx 1.1231$$

Example:



Example:



$$\hat{w}_0 \approx 2.4889$$

We will discuss how to compute a better approximation now

