Assignment

Karthikeya hanu prakash kanithi (EE22BTECH11026)

Ouestion: Let {0.13, 0.12, 0.78, 0.51} be a realization of a random sample of size 4 from a population with cumulative distribution function F(.). Consider testing

$$H_0: F = F_0$$
 against $H_1: F \neq F_0$ (1)

where,

$$F_0(x) = \begin{cases} 0 & x < 0 \\ x & 0 \le x < 1 \\ 1 & x \ge 1 \end{cases}$$
 (2)

Let D denote the Kolmogorov-Smirnov test statistic. If P(D > D)0.669) = 0.01 under H_0 and

$$\psi = \begin{cases} 1 & \text{if } H_0 \text{ is accepted at level 0.01} \\ 0 & \text{otherwise} \end{cases}$$
 (3)

then based on the given data, the observed value of $D + \psi$ (rounded off to two decimal places) equals

Solution: Its given that random sample is of size 4, So

$$n = 4 \tag{4}$$

The cdf of the random sample is given as

$$F_0(x) = \begin{cases} 0 & x < 0 \\ x & 0 \le x < 1 \\ 1 & x \ge 1 \end{cases}$$
 (5)

The empirical distribution function(edf) F_n for n independent and identically distributed (i.i.d.) ordered observations X_i is defined as

$$F_n(x) = \frac{\text{no of (elements in the sample } \le x)}{n}$$
 (6)

From (4), (5) and (6), the edf for the given data will be

$$F_n(0.13) = \frac{\text{no of (elements in the sample} \le 0.13)}{4} = \frac{1}{2} \quad (7)$$

$$F_n(0.12) = \frac{\text{no of (elements in the sample} \le 0.12)}{4} = \frac{1}{4} \quad (8)$$

$$F_n(0.12) = \frac{\text{no of (elements in the sample} \le 0.12)}{4} = \frac{1}{4} \quad (8)$$

$$F_n(0.78) = \frac{\text{no of (elements in the sample} \le 0.78)}{4} = 1$$
 (9)

$$F_n(0.78) = \frac{\text{no of (elements in the sample} \le 0.78)}{4} = 1 \quad (9)$$

$$F_n(0.51) = \frac{\text{no of (elements in the sample} \le 0.51)}{4} = \frac{3}{4} \quad (10)$$

The Kolmogorov–Smirnov statistic for a given cdf $F_0(x)$ is

$$D_n = \sup |F_n(x) - F_0(x)| \tag{11}$$

The difference between cdf and edf for the given data will be

$$F_n(0.13) - F_0(0.13) = 0.37$$
 (12)

$$F_n(0.12) - F_0(0.12) = 0.25$$
 (13)

$$F_n(0.78) - F_0(0.78) = 0.22$$
 (14)

$$F_n(0.51) - F_0(0.51) = 0.24$$
 (15)

Then

$$D_n = \sup(0.37, 0.25, 0.22, 0.24) = 0.37$$
 (16)

Given that,

$$P(D > 0.669) = 0.01 \tag{17}$$

Then

$$H_0 = \begin{cases} \text{accepted at level } 0.01 & \text{if } D_n \le 0.669 \\ \text{rejected at level } 0.01 & \text{if } D_n > 0.669 \end{cases}$$
 (18)

From (16) and (18); We can say that H_0 is accepted at level 0.01 and

$$\psi = 1 \tag{19}$$

: the value will be

$$\psi + D_n = 1 + 0.37 = 1.37 \tag{20}$$