

# Lab Course Scientific Computing

## Worksheet 3

distributed: 30.11.2017

due to: 10.12.2017, midnight

oral examination: 12.12.2017 (register for slots in Moodle)

We again examine a similar population equation as in worksheet 2 but with different parameters

$$\dot{p} = 7 \left( 1 - \frac{p}{10} \right) \cdot p \quad (1)$$

and with a different initial condition

$$p(0) = 20. \quad (2)$$

The analytical solution is given by

$$p(t) = \frac{200}{20 - 10e^{-7t}}.$$

- a) Plot the function  $p(t)$  in a graph.
- b) Reuse the Euler method and the method of Heun implemented in worksheet 2 to compute approximate solutions for equation (1) with initial conditions (2), end time  $t_{end} = 5$ , and  $\delta t = \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}$ .

Plot your solutions in one graph per method (together with the given solution from a)). Plot the function in the range  $t \in [0, 5]$  and  $p \in [0, 20]$ .

- c) Implement the following implicit numerical methods with variable stepsize  $\delta t$  and end time  $t_{end}$

- 1) implicit Euler method,
- 2) second order Adams-Moulton method.

for the solution of the initial value problem

$$\dot{y} = f(y), \quad y(0) = y_0$$

as a function of the right hand side  $f$ , the first derivative of the right hand side with respect to  $y$ , initial value  $y_0$ , the stepsize  $\delta t$  and the end time  $t_{end}$ . The output of the function is a vector containing all computed approximate values for  $y$ . Use an accuracy limit of  $10^{-4}$  for the Newton iteration in each time step. Stick to the signatures of the functions from worksheet 2 as far as possible.

**Hint:** Examine if the equation to be solved in each time step of these methods is solvable. If necessary, implement a stopping criterion to prevent the Newton solver from trying to solve unsolvable equations. In these cases, stop the time stepping with the method and the time step concerned and do not consider the associated approximations of  $y$  in your further examinations.

- d) For both methods implemented, compute – as far as possible – approximate solutions for equation (1) with initial conditions (2) and with time steps  $\delta t = \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}$ . Plot your solutions in one graph per method (together with the given solution from a)).
- e) To be able to handle also the cases for which you did not find a solution in d), implement the following linearised versions of the Adams Moulton method for equation (1)

- 1) linearisation 1:

$$y^{(n+1)} = y^{(n)} + \frac{\delta t}{2} \left( 7 \cdot \left( 1 - \frac{y^{(n)}}{10} \right) \cdot y^{(n)} + 7 \cdot \left( 1 - \frac{y^{(n+1)}}{10} \right) \cdot y^{(n)} \right),$$

- 2) linearisation 2:

$$y^{(n+1)} = y^{(n)} + \frac{\delta t}{2} \left( 7 \cdot \left( 1 - \frac{y^{(n)}}{10} \right) \cdot y^{(n)} + 7 \cdot \left( 1 - \frac{y^{(n)}}{10} \right) \cdot y^{(n+1)} \right),$$

for the solution of the initial value problem

$$\dot{y} = 7 \cdot \left(1 - \frac{y}{10}\right) \cdot y, \quad y(0) = y_0$$

as a function of the initial value  $y_0$ , the stepsize  $\delta t$  and the end time  $t_{end}$ . The output of the function is a vector containing all computed approximate values for  $y$ .

- f)** For both methods implemented, compute approximate solutions for equation (1) with initial conditions (2) and with time steps  $\delta t = \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}$ . Plot your solutions in one graph per method (together with the given solution from **a)**).
- g)** Compare the results of the implicit methods to those computed with the explicit methods:

Compute the approximation error

$$E = \sqrt{\frac{\delta t}{5} \sum_k (p_k - p_{k,exact})^2}$$

for each case in **b)** and **d)**, where  $p_k$  denotes the approximation of  $p(\delta t \cdot k)$ ,  $p_{exact,k}$  the exact values of  $p$  at  $t = \delta t \cdot k$ .

Collect the results in the tabulars below. Also write all information needed in the tabulars to the Matlab console in a readable way.

- h)** For each of the following methods, determine the factor by which the error is reduced if the step size  $\delta t$  is halved.
- 1) explicit Euler method,
  - 2) method of Heun,
  - 3) implicit Euler method,
  - 4) Adams Moulton method,
  - 5) Adams moulton method – linearisation 1,
  - 6) Adams moulton method – linearisation 2

Write down the results in the tabular below.

- i) In addition to accuracy, we examine an additional aspect of 'quality' of a method: the *stability*. Descriptively spoken, stability denotes the applicability of a method for varying parameters, whereas at least results similar to the exact/correct solution have to be achieved (In particular, unphysical oscillations should not occur). With this heuristic definition, decide for which of the used values for  $\delta t$  each of the four examined methods is stable (in the case of our problem).

Mark stable cases by a cross in the last tabular. Try to find a simple criterion to determine whether a solution is stable or not and write the result to the Matlab console.

| explicit Euler |               |               |               |                |                |
|----------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$     | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error          |               |               |               |                |                |
| error red.     |               |               |               |                |                |

| Heun       |               |               |               |                |                |
|------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$ | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error      |               |               |               |                |                |
| error red. |               |               |               |                |                |

| implicit Euler |               |               |               |                |                |
|----------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$     | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error          |               |               |               |                |                |
| error red.     |               |               |               |                |                |

| Adams Moulton |               |               |               |                |                |
|---------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$    | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error         |               |               |               |                |                |
| error red.    |               |               |               |                |                |

| Adams Moulton – linearisation 1 |               |               |               |                |                |
|---------------------------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$                      | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error                           |               |               |               |                |                |
| error red.                      |               |               |               |                |                |

| Adams Moulton – linearisation 2 |               |               |               |                |                |
|---------------------------------|---------------|---------------|---------------|----------------|----------------|
| $\delta t$                      | $\frac{1}{2}$ | $\frac{1}{4}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{32}$ |
| error                           |               |               |               |                |                |
| error red.                      |               |               |               |                |                |

| Stable cases              |                |      |                |               |                  |                  |
|---------------------------|----------------|------|----------------|---------------|------------------|------------------|
|                           | explicit Euler | Heun | implicit Euler | Adams-Moulton | Adams Moulton l1 | Adams Moulton l2 |
| $\delta t = \frac{1}{2}$  |                |      |                |               |                  |                  |
| $\delta t = \frac{1}{4}$  |                |      |                |               |                  |                  |
| $\delta t = \frac{1}{8}$  |                |      |                |               |                  |                  |
| $\delta t = \frac{1}{16}$ |                |      |                |               |                  |                  |
| $\delta t = \frac{1}{32}$ |                |      |                |               |                  |                  |

### Questions:

1) For which integer  $q$  can you conclude that the accuracy of the

- a) explicit Euler method,
- b) method of Heun,
- c) implicit Euler method,
- d) Adams Moulton method,
- e) Adams moulton method – linearisation 1,
- f) Adams moulton method – linearisation 2

behaves like  $O(\delta t^q)$ ?

2) In the lecture, we saw that both the implicit Euler and the Adams Moulton method are unconditionally stable and, thus, give us stable solutions for every choice of  $\delta t$ . For the example of this worksheet we stated in **c)** and **d)** that the resulting equation for each timestep is sometimes not solvable and, thus, the method cannot be applied for certain  $\delta t$ . Can you explain this apparent discrepancy?

3) Can you give a reason why the linearisation 1 of the Adams Moulton method works better?

4) Which type of methods (explicit/implicit) would you choose for

- 1) the initial value problem from worksheet 2,
- 2) the initial value problem (1,2) from this worksheet? Give a reason why you choose a certain type of methods and why you do not choose the other type in each case?