



Linear Discriminant Analysis - Introduction

- Linear Discriminant Analysis or Normal Discriminant Analysis or Discriminant Function Analysis is a dimensionality reduction technique that is commonly used for supervised classification problems.
- It is used to project the features in higher dimension space into a lower dimension space.
- Suppose we have two sets of data points belonging to two different classes that we want to classify.
- When the data points are plotted on the 2D plane, there's no straight line that can separate the two classes of the data points completely.
- Here, Linear Discriminant Analysis uses both the axes (X and Y) to create a new axis and projects data onto a new axis in a way to maximize the separation of the two categories and hence, reducing the 2D graph into a 1D graph.



Linear Discriminant Analysis - Introduction

- Two criteria are used by LDA to create a new axis:
 - Maximize the distance between means of the two classes.
 - Minimize the variation within each class.



1. Compute the class means of dependent variable
2. Derive the covariance matrix of the class variable
3. Compute the within class — scatter matrix
($S_1 + S_2$)
4. Compute the between class scatter matrix
5. Compute the Eigen values and eigen vectors
from the within class and between class scatter
matrix

$$\mu_1 = \frac{1}{N_1} \sum_{x \in \mathcal{X}_1} x$$

$$S_1 = \sum_{x \in \mathcal{X}_1} (x - \mu_1)(x - \mu_1)^T$$

$$S_w = S_1 + S_2$$

$$S_B = (\mu_1 - \mu_2)(\mu_1 - \mu_2)^T$$

$$S_w^{-1} S_B w = \lambda w$$

6. Sort the values of eigen values and select
the top k values
7. Find the eigen vectors corresponds to the
top k eigen vectors
8. Obtain the LDA by taking the dot product of
eigen vectors and original data


$$\left(S_w^{-1} S_B - \lambda I \right) \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = 0$$

Linear Discriminant Analysis - Solved Example

- Compute the Linear Discriminant projection for the following two dimensional dataset.

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$



Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

The classes mean are :

$$\mu_1 = \frac{1}{N_1} \sum_{x \in \omega_1} x = \frac{1}{5} \left[\begin{pmatrix} 4 \\ 2 \end{pmatrix} + \begin{pmatrix} 2 \\ 4 \end{pmatrix} + \begin{pmatrix} 2 \\ 3 \end{pmatrix} + \begin{pmatrix} 3 \\ 6 \end{pmatrix} + \begin{pmatrix} 4 \\ 4 \end{pmatrix} \right] = \begin{pmatrix} 3 \\ 3.8 \end{pmatrix}$$

$$\mu_2 = \frac{1}{N_2} \sum_{x \in \omega_2} x = \frac{1}{5} \left[\begin{pmatrix} 9 \\ 10 \end{pmatrix} + \begin{pmatrix} 6 \\ 8 \end{pmatrix} + \begin{pmatrix} 9 \\ 5 \end{pmatrix} + \begin{pmatrix} 8 \\ 7 \end{pmatrix} + \begin{pmatrix} 10 \\ 8 \end{pmatrix} \right] = \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix}$$

Linear Discriminant Analysis - Solved Example

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

Covariance matrix of the first class:

$$S_1 = \sum_{x \in \omega_1} \frac{(x - \mu_1)(x - \mu_1)^T}{N-1} = \frac{1}{4} \left[\begin{pmatrix} 4 \\ 2 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right] \left[\begin{pmatrix} 4 \\ 2 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right]^T + \begin{pmatrix} 2 \\ 4 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \left[\begin{pmatrix} 2 \\ 4 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right]^T + \begin{pmatrix} 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \left[\begin{pmatrix} 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right]^T + \begin{pmatrix} 3 \\ 6 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \left[\begin{pmatrix} 3 \\ 6 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right]^T + \begin{pmatrix} 4 \\ 4 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \left[\begin{pmatrix} 4 \\ 4 \end{pmatrix} - \begin{pmatrix} 3 \\ 3.8 \end{pmatrix} \right]^T \right]$$

$$= \begin{pmatrix} 1 & -0.25 \\ -0.25 & 2.2 \end{pmatrix}$$

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

Covariance matrix of the second class:

$$S_2 = \sum_{x \in \omega_2} \frac{(x - \mu_2)(x - \mu_2)^T}{N-1} = \frac{1}{4} \left[\begin{pmatrix} 9 \\ 10 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right] \left[\begin{pmatrix} 9 \\ 10 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T + \begin{pmatrix} 6 \\ 8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \left[\begin{pmatrix} 6 \\ 8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T + \begin{pmatrix} 9 \\ 5 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \left[\begin{pmatrix} 9 \\ 5 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T + \begin{pmatrix} 8 \\ 7 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \left[\begin{pmatrix} 8 \\ 7 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T + \begin{pmatrix} 10 \\ 8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \left[\begin{pmatrix} 10 \\ 8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T \right]$$

$$= \begin{pmatrix} 2.3 & -0.05 \\ -0.05 & 3.3 \end{pmatrix}$$

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

Within-class scatter matrix:

$$S_w = S_1 + S_2 = \begin{pmatrix} 1 & -0.25 \\ -0.25 & 2.2 \end{pmatrix} + \begin{pmatrix} 2.3 & -0.05 \\ -0.05 & 3.3 \end{pmatrix}$$

$$= \begin{pmatrix} 3.3 & -0.3 \\ -0.3 & 5.5 \end{pmatrix}$$

Linear Discriminant Analysis - Solved Example

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

$$\begin{aligned} S_B &= (\mu_1 - \mu_2)(\mu_1 - \mu_2)^T \\ &= \left[\begin{pmatrix} 3 \\ 3.8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right] \left[\begin{pmatrix} 3 \\ 3.8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right]^T \\ &= \begin{pmatrix} -5.4 \\ -3.8 \end{pmatrix} \begin{pmatrix} -5.4 & -3.8 \end{pmatrix} \\ &= \begin{pmatrix} 29.16 & 20.52 \\ 20.52 & 14.44 \end{pmatrix} \end{aligned}$$

- Find Eigen Values

$$\begin{aligned} S_W^{-1} S_B w &= \lambda w \\ \Rightarrow |S_W^{-1} S_B - \lambda I| &= 0 \\ \Rightarrow \begin{vmatrix} 3.3 & -0.3 \\ -0.3 & 5.5 \end{vmatrix}^{-1} \begin{pmatrix} 29.16 & 20.52 \\ 20.52 & 14.44 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} &= 0 \\ \Rightarrow \begin{pmatrix} 0.3045 & 0.0166 \\ 0.0166 & 0.1827 \end{pmatrix} \begin{pmatrix} 29.16 & 20.52 \\ 20.52 & 14.44 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} &= 0 \\ \Rightarrow \begin{pmatrix} 9.2213 - \lambda & 6.489 \\ 4.2339 & 2.9794 - \lambda \end{pmatrix} & \\ = (9.2213 - \lambda)(2.9794 - \lambda) - 6.489 \times 4.2339 &= 0 \\ \Rightarrow \lambda^2 - 12.2007\lambda = 0 \Rightarrow \lambda(\lambda - 12.2007) &= 0 \\ \Rightarrow \lambda_1 = 0, \lambda_2 = 12.2007 & \end{aligned}$$

- Find Eigen Vector

$$\left(S_W^{-1} S_B - \lambda I \right) \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = 0$$

$$w_1 = \begin{pmatrix} -0.5755 \\ 0.8178 \end{pmatrix}$$

$$w_2 = \begin{pmatrix} 0.9088 \\ 0.4173 \end{pmatrix} = w^*$$

- Find Eigen Vector

$$\left(\begin{matrix} S_W^{-1} S_B - \lambda I \\ - \quad - \quad - \quad - \end{matrix} \right) \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = 0$$

$$\underline{w_1} = \begin{pmatrix} -0.5755 \\ 0.8178 \end{pmatrix}$$

$$\underline{w_2} = \begin{pmatrix} 0.9088 \\ 0.4173 \end{pmatrix} = w^*$$

Or directly;

$$\begin{aligned} \underline{w^*} &= S_W^{-1} (\mu_1 - \mu_2) = \begin{pmatrix} 3.3 & -0.3 \\ -0.3 & 5.5 \end{pmatrix}^{-1} \left[\begin{pmatrix} 3 \\ 3.8 \end{pmatrix} - \begin{pmatrix} 8.4 \\ 7.6 \end{pmatrix} \right] \\ &= \begin{pmatrix} 0.3045 & 0.0166 \\ 0.0166 & 0.1827 \end{pmatrix} \begin{pmatrix} -5.4 \\ -3.8 \end{pmatrix} \\ &= \begin{pmatrix} 0.9088 \\ 0.4173 \end{pmatrix} \end{aligned}$$

- Obtain the LDA by taking the dot product of eigen vectors and original data

$$w_2 = \begin{pmatrix} 0.9088 \\ 0.4173 \end{pmatrix} = w^*$$

Samples for class ω_1 : $\mathbf{X}_1 = (x_1, x_2) = \{(4,2), (2,4), (2,3), (3,6), (4,4)\}$

Sample for class ω_2 : $\mathbf{X}_2 = (x_1, x_2) = \{(9,10), (6,8), (9,5), (8,7), (10,8)\}$

X1	4	2	2	3	4	9	6	9	8	10
X2	2	4	3	6	4	10	8	5	7	8
1 st LD	4.46	3.48	3.06	5.2	5.3	12.35	8.8	10.2	10.19	12.42

