

Date: September 28, 2026
To: Editor-in-Chief, *Computers & Fluids*
Elsevier Publishing

Subject: Submission of Original Research Article for Publication in *Computers & Fluids*

Dear Editor-in-Chief and Editorial Board,

We are pleased to submit our original research manuscript entitled:

“A High-Resolution Finite-Difference Framework for Steady Continuation and Finite-Window Unsteady Dynamics up to $Re = 100,000$ in 2D Lid-Driven Cavity Flows”

for consideration as a regular research paper in *Computers & Fluids*.

Significance and Novelty of the Work:

The incompressible 2D lid-driven cavity flow remains the quintessential proving ground for computational fluid dynamics. However, established benchmarks have largely stagnated at $Re \leq 10,000$ (e.g., Ghia et al., 1982) or $Re \leq 25,000$ (Erturk et al., 2005) due to numerical divergence stemming from the lid corner singularities and escalating cell Péclet numbers ($Pe \gg 2$).

In this manuscript, we push the resolution and Reynolds frontiers of the standard finite-difference streamfunction-vorticity (ψ - ω) formulation through several key technical advances:

1. **Exact Spectral Poisson Acceleration:** We integrate an exact Discrete Sine Transform (DST-I) Poisson solver operating at optimal $\mathcal{O}(N^2 \log N)$ complexity, eliminating iterative convergence errors, pressure-velocity decoupling, and artificial numerical dissipation ($\nu_{\text{art}} = 0$).
2. **Asymptotic Batchelor Core Verification at $Re = 100,000$:** We compute stationary continuation solutions up to $Re = 100,000$ on ultra-fine grids (513×513) and a high-resolution grid (1025×1025 , 1,050,625 nodes). We observe core vorticity behavior qualitatively consistent with Batchelor’s (1956) asymptotic theorem: the primary vortex center migrates toward the geometric mid-plane ($x_c \rightarrow 0.50000$), while core vorticity stabilizes to a plateau value of $\omega_0 \approx -2.54$.
3. **Interior Hybrid Differencing & Numerical Dissipation Mapping:** At $Re = 100,000$ ($Pe = 195.3$), an interior hybrid differencing scheme regularizes corner singularity dispersion with added numerical viscosity tensor $\boldsymbol{\nu}_{\text{art}} = \text{diag}(\nu_{\text{art},x}, \nu_{\text{art},y})$, yielding directional mean ratio $\bar{\nu}_{\text{art}}/\nu = 1.485$ at the geometric cavity center (0.5,0.5) and 48.93 near the lid on 513×513 (dropping to 0.74 and 24.46 on 1025×1025), with pure second-order central differencing strictly preserved everywhere for all $Re \leq 50,000$.
4. **Unsteady Dynamics on 513×513 Meshes:** Through physical time-accurate marching ($t = 16.0$ – 22.0 c.t.u., $St \equiv fL/U$), we systematically resolve the primary periodic shedding mode ($Re = 10,000$, $St = 0.6249 \pm 0.0416$), modulated multi-frequency dynamics ($Re = 15,000$, $St = 0.5364 \pm 0.0278$), folded multi-frequency orbits ($Re = 25,000$, $St = 0.0953 \pm 0.0281$), filtered-boundary vortex shedding ($Re = 50,000$, $St = 0.2581 \pm 0.0300$), and hybrid filtered-boundary shedding at $Re = 100,000$ ($St = 0.1908 \pm 0.0379$), reporting discrete Fourier bin half-width resolution ($\pm \Delta f/2$).

5. **Commitment to Open Science and Reproducibility:** We openly release thirty (30) verified benchmark full 2D coordinate/flow field datasets (23 steady continuation and 7 unsteady records), phase portrait scripts, and automated visualization tools under the MIT License with permanent Zenodo DOIs.

Suggested Potential Reviewers:

1. **Prof. Ercan Erturk** (Bahçeşehir University, Istanbul, Turkey) — World authority on high-Reynolds lid-driven cavity benchmarking and steady Navier–Stokes solutions up to $Re = 25,000$.
2. **Prof. Olivier Botella** (Université de Lorraine, LEMTA, France) — Renowned specialist in high-order compact finite difference schemes and incompressible cavity benchmarks.
3. **Prof. Chang Shu** (National University of Singapore, Singapore) — Pioneer of differential quadrature and lattice Boltzmann / FDM formulations for high-Reynolds cavity flows.
4. **Prof. Jie Shen** (Eastern Institute of Technology, China / Purdue University, USA) — Expert in spectral methods, projection schemes, and Navier–Stokes stability.

Declarations:

This manuscript is an original work, has not been published previously, and is not under simultaneous consideration for publication elsewhere. All authors have read and approved the final manuscript. There are no competing financial or non-financial interests to declare.

Sincerely yours,

Kartikeya Gangwar

Department of Mathematics

University of Delhi, Delhi, 110007, India

Email: kartikeyasingh525@protonmail.com

ORCID: <https://orcid.org/0009-0009-1973-7532>