

Documents - Vensopro x

172.18.0.17:8080/VENSOPRO/Document/ViewDoc?id=8BF5AFA5-2F49-4E06-8461-B975BA4261B6&q=55342c

DMS View Document

7 of 55 Automatic Zoom

COMPLAINTS
Fever since 5 days
Headache and bodyache since 5 days

HISTORY OF PRESENTING ILLNESS
patient developed fever since 5 days which was sudden in onset, high grade type, intermittent in nature, gradually progressive. Associated with chills and rigors.
patient c/o headache since 5 days, throbbing type, in the frontal region.
patient also complains of generalized weakness.
no h/o abdominal pain /vomitting/loose stools/constipation
no h/o cough / breathlessness
no h/o burning micturition / hematuria
patient gives a h/o irregular menstrual cycles.
patient went to a local hospital where routine blood investigations were done.she was found to be NS 1 antigen postive (dengue fever)

PAST HISTORY
Patient is a K/C/O multinodular colloid goitre.
No history of Diabetes Mellitus, Hypertension, Tuberculosis, Asthma or IHD

FAMILY HISTORY
No history of Diabetes Mellitus, Hypertension, Tuberculosis, Asthma or IHD in the Family

PERSONAL HISTORY
Diet: Mixed. Appetite: Good. Sleep: Adequate. Bowel and Bladder: Regular. No substance abuse.

GENERAL EXAMINATION
Patient is consious and cooperative, well oriented to time, place and person.
Moderately built and nourished. No pallor, icterus, cyanosis, clubbing, lymphadenopathy or oedema
PULSE:80bpm
BP :120/80mmhg
RR:20/min

Phone : 0834-2238124, 2238090 Fax : 2436661, 2437402

Start Documents - Vensopro... Microsoft Excel - 201- 10:59 AM

DMS View Document

172.18.0.17:8080/VENSOPRO/Document/ViewDoc?id=8BF5AFA5-2F49-4E06-8461-B975BA4261B6&q=55342c

REQUIREMENTS OF GENERAL INVESTIGATIONS

TEST NAME	TESTED DATE	UNIT	REFERENCE RANGE
HEMATOGRAM			
WBC Count			
RBC Count			
Hb %			
Hct %			
MCV			
MCH			
MCHC			
RDW			
URINE			
The patient was advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection. The patient was also advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection. The patient was also advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection.			
OTHER			
The patient was advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection. The patient was also advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection. The patient was also advised to collect a clean midstream urine sample in a sterile container. The sample should be delivered to the laboratory within 2 hours of collection.			

