

Pressure-aware signed third-order heat lift: exact coefficient interfaces




Two independent exact Fourier producers | $q = \exp(-s)$ | coefficientwise evidence | no PDE simulation

A Exact pressure-aware interface

four-site target $\hat{h}_* = (1, 2, 0)$ | exact q -polynomials

$$N = P \nabla \cdot (u \otimes u), \quad \chi_s = P_s(u \odot N) - v_s \odot N_s$$




$$K = [[-2, 1, 0], [1, 0, 0], [0, 0, 0]], \quad \|K\|_F = \sqrt{6}$$

SIGN-PAIR DIFFERENCE
 $\hat{\chi}_s(u) - \hat{\chi}_s(-u) = 2(q^3 - q^5) K$

exact target coefficient; distinct from the Germano stress source

B Four-site mode decomposition

$\hat{h}_* = (1, 2, 0)$ | exact matrices and certified bottom-scale orders




exact orders: s^2 vs $s \Rightarrow$ coefficientwise absorption costs at least s^{-1}

C Six-site zero-mode witness

same output coefficient $\hat{h} = 0$ | exact cancellation / survival

$$u = (6\sin y - 4\sin(x + y), 4\sin x + 4\sin(x + y), 0)$$




nonzero for $0 < s < \infty$

INTERPRETATION BOUNDARY
 coefficientwise witness only — not a whole-field information collision

D Selected quartic remainder

κ_{112} at $(0, 2, 0)$ | exact formula; line is not a fit



Parabolic lift: $\|\Delta \hat{\chi}\|_F = 2\sqrt{6}L(e^{-3\theta} - e^{-5\theta})$; **selected quartic tangent** $= 2iLe^{-2\theta}(1 - e^{-2\theta})^2$.

Exact two-path finite Fourier certificate | coefficientwise / selected-coefficient scope | no simulation, fit, closure theorem, or global regularity result | NOT CLAY