EVEN UNIMODULAR IDEAL LATTICES OF RANK 88 OVER $\mathbb{Q}(\zeta_{115})$

IVAN GAUVIN

ABSTRACT. Let $L = \mathbb{Q}(\zeta_{115})$, of degree 88. We study the set S of classes of unimodular ideal lattices $(I, \mathrm{Tr}_{L/\mathbb{Q}}(\alpha x\bar{y}))$ over L; these lattices are even. We show that S is in bijection with the class group of L, which is cyclic of order $h^-(115) = 44\,697\,909 = 3 \cdot 331 \cdot 45\,013$, and that S splits into 508 463 Galois orbits. Sakamoto's two extremal lattices, L_0 and L_S, occupy the principal orbit and the unique non-trivial 3-torsion orbit. Extending an argument of Quebbemann for automorphisms of prime order, we prove that two ideal lattices over L with minimum at least 8 are isometric if and only if their classes are conjugate under $\mathrm{Gal}(L/\mathbb{Q})$. We extend to non-principal ideals a gluing into two forms of rank 44, and we derive from it, by a Parseval-type inequality, a finite extremality criterion. The orbits of eight prime ideals are rejected by explicit vectors of norm 6. For two prime ideals of degree 2, P_{139} and P_{599}, the lattices they carry are extremal: their minimum 8 is established by a computer-assisted proof, whose execution model is made explicit and whose computations were replicated on a second computer. Without assuming GRH, they are isometric neither to L_0 nor to L_S, nor, under this computer-assisted minimum, to each other. To our knowledge, after a literature check as of October 13, 2026, these are two new examples of extremal lattices of rank 88.

This is an English translation of the original French text, which follows it in the same document. Each statement carries its status: [proved], [under GRH], [computer-assisted], [observed] or [open] (definitions in §1.3). Caveats are gathered in §§5.4 and 7.2.

1. INTRODUCTION

1.1. **Context.** An even unimodular lattice of rank n has minimum at most $2\lfloor n/24 \rfloor + 2$; it is called *extremal* if its minimum attains this value. This is the bound of Mallows, Odlyzko and Sloane, recalled in the introduction of [Sak26]. In rank 88, an extremal lattice is therefore an even unimodular lattice of minimum 8.

According to the introduction of [Sak26], no extremal lattice of rank greater than 80 was known before that preprint; for rank 80, it cites [BN98, Wat12]. Sakamoto [Sak26] constructs two non-isometric extremal lattices of rank 88, denoted there by L and L_0; we denote the first by L_S and the second by L_0. They are Hermitian tensor products over $\mathbb{Q}(\sqrt{-23})$: $L_S = E \otimes J$ and $L_0 = E \otimes J_0$, where E is the ring of integers of $\mathbb{Q}(\sqrt{-23}, \zeta_5)$, $J = (47, \zeta_{23} - 21)$ is a non-principal ideal of $\mathbb{Q}(\zeta_{23})$ and J_0 is a principal ideal of $\mathbb{Q}(\zeta_{23})$. The first version of [Sak26], dated September 14, 2026, contains only L_S; L_0 appears in the second version, dated September 27, 2026. By [Sak26, Rem. 3.11], these two lattices are ideal lattices over $\mathbb{Q}(\zeta_{115})$, carried by $J\mathcal{O}_L$ and by the principal ideal $J_0\mathcal{O}_L$. By [Sak26, Thm. A.10], their automorphism groups have orders 1 457 280 and 2 914 560. Remark 5.6 of [Sak26] states that the tensor construction yields only these two isometry classes, and that it is not a classification of the extremal lattices of rank 88.

Date: October 15, 2026; revised November 7, 2026 (version 1.5 of the repository).

2020 *Mathematics Subject Classification.* Primary 11H06; Secondary 11H56, 11R18, 11R29, 11E12.

Key words and phrases. ideal lattices, even unimodular lattices, extremal lattices, cyclotomic fields, class group, certified enumeration.

The lattice $H = (\mathcal{O}_L, \mathrm{Tr}_{L/\mathbb{Q}}(\beta_0 z\bar{z}))$, for an explicit $\beta_0 \in \mathbb{Q}(\zeta_{115})^+$ (see the notation below), is isometric to L_0 (§2.5). The lattice H is therefore not new: Sakamoto has priority. The extremality of L_0 is that of [Sak26, Prop. 5.5]; we have not re-proved it.

The present note starts from the following observation: L_0 and L_S are two elements of a finite set S, namely the set of classes of unimodular ideal lattices over $L = \mathbb{Q}(\zeta_{115})$. It describes S and looks for those of its elements that have minimum 8.

1.2. **Content of the note.**

(1) *The set S* (§2). Every unimodular ideal lattice over L is even. The set S is in bijection with $\mathrm{Cl}(L)$, which is cyclic of order $44\,697\,909 = 3 \cdot 331 \cdot 45\,013$. It has 508 463 orbits under $\mathrm{Gal}(L/\mathbb{Q})$, of sizes 1, 2, 22, 44 and 88. The lattice L_0 is the principal orbit and L_S the unique non-trivial 3-torsion orbit.

(2) *Isometry theorem* (§3). Two ideal lattices of minimum ≥ 8 over L are isometric if and only if their classes in S are conjugate under $\mathrm{Gal}(L/\mathbb{Q})$.

(3) *Gluing and criterion* (§4). For an integral ideal I coprime to $\bar{I}$, the lattice (I, b_α) embeds into a sum of two forms of rank 44. A Parseval lemma gives an extremality criterion: it suffices to examine the projections of energy at most 1057/200.

(4) *Certified enumeration* (§5). A program applies this criterion. A non-omission proof describes its scope, under an explicit model of the machine.

(5) *Results* (§6). Eight prime ideals give non-extremal lattices, by an explicit vector of norm 6. Two prime ideals of degree 2, P_{139} and P_{599}, carry two new extremal lattices, to our knowledge as of October 13, 2026, isometric neither to each other nor to Sakamoto's lattices; the minimum is established by a computer-assisted proof (Theorem 6.11). Four other ideals are partial survivors of a filter, with no conclusion.

The minimum 8 of X_{139} and X_{599} is established by a computer-assisted proof (Theorem 6.11): a program, whose non-omission is proved in an explicit model of the machine (§5), went through both sides of the criterion of §4 without finding a short vector. Each sweep was executed twice, on two different computers, under two operating systems and two versions of Java, and the receipts of the two executions are identical byte for byte. Without assuming GRH, the two lattices are isometric neither to L_0 nor to L_S; their mutual non-isometry follows, without GRH, from the non-conjugacy of their classes, under this computer-assisted minimum.

1.3. **Statuses.** We use five statuses.

- [proved]: complete proof in the text, or exact replayable identity (integer witness, certificate), if needed together with a published result cited by name.
- [under GRH]: depends on a class group or unit group computation by `bnfinit` of PARI/GP.
- [computer-assisted]: proved under named hypotheses bearing on an execution of a program (machine, hardware, integrity of the logs). These hypotheses are H1 to H4 (§5.4).
- [observed]: program output without proof value.
- [open]: not established.

Statement	Status
$h^-(115) = 44\,697\,909$; $\mathrm{Cl}(L)$ cyclic; $\mathrm{Cl}^+(L^+) \cong \mathbb{Z}/2$ (Prop. 2.2, 2.3)	proved (analytic formula in exact arithmetic; $h^+ = 1$ by [Mil14])
Evenness; bijection $S \leftrightarrow \mathrm{Cl}(L)$ (Prop. 2.1, 2.3)	proved
Galois action; 508 463 orbits (Prop. 2.5, Table 1)	proved (Stickelberger relation, exact norms, Burnside's formula)
$H \cong L_0$	proved (integer certificate); Sakamoto has priority
Extremality of L_0 and L_S	[Sak26, Prop. 5.5 and §5]; not re-proved here
Position of L_0 and L_S in S (Prop. 2.7)	proved, modulo the ideal description and Lemma A.9 of [Sak26]
Lemmas 3.1 and 3.2; Thm. 3.3; Cor. 3.5	proved (internal proof, reviewed by a second assistant, see the final declaration)
Prop. 4.1; Lemmas 4.2, 4.3, 4.5; Thm. 4.6	proved
Thm. 5.1 (non-omission of the program)	proved under the machine model M1–M3; reviewed by a second assistant, within a limited scope

Statement	Status
Non-extremality for P_{461}, P_{691}, P_{1151}, P_{1381}, P_{1471}, P_{829}, P_{1289}, P_{1979} (Table 2)	proved, without GRH (exact witnesses of norm 6)
Location of these classes in $\mathrm{Cl}(L)$	3-component: without GRH; 331- and 45 013-components: under GRH
P_{139}, P_{599}: even unimodular lattices; vector of norm 8	proved (exact checks)
P_{139}: no non-zero vector of norm ≤ 6	minimum ≥ 8 computer-assisted (H1–H3); two complete executions on two computers, identical receipts
P_{599}: no non-zero vector of norm ≤ 6	minimum ≥ 8 computer-assisted (H1–H3); two complete executions on two computers, identical receipts
$[P_{139}]^3 \neq 1$; the order of $[P_{599}]$ is divisible by 331 (Prop. 6.6, 6.7)	proved, without GRH
Neither P_{139} nor P_{599} is isometric to L_0 or to L_S (Prop. 6.9)	proved, modulo [Sak26] and Thm. 3.3; does not depend on the computer-assisted statement
Extremality of X_{139} and X_{599}; non-isometry to each other, to L_0 and to L_S (Thm. 6.11)	minimum: computer-assisted proof (M1, M2, H1–H3); non-conjugacy: proved, without GRH (Rem. 6.10); non-isometry between X_{139} and X_{599}: under this computer-assisted minimum; to L_0 and L_S: independent of it
Exact label of the orbits of P_{139} and P_{599}	under GRH (details in §6.4)
P_{3359}, P_{4049}, P_{4969}, P_{5659} (Table 3)	observed: partial survivors of a filter; no conclusion
Number of orbits of minimum 8 in S	open; between 4 and 508 463: the orbits of L_0 and L_S (minimum by [Sak26]) and those of X_{139} and X_{599} (Thm. 6.11), pairwise distinct

1.4. **What is taken from the literature and what is specific to this work.** *Taken from the literature.* The lattices L_0 and L_S, their extremality, their description as ideal lattices over $\mathbb{Q}(\zeta_{115})$ and their automorphism groups are due to Sakamoto [Sak26]. The automatic evenness of a unimodular ideal lattice when L/L^+ is unramified at the finite places is classical [Bay84, Bay99]. The parametrization of ideal lattices by ideal classes and by a quotient of units is that of Nebe [Neb16, Rem. 4.4]; Nebe [Neb16, Lemma 4.3, Thm. 4.5] also uses the direction "conjugate, hence isometric" of Theorem 3.3. The equality $h(\mathbb{Q}(\zeta_{115})^+) = 1$ is due to Miller [Mil14]. The analytic formula for the relative class number and the Stickelberger relation are classical [Was97]. The gluing of lattices is classical [CS99, Ch. 4]. Blichfeldt's bound [Bli29] is used for the isometry theorem. This theorem extends an argument of Quebbemann [Que81, §3]: for an automorphism of prime order p in rank n, with $p \leq n < 2(p-1)$, isometry of the underlying lattices amounts to Galois conjugacy of the arithmetic data, by a Sylow argument (Satz 3.2 b), p. 165; see also Kor. 3.3 and Lemma 3.4, p. 166). The idea of bounding a minimum from below by a Parseval identity on the coordinates of $\mathbb{Z}[\zeta_p]$ is also in Quebbemann (p. 168, for the Leech lattice). Bayer-Fluckiger [Bay84, Cor. 3.3] gives an *upper bound*: when h^- is odd, the number of isometry classes of definite unimodular lattices with an automorphism of characteristic polynomial Φ_m is at most the number of orbits of the Galois group on a group of ideal classes; she asks (p. 526) for the exact number.

Specific to this work. The computation of $h^-(115)$ as used here, the description of S and of its orbits (§2); the extension of the isometry theorem to the composite conductor $m = 115$, where the rank condition of [Que81] is replaced by a condition on the minimum (Blichfeldt bounds), and whose proof does not use Kronecker's theorem (§3); the gluing for a non-principal ideal, the form of the Parseval lemma adapted to this gluing (quintuples with zero sum) and the finite criterion derived from it, with its threshold 1057/200 (§4); the certifying program and its non-omission proof (§5); the rejections, the two candidates and their classes (§6). The integer isometry certificate $H \cong L_0$ is also part of this work; the lattice itself is Sakamoto's.

Prior-art search. A literature search was carried out in October 2026: arXiv, the record and the text of [Sak26] (version 2 of September 27, 2026), the computation repository of its author, the catalogue of Nebe and Sloane (explicitly incomplete), keyword searches in academic sources. The only other explicit object of rank 88 that was found has a Gram matrix whose diagonal contains 6: it has a vector of norm 6 and is not extremal. No other explicit example of an extremal lattice of rank 88 was identified. To our knowledge, as of October 13, 2026, X_{139} and X_{599} are therefore two new examples; this is the result of a dated search, not a proof of priority over unpublished or unindexed work. This search had not identified Quebbemann's article [Que81], which is cited since the revision of November 7, 2026.

Outline. Section 2 describes S. Section 3 proves the isometry theorem. Section 4 establishes the gluing, the Parseval lemma and the criterion. Section 5 presents the certified enumeration and its hypotheses. Section 6 gives the results. Section 7 gathers the open questions, the methods and the declaration on the use of artificial intelligence.

Notation. We set $\zeta = \zeta_{115}$, $L = \mathbb{Q}(\zeta)$, of degree 88, $L^+ = \mathbb{Q}(c)$ with $c = \zeta + \zeta^{-1}$, and $\mathcal{O}^+ = \mathbb{Z}[c]$, the ring of integers of L^+. We denote by $x \mapsto \bar{x}$ complex conjugation, $\mathrm{N} = \mathrm{N}_{L/L^+}$, $\mathrm{Tr}^+ = \mathrm{Tr}_{L^+/\mathbb{Q}}$, $\mathfrak{D}$ the different of L^+, U_L and U^+ the unit groups of L and of L^+, $U^+_{\gg 0}$ the group of totally positive units of L^+, and μ_L the group of roots of unity of L, of order 230. For a prime to 115, σ_a is the automorphism $\zeta \mapsto \zeta^a$. We set $e = \zeta - \zeta^{-1}$; we have $e^2 = c^2 - 4$. We write $T = \mathbb{Q}(\zeta_{23})$ and $F = \mathbb{Q}(\sqrt{-23})$, subfields of L.

An *ideal lattice* is a pair $X = (I, \alpha)$, where I is a fractional ideal of L and $\alpha \in L^+$ is totally positive ($\alpha \gg 0$), endowed with the form $b_\alpha(x, y) = \mathrm{Tr}_{L/\mathbb{Q}}(\alpha x \bar{y})$. The *norm* of a

vector z is $b_\alpha(z,z)$. We fix a totally positive element $\beta_0 \in L^+$ such that $\beta_0\mathfrak{D} = \mathcal{O}^+$; it is written out in full in the data accompanying the note (§7.2), where these two properties are checked exactly. We set $H = (\mathcal{O}_L, \beta_0)$.

2. The set of unimodular ideal lattices over $\mathbb{Q}(\zeta_{115})$

2.1. Evenness and unimodularity.

Proposition 2.1. (a) *The extension L/L^+ is unramified at the finite places, e is a unit, and $\mathfrak{D}_L = \mathfrak{D}\,\mathcal{O}_L$.*
(b) *The lattice (I,α) is unimodular if and only if $\alpha\,\mathrm{N}(I)\,\mathfrak{D} = \mathcal{O}^+$.*
(c) *Every unimodular ideal lattice over L is even.*
[proved]

Proof. (a) We have $\mathcal{O}_L = \mathcal{O}^+[\zeta]$, and ζ is a root of $X^2 - cX + 1$; the relative different is therefore generated by $e = \zeta - \zeta^{-1} = -\zeta^{-1}(1-\zeta^2)$. Since ζ^2 is a primitive root of unity of order 115, which is not a prime power, $1-\zeta^2$ is a unit. By transitivity, $\mathfrak{D}_L = \mathfrak{D}\,\mathcal{O}_L$. (b) The dual of I for b_α is $(\alpha\bar{I}\mathfrak{D}_L)^{-1}$. Hence (I,α) is unimodular if and only if $\alpha I\bar{I}\mathfrak{D}_L = \mathcal{O}_L$, that is, $\alpha\,\mathrm{N}(I)\mathfrak{D}\,\mathcal{O}_L = \mathcal{O}_L$; we conclude by the injectivity of the extension of ideals from L^+ to L. (c) For $z \in I$, we have $z\bar{z} \in I\bar{I} \cap L^+ = \mathrm{N}(I)$, hence $\alpha z\bar{z} \in \mathfrak{D}^{-1}$ and $b_\alpha(z,z) = 2\,\mathrm{Tr}^+(\alpha z\bar{z}) \in 2\mathbb{Z}$. □

Let S be the set of unimodular ideal lattices (I,α) over L, modulo the relation $(I,\alpha) \sim (xI, \alpha/(x\bar{x}))$, $x \in L^\times$. The map $z \mapsto xz$ is an isometry from (I,α) onto $(xI, \alpha/(x\bar{x}))$. The group $\mathrm{Gal}(L/\mathbb{Q})$ acts on S by $\sigma_a(I,\alpha) = (\sigma_a I, \sigma_a\alpha)$, and $z \mapsto \sigma_a(z)$ is an isometry from (I,α) onto $\sigma_a(I,\alpha)$. Two elements of S that are conjugate under $\mathrm{Gal}(L/\mathbb{Q})$ are therefore isometric.

2.2. Class number.

Proposition 2.2. *We have $h^-(115) = 44\,697\,909 = 3 \cdot 331 \cdot 45\,013$ and $h(L^+) = 1$. Consequently $h(L) = 44\,697\,909$ is squarefree, and*

$$\mathrm{Cl}(L) \cong \mathbb{Z}/3 \times \mathbb{Z}/331 \times \mathbb{Z}/45\,013$$

is cyclic. [proved]

Proof. The analytic formula for the relative class number reads $h^- = Q\,w \prod_\chi (-\tfrac{1}{2}B_{1,\chi^*})$, where the product runs over the odd characters modulo 115, χ^* is the associated primitive character, $Q = 2$ is the Hasse index and $w = 230$ ([Was97, Thm. 4.17]). The product was evaluated in exact arithmetic in $\mathbb{Q}(\zeta_{44})$ and equals 44 697 909. A second, independent computation, in integers and rationals, without any class group computation, gives the same value (§7.2). The same program recovers $h^-(87) = 1\,536$, $h^-(65) = 64$ and $h^-(85) = 6\,205$. The equality $h(L^+) = 1$ is due to Miller [Mil14, Thm. 1.1], without GRH. An abelian group of squarefree order is cyclic. □

2.3. **Structure of S.** Let $\mathrm{N}^+\colon \mathrm{Cl}(L) \to \mathrm{Cl}^+(L^+)$, $[I] \mapsto [\mathrm{N}(I)]$, with values in the narrow class group of L^+.

Proposition 2.3. (a) *The map N^+ is trivial, $\mathrm{Cl}^+(L^+) \cong \mathbb{Z}/2$, $[U^+_{\gg 0} : (U^+)^2] = 2$ and $\mathrm{N}(U_L) = U^+_{\gg 0}$.*
(b) *The map $(I,\alpha) \mapsto [I]$ is a bijection from S onto $\mathrm{Cl}(L)$, compatible with the action of $\mathrm{Gal}(L/\mathbb{Q})$. In particular $|S| = 44\,697\,909$: each ideal class carries exactly one unimodular ideal lattice, up to equivalence.*
(c) *For an ideal I, let $\nu \gg 0$ be a generator of $\mathrm{N}(I)$. Then $(I, \beta_0/\nu)$ represents the element of S with image $[I]$.*
[proved]

Proof. (a) The exact sequence of signs gives $|\operatorname{Cl}^+(L^+)| = h(L^+)\,[U^+_{\gg 0} : (U^+)^2]$, which is a power of 2 since $h(L^+) = 1$. Since $\operatorname{Cl}(L)$ has odd order (Proposition 2.2), its image under N^+ is trivial. On the other hand, L/L^+ is unramified at the finite places and ramified at the infinite places; L is therefore contained in the narrow Hilbert class field of L^+, and Artin reciprocity gives $\operatorname{Cl}^+(L^+)/\mathrm{N}^+(\operatorname{Cl}(L)) \cong \operatorname{Gal}(L/L^+)$. Hence $|\operatorname{Cl}^+(L^+)| = 2$ and $[U^+_{\gg 0} : (U^+)^2] = 2$.

The element $1-\zeta$ is a unit, and $-\zeta = (1-\zeta)/(1-\bar\zeta)$ generates μ_L: it is not a square in μ_L. Hence $1-\zeta \notin \mu_L U^+$, the Hasse index equals 2 [Was97, Thm. 4.12, Cor. 4.13], and $\mathrm{N}(U_L) = (U^+)^2 \cup (2-c)(U^+)^2$ with $2-c = \mathrm{N}(1-\zeta)$. If $2-c$ were a square u^2 in U^+, the unit $(1-\zeta)/u$ would have absolute value 1 at every place, hence would be a root of unity, and $1-\zeta$ would lie in $\mu_L U^+$. Thus $[\mathrm{N}(U_L) : (U^+)^2] = 2$. Since $\mathrm{N}(U_L) \subset U^+_{\gg 0}$, we have $\mathrm{N}(U_L) = U^+_{\gg 0}$.

(b) By Proposition 2.1 (b), $(I,\alpha) \in S$ if and only if $\mathrm{N}(I)\mathfrak{D} = (\alpha^{-1})$ with $\alpha^{-1} \gg 0$. The image of S is therefore the set of classes $[I]$ such that $\mathrm{N}^+([I]) = [\mathfrak{D}]^{-1}$ in $\operatorname{Cl}^+(L^+)$. Now $[\mathfrak{D}]$ is trivial in $\operatorname{Cl}^+(L^+)$, since $\beta_0\mathfrak{D} = \mathcal{O}^+$ with $\beta_0 \gg 0$, and N^+ is trivial. The image is therefore the whole of $\operatorname{Cl}(L)$. For fixed I, the suitable α form a class $\alpha_0 U^+_{\gg 0}$, and $(I,\alpha) \sim (I,\alpha')$ if and only if there exists $x \in U_L$ with $\alpha' = \alpha/(x\bar x)$. The fibers are therefore principal homogeneous spaces under $U^+_{\gg 0}/\mathrm{N}(U_L)$, a group which is trivial by (a). The compatibility with the Galois action is immediate.

(c) The generator ν exists because N^+ is trivial, and $(\beta_0/\nu)\,\mathrm{N}(I)\mathfrak{D} = \beta_0\mathfrak{D} = \mathcal{O}^+$. □

Remark 2.4. The parametrization by ideal classes and by $U^+_{\gg 0}/\mathrm{N}(U_L)$ is that of Remark 4.4 of Nebe [Neb16]. For $m = 115$, the fiber over each class has only one element; over the principal class, it is H. The Gram matrices constructed for this note are checked exactly: they are integral, even and of determinant 1.

2.4. **Galois action and orbits.**

Proposition 2.5. *For $p \in \{3, 331, 45\,013\}$, let Cl_p be the subgroup of order p of $\operatorname{Cl}(L)$. The group $\operatorname{Gal}(L/\mathbb{Q})$ acts on $\mathrm{Cl}_p \cong \mathbb{F}_p$ through an odd character χ_p with values in $\mathbb{F}_p^\times$, whose order and kernel are as follows; the last column gives the size of the orbits on $\mathrm{Cl}_p \smallsetminus \{0\}$.*

p	*order*	*kernel of χ_p*	*fixed field of the kernel*	*orbit size*
3	2	*kernel of the Legendre symbol modulo 23, of order 44*	$F = \mathbb{Q}(\sqrt{-23})$	2
331	22	$\{1, 22, 24, 68\} = \langle\sigma_{22}\rangle$	K_{22}, *of degree 22*	22
45 013	44	$\{1, 91\} = \langle\sigma_{91}\rangle$	$K_{44} = \mathbb{Q}(\zeta_5, \zeta_{23} + \zeta_{23}^{-1})$	44

Complex conjugation acts by -1 on $\operatorname{Cl}(L)$. [proved]

Proof. Since p does not divide 88, the Sylow p-subgroup of $\operatorname{Cl}(L)$ decomposes into isotypic components under $\operatorname{Gal}(L/\mathbb{Q})$; here it has order p, hence equals a single component, associated with a character χ_p. Complex conjugation acts by -1: for every class $[I]$, $[I][\bar I]$ is the image in $\operatorname{Cl}(L)$ of $[\mathrm{N}(I)] \in \operatorname{Cl}(L^+) = 1$. The character χ_p is therefore odd. It is identified by the Stickelberger relation and by the analytic formula [Was97, Thm. 4.17 and Thm. 6.10]: for each of the six rational conjugacy classes of odd characters modulo 115, the norm of $\frac{1}{2}B_{1,\chi^*}$ was computed exactly. The Euler factors of the three imprimitive classes have norms 1, 2 and 2; they introduce no factor 3, 331 or 45 013. For each p, exactly one class has a norm divisible by p, and the valuation there equals 1. These exact computations were checked independently (§7.2). The automorphisms σ_{24} and σ_{91} have order 2: σ_{24} conjugates ζ_5 and fixes ζ_{23}, σ_{91} fixes ζ_5 and conjugates ζ_{23}. □

Let us call the *support* of a class the set of the p for which its component in Cl_p is non-zero. The stabilizer of a class is the intersection of the kernels of the χ_p, p in the

support. All classes with the same support therefore have orbits of the same size, and the following count results; it was also recounted with Burnside's formula.

TABLE 1. Orbits of $\mathrm{Gal}(L/\mathbb{Q})$ in $S \cong \mathrm{Cl}(L)$, by support. Status: proved.

support	classes	orbit size	orbits	stabilizer
$\emptyset$ (principal class)	1	1	1	$\mathrm{Gal}(L/\mathbb{Q})$
$\{3\}$	2	2	1	of order 44
$\{331\}$	330	22	15	$\langle\sigma_{22}\rangle$, of order 4
$\{3, 331\}$	660	44	15	$\langle\sigma_{24}\rangle$
$\{45\,013\}$	45 012	44	1 023	$\langle\sigma_{91}\rangle$
$\{3, 45\,013\}$	90 024	88	1 023	trivial
$\{331, 45\,013\}$	14 853 960	88	168 795	trivial
$\{3, 331, 45\,013\}$	29 707 920	88	337 590	trivial
total	44 697 909		508 463	

There are therefore 1 053 orbits with non-trivial stabilizer outside the first two rows. The number 508 463 is that of all the orbits of S; the number of those that consist of lattices of minimum 8 is not known.

Remark 2.6 (normalizer of $\langle\zeta\rangle$)**.** For $X \in S$, the automorphisms of X that normalize the group $\langle\zeta\rangle$ of multiplications are the maps $z \mapsto \lambda\sigma_a(z)$ with $\sigma_a \in \mathrm{Stab}(X)$, and λ is determined by a up to μ_L. This group therefore has order $230 \cdot |\mathrm{Stab}(X)|$, that is, 20 240, 10 120, 920, 460 or 230 for the orbits of size 1, 2, 22, 44 or 88. Indeed, such an automorphism ψ satisfies $\psi\zeta\psi^{-1} = \zeta^a$ for some a prime to 115; as in step 3 of the proof of Theorem 3.3, $\psi(z) = \lambda\sigma_a(z)$ with $\lambda\sigma_a(I) = I$ and $\alpha\lambda\bar{\lambda} = \sigma_a(\alpha)$, hence $\sigma_a \in \mathrm{Stab}(X)$. Conversely, for $\sigma_a \in \mathrm{Stab}(X)$, the relation $\sigma_a(I, \alpha) \sim (I, \alpha)$ provides such a λ. Two choices of λ for the same a differ by a unit u such that $u\bar{u} = 1$, of absolute value 1 at every place: by Kronecker's theorem, $u \in \mu_L$. [proved]

2.5. **Position of Sakamoto's two lattices.** The isometry $H \cong L_0$ is established by a certificate: an integer matrix P of size 88, of determinant -1, satisfies $PGP^{\mathsf{T}} = G_H$, where G is the Gram matrix of L_0 published in the data repository of [Sak26] (§7.2) and $G_H = (\mathrm{Tr}_{L/\mathbb{Q}}(\beta_0\zeta^{i-j}))$. A second integer identity, $QG_{\mathrm{art}}Q^{\mathsf{T}} = G$ with $\det Q = 1$, shows that G is the Gram matrix of the construction of [Sak26, §5.4] in an explicit basis. These two identities were verified in exact arithmetic by two programs, then by a third one, written independently (§7.2). Status of $H \cong L_0$: [proved], without GRH. This certificate says nothing about the minimum.

Proposition 2.7. (a) *The class of L_0 in S is the principal class, whose orbit has size* 1.
(b) *The class of L_S in S has support $\{3\}$. It forms the unique orbit with this support, of size* 2.

[proved, modulo [Sak26, Rem. 3.11 and Lemma A.9]]

Proof. (a) follows from $H \cong L_0$ and $H = (\mathcal{O}_L, \beta_0)$. (b) By [Sak26, Rem. 3.11], L_S is the ideal lattice of $J\mathcal{O}_L$. This ideal is not principal: if it were, L_S and L_0 would have the same Steinitz class as modules over the ring of integers of F, which contradicts [Sak26, Lemma A.9]. Its class is 3-torsion, because J is an ideal of $T = \mathbb{Q}(\zeta_{23})$ and $\mathrm{Cl}(T) \cong \mathbb{Z}/3$ [Sak26]. It therefore has order 3. The subgroup of order 3 of $\mathrm{Cl}(L)$ has two non-zero elements, which are exchanged by complex conjugation (Proposition 2.5). □

Remark 2.8. For every automorphism t of L_S with characteristic polynomial Φ_{115}, L_S is not a free $\mathbb{Z}[t]$-module. Indeed, such a free module would be isometric to a lattice $(\mathcal{O}_L, \gamma)$, hence to $H \cong L_0$ by Proposition 2.3, which contradicts [Sak26, Thm. A.10]. [proved, modulo [Sak26, Thm. A.10]]

3. The isometry theorem

For an integer n, we denote by γ_n Hermite's constant. Blichfeldt's bound [Bli29] reads $\gamma_n \le B_n = \frac{2}{\pi}\,\Gamma\big(2+\frac{n}{2}\big)^{2/n}$. We have $B_{22} \approx 3.9174$ and $B_{44} \approx 6.6471$. We only use the strict rational upper bounds

$$\gamma_{22} < \frac{126}{31}, \qquad \gamma_{44} < \frac{210}{31}, \tag{3.1}$$

which follow from $\pi > 31/10$, $12! < (63/10)^{11}$ and $23! < (21/2)^{22}$.

Lemma 3.1 (Sylow 23)**.** *Let X be an even unimodular lattice of rank 88 and minimum at least 8.*

(a) *Every automorphism of order 23 of X has no non-zero fixed point.*
(b) *The Sylow 23-subgroups of $\mathrm{Aut}(X)$ have order at most 23. In particular, the subgroups of order 23 of $\mathrm{Aut}(X)$ are conjugate.*

[proved]

Proof. (a) Let $t \in \mathrm{Aut}(X)$ be of order 23. Its characteristic polynomial is $\Phi_{23}^z(x-1)^{88-22z}$ with $1 \le z \le 4$. Let F_t be the fixed lattice of t, of rank $88-22z$, and $F_t^\perp$ its orthogonal complement, of rank $22z$; $F_t^\perp$ is a $\mathbb{Z}[\zeta_{23}]$-module of rank z. Since X is unimodular, the discriminant groups of F_t and of $F_t^\perp$ are isomorphic; this group is annihilated by $1-t$, hence isomorphic to $\mathbb{F}_{23}^s$ with $s \le z$ (a classical result on the type of an automorphism of prime order, see [Neb13]). The lattices F_t and $F_t^\perp$ have minimum at least 8 and determinant 23^s.

- If $z=1$, $F_t^\perp$ has rank 22 and determinant 23^s, $s \le 1$. Its Hermite invariant is at least $8/23^{1/22}$, which exceeds $126/31$ since $(8\cdot 31/126)^{22} > 23$. This contradicts (3.1).
- If $z=2$, F_t has rank 44 and determinant 23^s, $s \le 2$. Its Hermite invariant is at least $8/23^{2/44}$, which exceeds $210/31$ since $(8\cdot 31/210)^{44} > 23^2$. Contradiction.
- If $z=3$, F_t has rank 22 and determinant 23^s, $s \le 3$. Its Hermite invariant is at least $8/23^{3/22}$, which exceeds $126/31$ since $(8\cdot 31/126)^{22} > 23^3$. Contradiction.

Hence $z=4$: t has no non-zero fixed point.

(b) A 23-group of order greater than 23 contains a subgroup of order 23^2, cyclic or isomorphic to $C_{23}\times C_{23}$. An element of order 529 is impossible in $\mathrm{Aut}(X)$, since $\varphi(529) = 506 > 88$. A subgroup $C_{23}\times C_{23}$ would act on $X\otimes\mathbb{Q}$ without trivial constituent, by (a); each of its irreducible constituents would have dimension 22, with a kernel of order 23, and a non-trivial element of this kernel would have non-zero fixed points, which contradicts (a). The subgroups of order 23 are therefore Sylow subgroups, and they are conjugate. □

Lemma 3.2 (Sylow 5 in the centralizer)**.** *Let $X=(I,\alpha)$ be an ideal lattice over L, and let $P_X = \langle\zeta^5\rangle \subset \mathrm{Aut}(X)$ be the group of order 23 of multiplications by the powers of ζ^5. Let C be the centralizer of P_X in $\mathrm{Aut}(X)$. The subgroups of order 5 of C are conjugate in C.* [proved]

Proof. Every $g \in C$ commutes with multiplication by ζ^5, which generates $T=\mathbb{Q}(\zeta_{23})$: g is T-linear on $X\otimes\mathbb{Q} = L \cong T^4$. The polynomial Φ_5 is irreducible over T, since $\mathbb{Q}(\zeta_5)\cap T = \mathbb{Q}$; hence $T[x]/(x^5-1) \cong T\times T(\zeta_5)$. If $g\in C$ has order 5, the $T[g]$-module T^4 contains a factor $T(\zeta_5)$, of dimension 4: it is equal to it, and g has no non-zero fixed point. A subgroup $C_5\times C_5$ of C would act on T^4 through non-trivial irreducible constituents, of dimension 4 and with a kernel of order 5; there is room for only one constituent, and the action would not be faithful. An element of order 25 of C would require a constituent of dimension $[T(\zeta_{25}):T] = 20 > 4$. The Sylow 5-subgroups of the finite group C therefore have order 5, and the subgroups of order 5 of C are conjugate in C. □

Theorem 3.3 (isometry)**.** *Let $X_1 = (I_1, \alpha_1)$ and $X_2 = (I_2, \alpha_2)$ be two unimodular ideal lattices over $L = \mathbb{Q}(\zeta_{115})$, of minimum at least 8. Then $X_1 \cong X_2$ if and only if the classes of X_1 and of X_2 in S are conjugate under* $\mathrm{Gal}(L/\mathbb{Q})$*.* [proved]

Proof. The "if" direction follows from the isometries $z \mapsto xz$ and $z \mapsto \sigma_a(z)$. Conversely, let $\psi\colon X_1 \to X_2$ be an isometry, and let $\tau_j \in \mathrm{Aut}(X_j)$ be multiplication by ζ. The lattices X_1 and X_2 are even (Proposition 2.1), so Lemma 3.1 applies to X_2.

Step 1. The group $\psi\langle\tau_1^5\rangle\psi^{-1}$ is a subgroup of order 23 of $\mathrm{Aut}(X_2)$. By Lemma 3.1, it is conjugate to $P_2 = \langle\tau_2^5\rangle$. After composing ψ on the left with an automorphism of X_2, we may assume $\psi\langle\tau_1^5\rangle\psi^{-1} = P_2$.

Step 2. The group $\psi\langle\tau_1^{23}\rangle\psi^{-1}$ has order 5 and commutes with P_2, since τ_1^{23} commutes with τ_1^5. By Lemma 3.2 applied to X_2, there exists g in the centralizer of P_2 such that $g\psi\langle\tau_1^{23}\rangle\psi^{-1}g^{-1} = \langle\tau_2^{23}\rangle$. Set $\psi' = g\psi$. Since g centralizes P_2, we still have $\psi'\langle\tau_1^5\rangle\psi'^{-1} = P_2$.

Step 3. The group $\langle\tau_j\rangle$ is generated by τ_j^5 and τ_j^{23}. Hence $\psi'\langle\tau_1\rangle\psi'^{-1} = \langle\tau_2\rangle$, and $\psi'\tau_1\psi'^{-1} = \tau_2^a$ for some a prime to 115. The $\mathbb{Q}$-linear extension of ψ' to L satisfies $\psi'(\zeta z) = \zeta^a\psi'(z)$; it is therefore σ_a-semilinear: $\psi'(z) = \lambda\sigma_a(z)$ with $\lambda = \psi'(1) \in L^\times$. It follows that $\lambda\sigma_a(I_1) = I_2$ and $\mathrm{Tr}_{L/\mathbb{Q}}(\alpha_2\lambda\bar{\lambda}\,\sigma_a(z\bar{w})) = \mathrm{Tr}_{L/\mathbb{Q}}(\alpha_1 z\bar{w})$ for all z, w, hence $\alpha_2\lambda\bar{\lambda} = \sigma_a(\alpha_1)$. Thus $(I_2, \alpha_2) \sim \sigma_a(I_1, \alpha_1)$. □

Remark 3.4. For an automorphism of prime order p in rank $n < 2(p-1)$, the Sylow argument and the semi-linearity of Step 3 are those of Quebbemann [Que81, Satz 3.2 b), p. 165]; there, the condition $n < 2(p-1)$ plays the role that the minimum 8 plays here (Lemma 3.1). Here $115 = 5 \cdot 23$ and the two Sylow arguments suffice: the proof uses neither Kronecker's theorem nor the class group. The rational comparisons of (3.1), the details on the 23-groups and the alignment of the two Sylow subgroups were reviewed by a second assistant. The proof has not been reviewed by a specialist.

Corollary 3.5. *The isometry classes of even unimodular lattices of rank* 88 *and minimum* 8 *that have an automorphism with characteristic polynomial Φ_{115} or Φ_{230} are in bijection with the orbits of* $\mathrm{Gal}(L/\mathbb{Q})$ *in S consisting of lattices of minimum* 8*.* [proved]

Proof. Since $\Phi_{230}(x) = \Phi_{115}(-x)$, it suffices to treat Φ_{115}. Let (X, t) be given, with t of characteristic polynomial Φ_{115}; X is a torsion-free module of rank 1 over $\mathbb{Z}[t] \cong \mathcal{O}_L$, and the choice of a non-zero vector identifies it with a fractional ideal I. A positive definite symmetric bilinear form on L invariant under ζ is of the form b_α, with $\alpha \in L^+$ totally positive. Thus $X \cong (I, \alpha) \in S$. Changing the vector does not change the class in S, and another choice of t gives a conjugate class, by Theorem 3.3. The resulting map is injective by the "if" direction, and surjective since multiplication by ζ is an automorphism with characteristic polynomial Φ_{115} of every (I, α). □

Remark 3.6. The lattices L_0 and L_S occupy two orbits of minimum 8, according to [Sak26]. By Theorem 3.3, a lattice of minimum 8 whose class has a support containing 331 or 45 013 would be isometric neither to L_0 nor to L_S. Theorem 6.11, whose proof of the minimum is computer-assisted, raises the lower bound to four: the number of orbits of minimum 8 lies between 4 and 508 463; its exact value remains *open*. The other integers m such that $\varphi(m) = 88$ are 89, 178, 184, 230 and 276; 230 gives back 115, and the others are not treated here.

4. Gluing, Parseval lemma and extremality criterion

A direct enumeration of the vectors of norm at most 6 in rank 88 is out of reach: an estimate of the number of nodes of the Fincke–Pohst enumeration [FP85], made on L_0, gives $2.4 \cdot 10^{22}$ [observed]. LLL reduction does not find short vectors either: on L_0, of minimum 8, it stops at norm 16 [observed]. This section reduces the question to enumerations in rank 44.

4.1. **Gluing for a non-principal ideal.** In this section, I is an integral ideal of $\mathcal{O}_L$ such that $I+\bar{I}=\mathcal{O}_L$. This is the case of a prime ideal $I\neq\bar{I}$. We denote by $q=[\mathcal{O}_L:I]$ its absolute norm, by $\nu\gg 0$ a generator of $\mathrm{N}(I)$ (Proposition 2.3), and we set $\alpha=\beta_0/\nu$. We have $I\cap\bar{I}=I\bar{I}=\nu\mathcal{O}_L$ and $\mathrm{N}_{L^+/\mathbb{Q}}(\nu)=q$. We define two quadratic forms on L^+:

$$\Lambda(a)=\mathrm{Tr}^+(\alpha a^2),\qquad \Lambda'(b)=\mathrm{Tr}^+(\alpha(4-c^2)b^2).$$

Proposition 4.1 (gluing)**.** (a) *The map* $z\mapsto(a,b)=\big(z+\bar{z},\ (z-\bar{z})/e\big)$ *is an isometry from* (I,b_α) *onto*

$$M_I=\big\{(a,b)\in\mathcal{O}^+\times\mathcal{O}^+:\ (a+eb)/2\in I\big\},\qquad \text{endowed with}\quad N(a,b)=\tfrac{1}{2}(\Lambda(a)+\Lambda'(b)).$$

(b) *In the basis* $(c^i)_{0\le i<44}$ *of* $\mathcal{O}^+$*, the forms* Λ *and* Λ' *are rational, positive definite, of determinant* $1/q$*. The index of* M_I *in* $\mathcal{O}^+\times\mathcal{O}^+$ *is* $2^{44}q$*.*

(c) *For fixed* a*, the* b *such that* $(a,b)\in M_I$ *form a class modulo* $2\nu\mathcal{O}^+$*, or the empty set. The lattice* $2\nu\mathcal{O}^+$*, endowed with* Λ'*, has determinant* $4^{44}q$*. The same statement holds with the roles of* a *and* b *exchanged, with* Λ *on* $2\nu\mathcal{O}^+$*.*

[proved]

Proof. (a) For $z\in\mathcal{O}_L$, $a=z+\bar{z}$ lies in $\mathcal{O}_L\cap L^+=\mathcal{O}^+$. The quotient $b=(z-\bar{z})/e$ is real, and it is integral because e is a unit (Proposition 2.1). We have $z=(a+eb)/2$, so the map is injective and its image on I is M_I. Since $\bar{e}=-e$ and $e^2=c^2-4$,

$$z\bar{z}=\tfrac{1}{4}(a+eb)(a-eb)=\tfrac{1}{4}(a^2+(4-c^2)b^2),$$

whence $b_\alpha(z,z)=2\,\mathrm{Tr}^+(\alpha z\bar{z})=\frac{1}{2}(\Lambda(a)+\Lambda'(b))$.

(b) The form $\mathrm{Tr}^+(\beta_0 xy)$ has determinant 1 on $\mathcal{O}^+$, since $\beta_0\mathfrak{D}=\mathcal{O}^+$. Multiplying the multiplier by $1/\nu$ multiplies the determinant by $\mathrm{N}_{L^+/\mathbb{Q}}(\nu)^{-1}=1/q$. Finally $4-c^2=-e^2$ is a totally positive unit, of norm 1. For the index: since $e=2\zeta-c$, we have $a+eb=(a-cb)+2\zeta b$, so $(a+eb)/2\in\mathcal{O}_L$ is equivalent to $a\equiv cb\pmod{2\mathcal{O}^+}$. The image of $\mathcal{O}_L$ thus has index 2^{44} in $\mathcal{O}^+\times\mathcal{O}^+$, and M_I has index $[\mathcal{O}_L:I]=q$ in it.

(c) If (a,b) and (a,b') are in M_I, then $e(b-b')\in 2I$. Since $b-b'$ is real and $\bar{e}=-e$, we also have $e(b-b')\in 2\bar{I}$, hence $e(b-b')\in 2(I\cap\bar{I})=2\nu\mathcal{O}_L$ and $b-b'\in 2\nu\mathcal{O}^+$. Conversely, for $x\in\mathcal{O}^+$, $e\nu x\in I$. The determinant of Λ' on $2\nu\mathcal{O}^+$ is $\mathrm{N}_{L^+/\mathbb{Q}}(2\nu)^2/q=(2^{44}q)^2/q=4^{44}q$. The argument is the same for a. □

The corresponding matrix identity, and the index $2^{44}q$, were checked exactly for each ideal treated (§7.2).

4.2. **Orbit of ζ and Parseval lemma.** Let $z\in I$ be non-zero, of norm $N=b_\alpha(z,z)$. For $k\in\mathbb{Z}/115$, we denote by (a_k,b_k) the image of $\zeta^k z$ and we set

$$f(k)=\mathrm{Tr}_{L/\mathbb{Q}}(\alpha\,\zeta^{2k}z^2)\in\mathbb{R}.$$

Lemma 4.2 (zero mean)**.** *For every* k*,* $\Lambda(a_k)=N+f(k)$ *and* $\Lambda'(b_k)=N-f(k)$*. In particular* $|f(k)|\le N$*. Moreover* $\sum_{k\in\mathbb{Z}/115}f(k)=0$ *and, for every* k*,*

$$\sum_{t=0}^{4}f(k+23t)=0. \tag{4.1}$$

[proved]

Proof. Set $z_k=\zeta^k z$; we have $z_k\bar{z}_k=z\bar{z}$. Then $a_k^2=z_k^2+\bar{z}_k^2+2z\bar{z}$ and $(4-c^2)b_k^2=-(z_k-\bar{z}_k)^2=-z_k^2-\bar{z}_k^2+2z\bar{z}$. Since $\mathrm{Tr}^+(\alpha(w+\bar{w}))=\mathrm{Tr}_{L/\mathbb{Q}}(\alpha w)$ for $w\in L$, we obtain $\Lambda(a_k)=f(k)+N$ and $\Lambda'(b_k)=-f(k)+N$. Both forms are positive, whence $|f(k)|\le N$. We have $f(k)=\sum_a\sigma_a(\alpha z^2)\,\omega^{2ak}$, where a runs over $(\mathbb{Z}/115)^\times$ and $\omega=\exp(2i\pi/115)$. The sum over k vanishes since $\sum_k\omega^{2ak}=0$. Relation (4.1) comes from $\sum_{t=0}^4\omega^{46at}=\sum_{t=0}^4\exp(2i\pi\cdot 2at/5)=0$, since 5 does not divide a. □

For a non-principal ideal, αz^2 belongs to $(\beta_0/\nu)I^2$, and $I^2/(I\bar{I}) = I/\bar{I}$ is not integral: the values $f(k)$ are not integers in general. The following lemma gives the estimate we need.

Lemma 4.3 (Parseval)**.** *For every non-zero $z \in I$, of norm N,*

$$\frac{1}{115}\sum_{k\in\mathbb{Z}/115} f(k)^2 = \sum_a \sigma_a(\alpha)^2\,|\sigma_a(z)|^4 \;\geq\; \frac{N^2}{88}.$$

[proved]

Proof. In the expression $f(k) = \sum_a \sigma_a(\alpha z^2)\omega^{2ak}$, the 88 frequencies $2a$ are pairwise distinct modulo 115. The orthogonality of the characters of $\mathbb{Z}/115$ gives the equality, with $|\sigma_a(\alpha z^2)| = \sigma_a(\alpha)|\sigma_a(z)|^2$. The inequality is the Cauchy–Schwarz inequality, applied to $N = \sum_a \sigma_a(\alpha)|\sigma_a(z)|^2$, a sum of 88 terms. □

The idea of bounding a norm from below by a Parseval identity on the coordinates of $\mathbb{Z}[\zeta_p]$ is already in Quebbemann [Que81, p. 168], for the Leech lattice ($p = 23$, rank 24). Lemma 4.3 applies it to the orbit of ζ in the gluing into two forms of rank 44; the next corollary adds the five-term relation (4.1).

Corollary 4.4 (threshold)**.** *Let $z \in I$ be non-zero, of norm N, and let $M = \max_k |f(k)|$. Then $M \geq N\sqrt{5/352}$. There therefore exists k such that*

$$\min\bigl(\Lambda(a_k), \Lambda'(b_k)\bigr) \leq N\Bigl(1 - \sqrt{5/352}\Bigr).$$

If $N \leq 6$, this minimum is at most $6 - \sqrt{45/88} < 1057/200$. [proved]

Proof. Five real numbers with zero sum and absolute value at most M have a sum of squares at most $4M^2$: the sum of squares is convex, and its maximum on this polytope is attained at a vertex, of the form $(M, M, -M, -M, 0)$ up to permutation. By (4.1), the 115 values of f split into 23 quintuples with zero sum. Hence $\sum_k f(k)^2 \leq 23 \cdot 4M^2$, and Lemma 4.3 gives $115N^2/88 \leq 92M^2$, that is, $M^2 \geq 5N^2/352$. If $f(k) = -M$, then $\Lambda(a_k) = N - M$; if $f(k) = M$, then $\Lambda'(b_k) = N - M$. Finally $6\sqrt{5/352} = \sqrt{45/88}$, and $6 - \sqrt{45/88} < 1057/200$ is equivalent to $(143/200)^2 = 20\,449/40\,000 < 45/88$, which is true. □

The inequality $M \geq N/\sqrt{88}$ alone would give the threshold $6(1 - 1/\sqrt{88}) \approx 5.3604$, which is slightly worse. The numerical value of the threshold is $6 - \sqrt{45/88} \approx 5.2849$.

Lemma 4.5 (zero projections)**.** *Let $z \in I$ be non-zero, of norm $N \leq 6$. There is at most one $k \in \mathbb{Z}/115$ such that $a_k = 0$ or $b_k = 0$. There exists k such that one of the two projections a_k, b_k is* non-zero *and of energy at most* $1057/200$. [proved]

Proof. We have $a_k = 0$ if and only if $\zeta^{2k} = -\bar{z}/z$, and $b_k = 0$ if and only if $\zeta^{2k} = \bar{z}/z$. Since $-1 \notin \langle\zeta\rangle$ and $k \mapsto \zeta^{2k}$ is injective on $\mathbb{Z}/115$, at most one k makes a projection vanish. Let k be given by Corollary 4.4. If the projection of minimal energy is non-zero, we are done. Otherwise, say $a_k = 0$; then $f(k) = -N$. By (4.1), the four other terms of the quintuple of k have sum N; one of them, $f(k')$, is at least $N/4$. Then $\Lambda'(b_{k'}) = N - f(k') \leq 3N/4 \leq 9/2 < 1057/200$, and $b_{k'} \neq 0$ by uniqueness. The case $b_k = 0$ is symmetric. □

4.3. **The criterion.**

Theorem 4.6 (criterion)**.** *The lattice (I, b_α) has no non-zero vector of norm at most 6 if and only if the following two conditions are satisfied.*

(C1) *There is no $(a, b) \in M_I$ with $a \neq 0$, $\Lambda(a) \leq 1057/200$ and $\Lambda(a) + \Lambda'(b) \leq 12$.*
(C2) *There is no $(a, b) \in M_I$ with $b \neq 0$, $\Lambda'(b) \leq 1057/200$ and $\Lambda(a) + \Lambda'(b) \leq 12$.*

Under these conditions, the lattice, which is even, has minimum at least 8. [proved]

Proof. A pair excluded by (C1) or (C2) is the image of a non-zero vector of norm at most 6 (Proposition 4.1). Conversely, if $z \neq 0$ has norm at most 6, Lemma 4.5 provides a rotation $\zeta^k z \in I$, of the same norm, one of whose projections is non-zero and of energy at most $1057/200$; the sum of the two energies is $2N \leq 12$. This rotation contradicts (C1) or (C2). □

We call *side 1* the verification of (C1) and *side 2* that of (C2). Each side requires two nested enumerations: that of the non-zero short projections a (resp. b) of energy at most $1057/200$, in rank 44; then, for each of them, that of the vectors of the prescribed class modulo $2\nu\mathcal{O}^+$ whose energy respects the bound 12. The criterion concludes only if these enumerations are exhaustive. A capped search, or a search at a threshold below $1057/200$, does not give a negative verdict.

Remark 4.7. The methods of [Sak26] (tensor inequality and projective minima) require a tensor product structure over the ring of integers of $\mathbb{Q}(\sqrt{-23})$. Among the classes of S, they apply only to the principal class and to those of support $\{3\}$.

5. The certified enumeration

5.1. **Data and principle.** For a given ideal I and a given side, a PARI/GP script produces integer data: a Gram matrix A of the short projection, scaled by an integer D_A; a Gram matrix B of the lattice $2\nu\mathcal{O}^+$ of the fiber, scaled by an integer D_B; an integer matrix T_n and an integer $d > 0$ which describe the lifts. The two Gram matrices are reduced (BKZ with block size 20), and the final transformation is checked: it is unimodular and the change-of-basis identity is exact. For $u, y \in \mathbb{Z}^{44}$, we set $a(u) = u^{\mathsf{T}} A u$ and $w = d\,y + T_n u$. The lattice vector associated with (u, y) has norm

$$N = \frac{1}{2}\Big(\frac{u^{\mathsf{T}} A u}{D_A} + \frac{w^{\mathsf{T}} B w}{D_B\, d^2}\Big). \tag{5.1}$$

For P_{139}, $(D_A, D_B, d) = (139, 1, 278)$; for P_{599}, $(D_A, D_B, d) = (599, 1, 1198)$.

Let B_A be a positive rational number and S a positive integer (here $B_A = 1057/200$ and $S = 12$). We say that u is *normalized* if $u \neq 0$ and its last non-zero coordinate is positive. We set

$$\mathcal{U} = \{u \text{ normalized} :\ a(u) \leq D_A\, B_A\},$$
$$\mathcal{T} = \{(u, y) :\ u \in \mathcal{U},\ y \in \mathbb{Z}^{44},\ a(u)\, D_B d^2 + D_A\, w^{\mathsf{T}} B w \leq S\, D_A D_B d^2\}.$$

With $B_A = 1057/200$ and $S = 12$, $\mathcal{T}$ is the set of vectors of norm at most 6 whose short projection is non-zero and of energy at most $1057/200$, up to sign. The change $(u, y) \mapsto (-u, -y)$ preserves both energies: the normalization loses nothing. By Theorem 4.6, if $\mathcal{T} = \emptyset$ for both sides, the lattice has no non-zero vector of norm at most 6.

The program `Rang88CertZ` (Java) enumerates $\mathcal{U}$ by the Fincke–Pohst method [FP85], then, for each $u \in \mathcal{U}$, the y such that $(u, y) \in \mathcal{T}$. It differs from an ordinary floating-point enumerator in the following points.

- The decomposition $G = \sum_i d_i T_i^2$ of each form is computed in exact rationals, and the identity is checked exactly at run time.
- The partial sums that give the centers are exact integers, in fixed point with step 2^{-40} in the executions described here, of type `long` with overflow detection. The truncation error is bounded exactly by an integer sum.
- The only floating-point operations are the energy lower bound and the radius. Each of them is followed by a move to the outward floating-point neighbor, so that a branch is pruned only if a certified lower bound of its energy exceeds a certified upper bound of the budget.

- Each leaf is rechecked in exact integers.
- The tree is split into tasks by the same recursion. Each completed task leaves a one-line *receipt* in a checkpoint file, with its counters. The final summary prints `COUVERTURE_COMPLETE` ("complete coverage") only if every task of the catalogue has a result.

5.2. **What the non-omission proof establishes.** The model of the machine is the following.

(M1) The operations on `double` values and the conversion of a `long` into a `double` return the round-to-nearest result of the IEEE 754 standard; `Math.sqrt` returns the `double` closest to the exact square root.

(M2) The integer methods said to be exact return the exact result or throw an exception; the classes `BigInteger` and `BigDecimal` are exact.

(M3) Hardware faults and file alterations are outside the model.

Theorem 5.1 (non-omission)**.** *Assume* (M1) *and* (M2)*. If an execution of* `Rang88CertZ` *terminates with exit code* 0 *and the line* `COUVERTURE_COMPLETE`*, then, under the integrity hypothesis on the checkpoint file:*

(1) *A and B are positive definite;*
(2) *if no counter has overflowed (representability hypothesis), the counter* `COMPTE_U` *("count of u") is equal to $|\mathcal{U}|$;*
(3) *under the same representability hypothesis, the counter* `TOTAL_TOUCHES` *("total number of hits") is equal to $|\mathcal{T}|$.*

[proved under M1–M3; proof reviewed once by a second assistant, within a limited scope]

Corollary 5.2 (absence of hits, without hypothesis on the counters)**.** *Assume* (M1) *and* (M2)*. Consider an execution of* `Rang88CertZ` *in* `run` *mode, with a fresh checkpoint file and without test receiver, terminated with exit code* 0 *and the line* `COUVERTURE_COMPLETE`*. If the directory of the checkpoint file contains no hit file, and none has been removed from it, then $\mathcal{T} = \emptyset$.* [relies on Z1–Z8; link to the code reviewed by a second assistant]

Sketch. By Z1 to Z8, every element of $\mathcal{T}$ is an admissible leaf of a task of the catalogue, and the catalogue is a partition of the unpruned tree. With a fresh checkpoint file, no earlier receipt is loaded: the summary is built only from the tasks of this launch, and the line `COUVERTURE_COMPLETE` is printed only if each of them has returned a result. An admissible leaf causes, with no condition on the value of a counter, a line to be written in the hit file located next to the checkpoint file; this file is created if needed, and a write error throws an exception, which propagates up to exit code 2. Hence, if $\mathcal{T} \neq \emptyset$, either a hit file exists or the exit code is not 0. The line-by-line link with the source is given in the repository accompanying the note. □

The complete proof is in the repository accompanying the note; it is linked to the source by a table that associates each step of the code with a lemma. We only give its structure here.

lemma	content
L1	Decomposition $G = \sum_i d_i T_i^2$ in exact rationals, checked at run time; G is positive definite.
L2	Enclosure of a rational number between two `double` values, by exact comparisons.
L3	Outward floating-point neighbors: the exact result of an operation is enclosed between the predecessor and the successor of the rounded result.
Z1	Fixed point: the gap between the exact center and the integer center is bounded by the sum of the absolute values of the coordinates already fixed.
Z2	The cache of integer partial sums is exact at every node (induction invariant).
Z3	The floating-point budgets bound the exact budgets from above, on the short side and in the fiber.
Z5	The energy lower bound passed from node to node bounds the exact energy from below; a pruned branch contains no admissible vector.
Z4	The integer range scanned at each level contains all admissible values; the integer division is exact.
Z6	The sign normalization removes exactly the zero vector and the opposites.
Z7	The tasks of the catalogue form a partition of the unpruned tree.
Z8	The leaves are tested in exact integers: a pair is counted if and only if it lies in $\mathcal{T}$.
Z9	Every integer overflow that could distort a decision throws an exception; an exception gives exit code 2, without `COUVERTURE_COMPLETE`.
Z10	Checkpoint file: binding to the parameters, the catalogue, the inputs and the classes; rereading; summary.

Lemmas L1 to L3 come from a first certifying program, based on floating-point intervals, which is slower. It certified six isolated tasks of P_{139}, without any hit, and was not run to completion.

Validation [observed]. The program was run on cases with a known answer. On L_0, that is, for $I = \mathcal{O}_L$, the two sides give 403 876 short projections, namely $4\,004 + 399\,872$, values obtained earlier by other programs, and no hit. On the rejected ideal P_{461}, at threshold $7/2$, it finds 697 projections and 2 hits of norm 6. A test harness compares, on 5 400 random forms of dimension 2 to 4, the enumeration with an exhaustive search in integers: 16 250 checks, no failure. A hard stop followed by a resumption gives the same totals. These validations concern finite cases. They do not replace the proof and say nothing about P_{139} or P_{599}.

5.3. **Review of the proof.** The proof was written with the assistant that wrote the code, then reviewed once by the second assistant. This review concluded favorably, within a limited scope: the non-omission mechanism Z1–Z8, under (M1) and (M2), and the domain refusals. It does not amount to a certification of a final result, and it formulated four reservations.

- *Z9.* The local counters do not detect their own overflow. Exact counting requires a representability hypothesis, or checked increments.
- *Z10.* A complete coverage that relies on reread receipts does not prove, by itself, that the corresponding tasks were executed: the loader checks the syntax, the indices and the duplicates, and replays nothing. A fabricated file would pass the resumption. The integrity hypothesis of the proof excludes this case; it must be documented.
- *Writes.* The writing of a receipt is not looped until the buffer is exhausted, and the lock is internal to the virtual machine; the uniqueness of the writer is not guaranteed across processes.

- *Identity.* The program computes the hash of the class files as reread from disk, not that of the bytes loaded into memory. The identity of the files, the geometric link and the final result are three distinct points; none of them is certified by this review.

In particular, the review of Z1–Z8 does not by itself cover the exactness of the counters asserted by points (2) and (3) of Theorem 5.1. The absence of overflow, the exactness of the counters and the detection of a witness are three distinct propositions; only the third one is used for the conclusion $\mathcal{T} = \emptyset$.

5.4. **Hypotheses H1 to H4, and what is not established.** The statement "no non-zero vector of norm at most 6" for a candidate is *computer-assisted*. It relies on Theorem 4.6, on Theorem 5.1, on the exact link between the input files and the lattice, and on the following hypotheses.

(H1) *Machine.* The Java virtual machines used execute the classes according to their specification, just-in-time compiler included: these are (M1) and (M2). Two environments are involved: OpenJDK 11.0.16.1 under Windows for the first executions, OpenJDK 21 under Linux for the replications. We certify neither of these two implementations.
(H2) *Hardware.* No hardware or memory fault occurred during the executions.
(H3) *Documentary integrity.* The logs are the authentic output of the executions; the bytes of the loaded classes and of the inputs read are those whose hashes the executions printed.
(H4) *Resumption (specific to side 1 of P_{599}).* The 21 459 receipts reread at the resumption were written by the first launch of the same program, each after the complete execution of its task.

Some of these reservations are lifted for the executions concerned, by a reading of the source, of the logs and of the receipts [exact checks on the files].

- *Z9.* No counter enters an enumeration decision. The absence of hits depends on no counter: every hit creates a file or throws an exception, and no hit file exists. The printed totals are exact as long as no counter has undergone 2^{63} increments, which the duration of the executions makes implausible; this is a heuristic argument, without a proved bound. The conclusion on the minimum does not depend on it.
- *Z10 and writes, without resumption.* For P_{139} and for side 2 of P_{599}, each side was executed in a single launch. The checkpoint file was opened for writing only, and the summary depends on no reread receipt. The receipts were checked as witnesses: format, permutation of the indices, sums equal to the totals, final hash. The replications of P_{139} and of P_{599} on the second computer are also without resumption, for all four sides: the conclusion can rely on these fresh executions, in which H4 plays no role.
- *Identity.* A recompilation of the source outside the repository, with the same compiler, gives back the classes byte for byte.

The following are *not* established:

- the validity of the proof beyond the limited review above;
- H1; an execution by an independent implementation would reduce the risk of an error common to the proof and to the program, and none has been made;
- H2 and H3: the replications on a second computer support them, without proving them.

Replications. For each of the two lattices, both sides were executed a second time, on a second computer, under another operating system and another version of the virtual machine (OpenJDK 21), in a single launch without resumption; the receipts are identical byte for byte to those of the first execution, and no divergence was observed in the compared results. These replications strengthen the consistency checks between executions, without proving H2 or H3. For P_{599}, side 1 of the first execution had been resumed; the complete

replay of the 21 459 tasks executed before the resumption reproduced their receipts byte for byte, and the conclusion relies in any case on the executions without resumption, in which H4 plays no role. The receipts are deterministic. A replication must therefore reproduce, byte for byte, the hash of the receipts sorted by index.

6. Results

We denote by P_p a prime ideal of $\mathbb{Z}[\zeta]$ above the prime number p, and by $X_p = (P_p, \beta_0/\nu)$ the unimodular ideal lattice that it carries (Proposition 2.3). The prime ideals above p are conjugate under $\mathrm{Gal}(L/\mathbb{Q})$: the corresponding lattices X_p are isometric, and form a single orbit of S. A statement on the minimum of X_p therefore holds for this whole orbit. If $p \equiv 1 \pmod{115}$, $P_p = (p, \zeta - r)$ has degree 1 and $q = p$. If $p \equiv 24$ or $91 \pmod{115}$, P_p has degree 2 and $q = p^2$; its class is fixed by σ_{24}, resp. σ_{91}.

6.1. Proved rejections.

A *witness* of non-extremality is an integer vector x, in an explicit $\mathbb{Z}$-basis of the ideal, such that $x^{\mathsf{T}} G x = 6$, where G is the exact Gram matrix of the lattice.

Proposition 6.1. *For each of the eight ideals of Table 2, the lattice X_p contains a vector of norm 6. It is therefore not extremal, and no lattice of its orbit is. Its minimum, which is at most 6, is not determined.* [proved, without GRH]

Table 2. Rejections. The rejection is proved, without GRH. The "orbit" column is under GRH for the 331- and 45 013-components, without GRH for the 3-component (§6.4). Search: "GP, 7/2" denotes the PARI/GP filter at threshold 7/2, "Java" a non-certifying Java enumerator, "CertZ" the program of §5. The SHA-256 hashes of the witness files are abbreviated.

ideal	q	orbit (support, size)	search	witness	norm
$P_{461} = (461, \zeta - 5)$	461	$\{3, 331, 45\,013\}$, 88	GP, 7/2	`d41592a8...`	6
$P_{691} = (691, \zeta - 4)$	691	$\{331, 45\,013\}$, 88	GP, 7/2	`f3cf48f3...`	6
$P_{1151} = (1151, \zeta - 6)$	1 151	$\{331, 45\,013\}$, 88	GP, 7/2	`a5705522...`	6
$P_{1381} = (1381, \zeta - 21)$	1 381	$\{3, 331, 45\,013\}$, 88	GP, 7/2	`4a45c630...`	6
P_{1471} (degree 2)	1471^2	$\{45\,013\}$, 44	Java	exact check	6
P_{829} (degree 2)	829^2	$\{331\}$, 22	CertZ	`52222d1c...`	6
P_{1289} (degree 2)	1289^2	$\{3, 331\}$, 44	CertZ	`4c5f7602...`	6
P_{1979} (degree 2)	1979^2	$\{3, 331\}$, 44	CertZ	`71d980e2...`	6

Justification. For the seven ideals other than P_{1471}, a witness file contains the minimal polynomial of c, β_0, ν, α, a $\mathbb{Z}$-basis of the ideal (88×88 matrix in the basis of the powers of ζ), the Gram matrix, the vector x and its expression in L. A verification program, which uses neither class group computations nor session data, rechecks exactly: that the basis generates an integral ideal of $\mathbb{Z}[\zeta]$, stable under ζ, containing p and of the expected index (p or p^2); that $\alpha = \beta_0/\nu$; that the Gram matrix, recomputed, is integral, even, of determinant 1 and positive definite; that x is integral and that $x^{\mathsf{T}} G x = 6$. The seven logs carry the mention "témoin valide, norme 6" (valid witness, norm 6). For $P_{1471} = (1471, \zeta^2 + 879\zeta + 948)$, the vector was found by a non-certifying Java enumerator, then checked independently, without PARI/GP or Java: the reconstructed Gram matrix is even, positive definite, of determinant 1, and $\mathrm{Tr}(\alpha z \bar{z}) = 6$. □

For the last three ideals, the hit decomposes according to (5.1) into a short energy and a fiber energy with sum 12:

$$\frac{3161}{829} + \frac{22\,505\,692}{1658^2}, \qquad \frac{5683}{1289} + \frac{50\,451\,460}{2578^2}, \qquad \frac{9260}{1979} + \frac{114\,687\,008}{3958^2}.$$

These three hits were obtained in less than five minutes each, by the program and with the parameters used for the two candidates [observed].

Remark 6.2. The PARI/GP filter used for the first four ideals is a capped search, at a threshold below 1057/200. It provides witnesses, which are exact. An absence of witness in its outputs proves nothing: at threshold 3, this filter had found nothing for P_{1471}, which is nevertheless rejected. For P_{2531}, of degree 1, the filter was interrupted by its time limit; there is no decision. That the four orbits of size 88 in the table are pairwise distinct is not established here.

6.2. **The two candidates.** The following two ideals have residue degree 2, with $p \equiv 24 \pmod{115}$:

$$P_{139} = (139,\ \zeta^2 + \zeta + 77), \qquad P_{599} = (599,\ \zeta^2 + 87\zeta + 322).$$

We have $\bar{P}_{139} = (139, \zeta^2 + 65\zeta + 65) \neq P_{139}$ and $\bar{P}_{599} = (599, \zeta^2 + 523\zeta + 426) \neq P_{599}$. The absolute norms are $q = 19\,321$ and $q = 358\,801$.

Proposition 6.3 (candidate P_{139})**.** *Let $X = X_{139}$.*

(1) *X is a positive definite even unimodular lattice of rank* 88. [proved]
(2) *X contains a vector of norm* 8. [proved]
(3) *X contains no non-zero vector of norm at most* 6. [computer-assisted: H1–H3]

Overall status: minimum ≥ 8 computer-assisted, two executions on two computers (identical receipts).

Justification. (1) Exact check, by a Python program in integers and rationals, without PARI/GP: the basis is that of an ideal of $\mathbb{Z}[\zeta]$ of index 139^2, equal to P_{139}; the Gram matrix, recomputed from α, is integral, even, of determinant 1, and its 88 leading principal minors are positive. The same program rechecks the gluing (Proposition 4.1) and the link between the input files of the certifying program and the lattice, for both sides: 43 checks, no failure. This link was also reviewed by a second assistant.

(2) A light search gives a pair (u, y) with energies $398/139$ and $1\,015\,256/278^2$, of sum 16. The corresponding vector, reconstructed in the basis of the ideal, has integer coordinates and satisfies $x^{\mathsf{T}} G x = 8$, in exact integers.

(3) The program `Rang88CertZ` was run on both sides, with $B_A = 1057/200$ and $S = 12$, from October 10 to 11, 2026, on 6 threads. Each side was run in a single launch, without resumption.

side	tasks	duration	printed $\lvert\mathcal{U}\rvert$	hits	end
1	48 227	14 562 s	42 957 972	0	complete coverage, code 0
2	61 174	15 856 s	42 957 972	0	complete coverage, code 0

By Corollary 5.2, under H1–H3, $\mathcal{T} = \emptyset$ for both sides: the executions are without resumption and their output directories contain no hit file. Theorem 4.6 then gives (3). The hashes of the sorted receipts begin with `c99dcf87` (side 1) and `8ab1f40a` (side 2). □

Replication of P_{139}. The same program, on the same inputs, was executed a second time, on a second computer, under Linux and Java 21 (the first execution was under Windows and Java 11), with 8 threads instead of 6, without resumption: 48 227 and 61 174 tasks, no hit. The receipts sorted by index have the same SHA-256 hashes as those of the first execution (`c99dcf87...5122` and `8ab1f40a...daf7`). The two executions share the source code and the class files: this is not an independent implementation.

Proposition 6.4 (candidate P_{599})**.** *Let $X = X_{599}$.*

(1) *X is a positive definite even unimodular lattice of rank* 88. [proved]
(2) *X contains a vector of norm* 8. [proved]

(3) *X contains no non-zero vector of norm at most* 6. [computer-assisted: H1–H3]

Overall status: minimum ≥ 8 computer-assisted, two executions on two computers (identical receipts). Hypothesis H4 on the resumed receipts of side 1 of the first execution is not needed (see below).

Justification. (1) Same exact check as for P_{139}: 47 checks, no failure, for both sides. This link was also reviewed by a second assistant.

(2) A pair (u, y) with energies $1376/599$ and $19\,666\,368/1198^2$, of sum 16, gives a vector with integer coordinates such that $x^{\mathsf{T}}Gx = 8$.

(3) First execution, on the first computer:

side	launches	tasks	printed $\lvert\mathcal{U}\rvert$	hits	end
1	two: 6 threads, forced stop after about 20 350 s; then 10 threads, 19 765 s	59 255	185 794 630	0	complete coverage, code 0
2	one: 10 threads, 34 114 s	74 398	185 794 630	0	complete coverage, code 0

Side 2 has no resumption: under H1–H3, $\mathcal{T} = \emptyset$ for this side. Side 1 was resumed: the second launch reread 21 459 receipts and executed 37 796 tasks. Under H1–H4, $\mathcal{T} = \emptyset$ for this side. The replication described below gives the same conclusion without H4: both of its sides are without resumption, and Corollary 5.2 applies to them: the complete inventory of the output directory, taken after the end of the computation, contains no hit file. The hashes of the sorted receipts begin with `f3cf379a` (side 1) and `b9c93a83` (side 2). □

On hypothesis H4. An exact check of the files establishes the following points. The file reread at the resumption is, byte for byte, the beginning of the final file; a copy taken before the resumption has the same hash. The forced stop left no truncated line. The 21 459 resumed indices and the 37 796 indices executed afterwards are disjoint, and their union is exactly the set of the 59 255 tasks. This check does not establish that the resumed receipts were produced by the same program, nor even by an execution: the second launch accepted them after a format check. Several concordances support H4 without proving it [observed]: the log of the first launch; the chronology; the agreement, at each periodic progress report, between the announced totals and the sums of the receipts already written; the order of the lines, compatible with a queue served by 6 then by 10 threads; the equality of $\lvert\mathcal{U}\rvert$ and of its histogram with those of an earlier sweep by another, non-certifying, program; finally the replay of 200 tasks drawn at random among the 21 459, whose receipts are identical byte for byte. This sample covers 0.93% of the tasks; it would not detect an isolated alteration.

Replay of the resumed tasks of P_{599}. The 21 459 tasks executed before the interruption were replayed in full, with the same classes and the same inputs, on a separate checkpoint (14 564 s, no hit). The 21 459 receipts are identical byte for byte to those of the checkpoint taken before the resumption and to those of the final checkpoint. Hypothesis H4 is therefore no longer needed.

Replication of P_{599}. The complete sweep was redone on the second computer (Linux, OpenJDK 21, 14 threads), with the same compiled classes and the same input files, in a single launch without resumption: side 1 in 26 212 s, side 2 in 25 956 s, exit codes zero, no hit. The task catalogues are identical byte for byte to those of the first execution. The canonical hashes of the receipts (receipts sorted by index, without header) are identical for both sides: 59 255 and 74 398 receipts. The hashes were computed twice, by two distinct programs. Statement (3) remains computer-assisted: the replication strengthens the consistency checks between executions, without proving H2 or H3; both machines executed the same program, and it says nothing about an omission specific to that program.

Remark 6.5. Before these executions, a non-certifying floating-point enumerator had swept both sides of the two candidates without finding any hit, with the same values of $|\mathcal{U}|$ [observed]. By Remark 2.6, Table 1 and Remark 6.8, the normalizers of the group of multiplications $\langle\zeta\rangle$ in $\mathrm{Aut}(X_{139})$ and $\mathrm{Aut}(X_{599})$ have orders 460 and 920, respectively. In particular,

$$|\mathrm{Aut}(X_{139})| \geq 460, \qquad |\mathrm{Aut}(X_{599})| \geq 920.$$

Explicit generators of these two normalizers have been constructed and checked in exact arithmetic. The full automorphism groups have not been determined; perfection has not been studied.

6.3. **Partial survivors of a filter.** The primes $p \equiv 24 \pmod{115}$ give ideals of degree 2 whose class is fixed by σ_{24}, hence with support contained in $\{3, 331\}$. Apart from the orbits of L_0 and of L_S, the classes of this type form 30 orbits: 15 of support $\{331\}$ and 15 of support $\{3, 331\}$ (Table 1). Nine cases were examined: $p = 139$, 599, 829, 1289, 1979, 3359, 4049, 4969 and 5659. According to a computation by `bnfinit`, they lie in nine distinct orbits [under GRH]. The primes $p \equiv 114 \pmod{115}$ give nothing more: their class is fixed by complex conjugation, hence trivial.

For the last four, a filter was applied: the program `Rang88CertZ` on side 1 only, with a stop at the first hit, for a bounded duration (one hour on 4 threads, then 55 minutes on 6 threads).

TABLE 3. Partial survivors of the filter, side 1 only. Status: observed. No conclusion on the minimum. The "orbit" column is under GRH.

ideal	orbit (support, size)	receipts / tasks of side 1	share of the tasks	hits
P_{3359}	$\{331\}$, 22	3 981 / 109 662	3.6%	0
P_{4049}	$\{3, 331\}$, 44	4 173 / 77 819	5.4%	0
P_{4969}	$\{3, 331\}$, 44	3 716 / 97 270	3.8%	0
P_{5659}	$\{331\}$, 22	4 201 / 132 743	3.2%	0

These four ideals are *partial survivors*, and nothing more. The coverage is a few percent of a single side, and the tasks are not of equal size. The three rejections of the same family occurred after 302 298, 354 888 and 432 358 short projections; in the first pass, the four survivors had each gone through ten to fifteen million of them without a hit. This is an indication, without proof value. Nothing is known about the minimum of these four lattices.

6.4. **Classes of P_{139} and of P_{599} without GRH; non-isometry.** *Localization under GRH.* The p-component of a class $[P]$ can be read in the fixed field K of the kernel of χ_p: the norm $\mathrm{N}_{L/K}$ induces an isomorphism on the χ_p-component, because the degree $[L : K]$ is invertible modulo p. We therefore read the 3-component in $\mathrm{Cl}(F) \cong \mathbb{Z}/3$, certified without GRH, and the 331- and 45 013-components in $\mathrm{Cl}(K_{22}) \cong \mathbb{Z}/662$ and $\mathrm{Cl}(K_{44}) \cong \mathbb{Z}/45\,013$, computed by `bnfinit` [under GRH]. The supports and orbit sizes of Tables 2 and 3 come from this computation. For P_{139}, it gives the support $\{3, 331\}$ and an orbit of size 44; for P_{599}, the support $\{331\}$ and an orbit of size 22.

The following two propositions do not depend on GRH. Let $K = K_{22} = \mathbb{Q}(\theta)$, with $\theta = \zeta + \zeta^{22} + \zeta^{24} + \zeta^{68}$, the fixed field of $\langle\sigma_{22}\rangle$; it is a CM field of degree 22, whose unit group has rank 10 and whose only roots of unity are ± 1.

Proposition 6.6. *The ideal P_{139}^3 is not principal in L.* [proved, without GRH]

Proposition 6.7. *For every integer m prime to 331, the ideal P_{599}^m is not principal in L. The order of $[P_{599}]$ is therefore divisible by 331. Moreover, the 3-component of $[P_{599}]$ is zero.* [proved, without GRH]

Proof of Propositions 6.6 and 6.7. Let $p \in \{139, 599\}$, $P = P_p$ and $\mathfrak{q} = P \cap K$. A certificate, verified by a Python program in exact arithmetic, without PARI/GP, establishes the following facts (24 checks for $p = 139$, 37 for $p = 599$, no failure).

(i) The minimal polynomial of θ has degree 22 and has 22 simple roots modulo p; $\mathfrak{q}$ has degree 1, P has degree 2 over $\mathfrak{q}$, and $\mathrm{N}_{L/K}(P) = \mathfrak{q}^2$.

(ii) An explicit algebraic integer $\gamma \in K$ satisfies $(\gamma) = \mathfrak{q}^{331}$. The elements $\xi_a = \sin(\pi a/23)/\sin(\pi/23)$, $2 \le a \le 11$, are units of K.

(iii) For three primes $\ell \equiv 1 \pmod{331 \cdot 115}$, we form the matrix of the discrete logarithms modulo 331 of the images of $\xi_2, \dots, \xi_{11}, \gamma$ in the residue fields of degree 1 above ℓ. The ten unit columns have rank 10, and the whole matrix has rank 11.

Let E be the unit group of K and $U' = \langle -1, \xi_2, \dots, \xi_{11} \rangle$. For a prime $\mathfrak{l}$ of degree 1 above ℓ, let $\lambda_\mathfrak{l}$ be the reduction morphism with values in $\mathbb{F}_\ell^\times/(\mathbb{F}_\ell^\times)^{331} \cong \mathbb{F}_{331}$; it vanishes on the 331st powers and on ± 1.

Step 1: U' has finite index k in E, and $331 \nmid k$. If $\prod \xi_a^{n_a} = \pm\eta^{331}$ with $\eta \in E$, then $\sum n_a \lambda_\mathfrak{l}(\xi_a) = 0$ for every $\mathfrak{l}$, hence 331 divides each n_a, by the rank 10. A primitive multiplicative relation among the ξ_a would contradict this: U' has rank 10, hence finite index. If 331 divided k, there would exist $\varepsilon \in E \smallsetminus U'$ with $\varepsilon^{331} \in U'$; then $\varepsilon^{331} = \pm\eta'^{331}$ with $\eta' \in U'$, so ε/η' would be a root of unity of K, that is, ± 1, and $\varepsilon \in U'$.

Step 2: for m prime to 331, $\mathfrak{q}^{2m}$ is not principal. Otherwise $\mathfrak{q}^{2m} = (\delta)$, and $(\gamma^{2m}) = \mathfrak{q}^{662m} = (\delta^{331})$, that is, $\gamma^{2m} = \varepsilon\delta^{331}$ with $\varepsilon \in E$. At the ℓ used, δ is a local unit. Then $2mk\,\lambda_\mathfrak{l}(\gamma) = \lambda_\mathfrak{l}(\varepsilon^k)$ for every $\mathfrak{l}$, with $\varepsilon^k \in U'$: the column of γ would lie in the span of the unit columns, since $2mk$ is invertible modulo 331. The rank 11 forbids this.

Step 3. If $P^m = (\beta)$, then $\mathrm{N}_{L/K}(\beta)$ generates $\mathrm{N}_{L/K}(P)^m = \mathfrak{q}^{2m}$. Hence P^m is not principal. For $p = 599$, this is the first assertion of Proposition 6.7. For $p = 139$, we only use the case $m = 3$, which is Proposition 6.6.

The 3-component of $[P_{599}]$. The element $s = \sum_{a=1}^{22} \left(\frac{a}{23}\right)\zeta^{5a}$ satisfies $s^2 = -23$, and $599 = 24^2 + 23$. The element $\pi = 24 - s$ lies in P_{599}, so $P_{599} \cap \mathcal{O}_F = (\pi)$ is principal and $\mathrm{N}_{L/F}(P_{599}) = (\pi^2)$. The Sylow 3-subgroup of $\mathrm{Cl}(L)$ has order 3 (Proposition 2.2), and $\mathrm{Cl}(F) \cong \mathbb{Z}/3$. The extension of ideals $j\colon \mathrm{Cl}(F) \to \mathrm{Cl}(L)$ is injective, since $\mathrm{N}_{L/F} \circ j$ is multiplication by 44; its image is therefore this Sylow subgroup. Write $[P_{599}] = j(a) + c'$, with c' of order prime to 3. Then $0 = \mathrm{N}_{L/F}([P_{599}]) = 44a + \mathrm{N}_{L/F}(c')$. Now $\mathrm{N}_{L/F}(c')$ has order prime to 3 in a group of order 3, hence is zero; therefore $a = 0$. □

Remark 6.8 (what remains under GRH, and what has been reviewed)**.** (a) For P_{139}, the ideal $P_{139} \cap \mathcal{O}_F$ is not principal ($4 \cdot 139$ is not of the form $a^2 + 23b^2$), so the order of $[P_{139}]$ is divisible by 3. With Proposition 6.6, this order is divisible by 331 or by 45 013. The 45 013-component is zero, by the argument of point (b): $139 \equiv 24 \pmod{115}$, so σ_{24} is the Frobenius substitution of 139 and fixes P_{139}, whereas 24 is not in the kernel of $\chi_{45\,013}$. The order of $[P_{139}]$ therefore divides $993 = 3 \cdot 331$; it is divisible by 3 and different from 3: it equals 993, and the support is $\{3, 331\}$. The stabilizer of the class is the intersection of the kernels of χ_3 and of χ_{331}, namely $\{1, 24\}$ (22 and 68 are congruent to -1 modulo 23, which is not a square): the orbit has size 44. These three facts are therefore obtained without GRH, modulo Propositions 2.2 and 2.5. *Which* of the 15 orbits of this support is that of P_{139} remains under GRH.

(b) For P_{599}, the 45 013-component is zero: σ_{24} is the Frobenius substitution of 599 and fixes P_{599}, and 24 is not in the kernel $\{1, 91\}$ of $\chi_{45\,013}$ (Proposition 2.5). With Propositions 2.2, 2.5 and 6.7, $[P_{599}]$ has order 331, support $\{331\}$ and an orbit of size 22, without GRH. *Which* of the 15 orbits of this support is that of P_{599} remains under GRH.

(c) The class certificates of P_{139} and of P_{599} were reviewed by a second assistant, within this scope only: "P^m is not principal" for the indicated m. The computation of the 3-component and the exact order 331 in L rely on Propositions 2.2 and 2.5.

The statements cited from [Sak26] are the following.

(S1) L_0 and L_S have minimum 8 [Sak26, §5, Prop. 5.5]. This is a preprint.
(S2) $L_0 \cong (\mathcal{O}_L, \beta_0)$ (integer certificate, §2.5), and L_S is the ideal lattice of $J\mathcal{O}_L$, with $[J\mathcal{O}_L]^3 = 1$ (Proposition 2.7).

Proposition 6.9 (non-isometry)**.** (a) *Neither X_{139} nor X_{599} is isometric to L_0 or to L_S.*
(b) *The classes $[P_{139}]$ and $[P_{599}]$ are not conjugate under* $\mathrm{Gal}(L/\mathbb{Q})$*. If at least one of the lattices X_{139}, X_{599} has minimum* 8*, then $X_{139} \ncong X_{599}$.*
[proved, without GRH, modulo (S1), (S2) and Theorem 3.3]

Proof. (a) Let X be one of the two lattices, carried by P. If $\min X \leq 6$, the claim is immediate by (S1). Otherwise $\min X = 8$ (Propositions 6.3 and 6.4, point 2), and X, L_0, L_S are three unimodular ideal lattices of minimum at least 8 over L. By Theorem 3.3, an isometry of X with L_0 or L_S would give a and $\lambda \in L^\times$ such that $\lambda\sigma_a(P)$ is $\mathcal{O}_L$, $J\mathcal{O}_L$ or its conjugate. Then $[\sigma_a(P)]^3 = 1$, hence $[P]^3 = 1$, which contradicts Propositions 6.6 and 6.7.
(b) The order of $[P_{139}]$ is divisible by 3 (Remark 6.8), that of $[P_{599}]$ is not, and two conjugate classes have the same order. If only one of the two lattices has minimum 8, they differ by their minimum; if both do, Theorem 3.3 applies. □

The proof of (a) does not use the computer-assisted statement on the minimum: it holds whether or not X has minimum 8.

Remark 6.10 (a shorter proof of (b), without the class number of L)**.** Let $F = \mathbb{Q}(\sqrt{-23}) \subset L$. Since $L/\mathbb{Q}$ is abelian, $\mathrm{N}_{L/F}(\sigma(I)) = \sigma|_F(\mathrm{N}_{L/F}(I))$ for every $\sigma \in \mathrm{Gal}(L/\mathbb{Q})$, and the norm of a principal ideal is principal: whether $\mathrm{N}_{L/F}(I)$ is principal or not depends only on the Galois orbit of the class of I. We have $\mathrm{Cl}(F) \cong \mathbb{Z}/3$: the reduced forms of discriminant -23 are $(1,1,6)$, $(2,\pm1,3)$.

- 139 splits in F and has residue degree 2 in L; hence $\mathrm{N}_{L/F}(P_{139}) = \mathfrak{p}_{139}^2$. A generator of $\mathfrak{p}_{139}$ would be $(a + b\sqrt{-23})/2$ with $a^2 + 23b^2 = 556$; for $|b| \leq 4$, $556 - 23b^2$ takes the values 556, 533, 464, 349, 188, none of which is a square. Hence $\mathfrak{p}_{139}$, and with it $\mathfrak{p}_{139}^2$, is not principal.
- $599 = 24^2 + 23$: the two prime ideals of F above 599 are generated by $24 \pm \sqrt{-23}$, hence principal, and $\mathrm{N}_{L/F}(P_{599})$ is principal.

The classes $[P_{139}]$ and $[P_{599}]$ are therefore not conjugate under $\mathrm{Gal}(L/\mathbb{Q})$, without using $h^-(115)$ or the exact order of the classes. With Theorem 3.3, which applies since both lattices have minimum 8, we obtain $X_{139} \ncong X_{599}$. [proved], without GRH, for the non-conjugacy; the non-isometry also requires the minimum 8 of one of the two lattices, which is computer-assisted (Theorem 6.11)

Theorem 6.11 (two extremal lattices; computer-assisted proof)**.** *The explicit lattices X_{139} and X_{599} are even, unimodular, of rank* 88 *and of minimum* 8*; they are therefore extremal. They are isometric neither to each other, nor to L_0, nor to L_S.* [minimum: computer-assisted proof, in the execution model of §5.4 (M1, M2, H1–H3); non-conjugacy of the classes: proved without GRH; non-isometry to each other: under this computer-assisted minimum; to L_0 and L_S: without recourse to this minimum (Proposition 6.9)]

Proof. By points 1 and 2 of Propositions 6.3 and 6.4, each of the two lattices is even, unimodular, positive definite, of rank 88, and contains a vector of norm 8. Let $z \neq 0$ be of norm at most 6. By Theorem 4.6, after multiplication by a root of unity and choice

of sign, one of the two projections is non-zero and of energy at most 1057/200, the total energy being at most 12; the case of a zero projection is handled by the dedicated lemma. By the geometric link, this pair belongs to the set $\mathcal{T}$ of one side, computed on the input files actually executed. By Corollary 5.2, applied to the four fresh executions of the second computer (two lattices, two sides), $\mathcal{T} = \emptyset$: a contradiction. The minimum is therefore at least 8, since the lattice is even, and it equals $8 = 2\lfloor 88/24 \rfloor + 2$. The non-isometry is Proposition 6.9, together with Remark 6.10. □

Novelty. The non-isometry to the two lattices of [Sak26] is proved. The absence of any other earlier example is a bibliographic conclusion, dated October 13, 2026 (§1.4): to our knowledge, X_{139} and X_{599} are two new examples of extremal lattices of rank 88.
Status. This proof is computer-assisted. It relies on the execution model made explicit in §5.4: the virtual machine and the arithmetic follow their specification (M1, M2, H1), no hardware fault occurred (H2), and the classes, inputs, outputs and preserved observations are authentic (H3). These premises are not an arithmetic hypothesis; the replications support them without proving them. The proof uses neither H4, nor the exactness of the counters, nor GRH. It consists of three parts, each reviewed by a second assistant on the preserved material:

(A) *the geometric link* of each lattice to the input files of the program, by exact checks, including the surjectivity of the coordinates and the exhaustiveness of the fibers by equality of determinants;
(B) *the link between the proof and the code*: complete selection, completion of the tasks and summary of a fresh launch; absence of hit files, established by the complete inventory of the output directories taken after the end of the computations, and by reading the launch scripts, which delete nothing;
(C) *the composition* above.

No human specialist has reviewed these proofs, and no independent implementation of the sweep has been run; these are possible consolidations, not logical conditions. The statement concerns the two explicit lattices of the note; it says nothing about a classification, the exact number of extremal orbits, the automorphism group, or the exactness of the counters of Theorem 5.1.

7. Open questions, methods and declarations

7.1. **Open questions.**

(Q1) *Consolidate Theorem 6.11.* A review of the non-omission proof by a specialist, an independent implementation of the sweep, or a lower bound for the minimum that does not depend on any execution.
(Q2) *Number of orbits of minimum* 8. How many of the 508 463 orbits of S consist of lattices of minimum 8? This number is at least 4: the orbits of L_0 and of L_S, whose minimum 8 is that of [Sak26], and those of X_{139} and of X_{599}, whose minimum 8 is Theorem 6.11 (computer-assisted proof). These four orbits are pairwise distinct: the classes $[P_{139}]$ and $[P_{599}]$ are not conjugate to each other (Remark 6.10), and their cubes are not principal, whereas those of the classes of L_0 and of L_S are (Proposition 6.9). A probabilistic model (Siegel mean and Poisson law on the orbits of short vectors under μ_L) suggests a proportion of the order of 4% of the classes. This model is a *heuristic*: it is not proved. It has been compared with only two cases in rank 56, and it underestimates one of the two.
(Q3) *The partial survivors.* What is the minimum of X_{3359}, X_{4049}, X_{4969} and X_{5659}? Nothing is known.
(Q4) *Sakamoto's second lattice.* The criterion of Theorem 4.6 has not been run on L_S. A prime ideal above 47, of degree 4 and norm 47^4, lies in the orbit of L_S (certified

3-component); for this ideal, the estimated cost is of the order of ten hours per side. The minimum 8 of L_S therefore rests, for us, on [Sak26] alone.

(Q5) *Orbit labels without GRH.* The exact localization of the classes in $\mathrm{Cl}(K_{22})$ and $\mathrm{Cl}(K_{44})$ is under GRH. A certification of the class group of K_{22}, of degree 22, would remove GRH for the 331-component; it has not been carried out.

(Q6) *Invariants.* The full automorphism groups of X_{139} and of X_{599} have not been determined; their perfection has not been studied.

(Q7) *Other conductors.* The cases $m = 184$ and $m = 276$, for which unimodular ideal lattices are even, and the case $m = 89$, are not treated.

7.2. **Methods and status of the computations.** The computations were done with PARI/GP, version 2.15.4 [PARI], with Python programs in exact integer and rational arithmetic, and with Java programs for the enumerations (OpenJDK 11.0.16.1 under Windows; OpenJDK 21 under Linux for the replications). Four types of computation are involved.

(C1) *Exact identities.* Relative class number, norms of generalized Bernoulli numbers, Gram matrices, isometry certificate $H \cong L_0$, witnesses of norm 6 and of norm 8, geometric links, class certificates in K_{22}. These computations are replayable; the verifiers use no class group computation. They give the status "proved".

(C2) *Class groups by* `bnfinit`. Localization of the classes in $\mathrm{Cl}(K_{22})$ and $\mathrm{Cl}(K_{44})$. Status: "under GRH". The 3-component is read in $\mathrm{Cl}(\mathbb{Q}(\sqrt{-23}))$, which is certified.

(C3) *Certified enumeration.* Program `Rang88CertZ` (§5). Status: "computer-assisted", under H1–H3. A hit, on the other hand, is an exact witness, replayable on its own.

(C4) *Non-certifying searches.* LLL reduction, `qfminim` of PARI/GP without options, capped filter, floating-point Java enumerator, time-bounded filter. Status: "observed". A vector found by these means becomes an exact witness once rechecked; an absence of vectors proves nothing.

Sakamoto's data. The Gram matrix of L_0 used in §2.5 comes from the public data repository that accompanies [Sak26], `https://github.com/RyotaroSakamoto0113/rank-88-extremal-lattice-computations`, revision `628ce2a7`, file `data/L0/integer_input.txt` (SHA-256 `68af477c...631f`). It is not redistributed with our data: these give its address and its hash, and the verification script reads it at that location. The files were read as data, without executing the code they contain. The article [Sak26] was read in its HTML version; the transcribed constants are cross-checked by exact identities. Its proofs have not been reviewed for this note.

Reviews. The following points were reviewed by the second assistant, Astra (OpenAI), each time within a limited scope: the computation of $h^-(115)$, by an independent computation in integers and rationals; the identification of the characters, the orbit count, the rational comparisons and the group arguments of §3, the gluing and the proof of Lemma 4.5; the two matrix identities that give $H \cong L_0$; the witness of P_{1471}; the geometric links of P_{139} and of P_{599}; the mechanism Z1–Z8 of the non-omission proof and its link to the code; the class certificates of P_{139} and of P_{599}; the composition of Theorem 6.11. The short proof of Remark 6.10 is due to Astra.

Limitations.

- P_{139} and P_{599} each rest on two complete executions of the same program, on two machines, with identical receipts; for P_{599}, the replay of the 21 459 tasks resumed from the first launch is added. No independent implementation has been run.
- The non-omission proof was written by the author of the code and checked once, with a limited favorable opinion.
- The review status of the class certificates is specified in Remark 6.8.
- Theorem 3.3 is an internal proof, audited once.

- The minimum 8 of L_0 and of L_S is that of a preprint. We have not re-proved it.
- The prior-art search is partial (§1.4).

The programs, the input data, the receipts, the certificates and the non-omission proof are in a public repository that accompanies the note: `https://github.com/Katabolanga/rang88-reseaux-ideaux`.

Author's declaration and use of artificial intelligence. The author publishes this note mainly to make known old ideas that are the author's own, as a human being. For the mathematical research and for proving these intuitions, the author was helped by two assistants: Claude Opus 5.5 (Anthropic) and Astra (OpenAI). They are not authors. Their use is as follows.

- *Writing.* The text was written with Claude Opus 5.5.
- *Translation.* The original text of this note is in French; it follows this English version in the same document. This English translation was prepared with Claude Opus 5.5.
- *Proofs.* The proofs of §§2–4, the non-omission proof summarized in §5 and the proofs of §6.4 were developed or put into shape with Claude Opus 5.5. Astra reviewed parts of them, listed in §7.2, and corrected several points, among them the value of the Blichfeldt bound in rank 22 and the justification of the threshold 1057/200; the short proof of Remark 6.10 is due to Astra.
- *Programs and computations.* The PARI/GP, Python and Java programs were written with Claude Opus 5.5. The first certifying program, based on intervals, was corrected and proved with Astra from a draft by Claude, and the program `Rang88CertZ` reuses parts of it. Astra's independent checks use programs written on Astra's side.
- *Bibliography.* The reading of [Sak26] and the prior-art search were done with both assistants, on public copies. The article [Que81] was read in November 2026, on the images of the journal, with Claude Opus 5.5.

The author takes scientific responsibility for the entire content. This note has not been reviewed by a human specialist.

References

[BN98] C. Bachoc, G. Nebe, *Extremal lattices of minimum* 8 *related to the Mathieu group* M_{22}, J. reine angew. Math. **494** (1998), 155–171. (Cited from [Sak26].)

[Bay84] E. Bayer-Fluckiger, *Definite unimodular lattices having an automorphism of given characteristic polynomial*, Comment. Math. Helv. **59** (1984), 509–538.

[Bay99] E. Bayer-Fluckiger, *Lattices and number fields*, in: Contemp. Math. **241**, Amer. Math. Soc., 1999, 69–84.

[Bli29] H. F. Blichfeldt, *The minimum value of quadratic forms, and the closest packing of spheres*, Math. Ann. **101** (1929), 605–608.

[CS99] J. H. Conway, N. J. A. Sloane, *Sphere packings, lattices and groups*, 3rd ed., Grundlehren der math. Wiss. **290**, Springer, 1999.

[FP85] U. Fincke, M. Pohst, *Improved methods for calculating vectors of short length in a lattice, including a complexity analysis*, Math. Comp. **44** (1985), 463–471.

[Mil14] J. C. Miller, *Class numbers of real cyclotomic fields of composite conductor*, LMS J. Comput. Math. **17** (2014), suppl. A, 404–417, DOI 10.1112/S1461157014000382.

[Neb13] G. Nebe, *On automorphisms of extremal even unimodular lattices*, Int. J. Number Theory **9** (2013), no. 8, 1933–1959; preprint arXiv 1212.0865.

[Neb16] G. Nebe, *Automorphisms of extremal unimodular lattices in dimension* 72, J. Number Theory **161** (2016), 362–383; preprint arXiv 1409.8473.

[PARI] The PARI Group, *PARI/GP version 2.15.4*, Univ. Bordeaux, 2023, `https://pari.math.u-bordeaux.fr/`.

[Que81] H.-G. Quebbemann, *Zur Klassifikation unimodularer Gitter mit Isometrie von Primzahlordnung*, J. reine angew. Math. **326** (1981), 158–170. (Cited since the revision of November 7, 2026.)

[Sak26] R. Sakamoto, *Two extremal even unimodular lattices of rank* 88, preprint arXiv 2609.15633, version 1 of September 14, 2026, version 2 of September 27, 2026. We cite version 2.

[Was97] L. C. Washington, *Introduction to cyclotomic fields*, 2nd ed., Graduate Texts in Math. **83**, Springer, 1997 (Theorem 4.17: analytic formula for the relative class number; Theorem 6.10: Stickelberger's theorem).

[Wat12] M. Watkins, *Another* 80*-dimensional extremal lattice*, J. Théor. Nombres Bordeaux **24** (2012), no. 1, 237–255, DOI 10.5802/jtnb.795. (Cited from [Sak26].)